

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number  
**WO 01/12600 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 209/04**,  
215/00, 235/04, 241/36, 265/36, 403/02, 498/02

(21) International Application Number: PCT/US00/21742

(22) International Filing Date: 10 August 2000 (10.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/148,627 12 August 1999 (12.08.1999) US  
60/202,202 5 May 2000 (05.05.2000) US

(71) Applicant (*for all designated States except US*): **COR THERAPEUTICS, INC.** [US/US]; 256 East Grand Avenue, South San Francisco, CA 94080 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ZHU, Bing-Yan** [CA/US]; 3325 Adelaide Way, Belmont, CA 94002 (US). **SCARBOROUGH, Robert, M.** [US/US]; 22 Greenbrier Court, Half Moon Bay, CA 94019 (US). **CLIZBE, Lane** [US/US]; 392 Newcastle Drive, Redwood City, CA 94061 (US). **DOUGHAN, Brandon** [US/US]; 650 Guerrero Street, San Francisco, CA 94110 (US). **JIA, Zhaozhong-Jon** [CN/US]; 849 West Orange Avenue #2007, San Francisco, CA 94110 (US). **KANE-MAGUIRE, Kim** [US/US]; 3425 Lodge Drive, Belmont, CA 94002 (US). **MARLOWE, Charles** [US/US]; 636 California Way, Redwood City, CA 94080 (US). **SONG, Yonghong** [CA/US]; 1144 Nimitz Lane,

Foster City, CA 94404 (US). **SU, Ting** [US/US]; 3222 Upper Lock Avenue, Belmont, CA 94002 (US). **TENG, Willy** [US/US]; 2355 Scott Street #103, San Francisco, CA 94115 (US). **ZHANG, Penglie** [CN/US]; 224 Serrano Drive, San Francisco, CA 94132 (US).

(74) Agent: **WEIMAR, Elizabeth, C.**; Morgan, Lewis & Bockius LLP, 1800 M Street NW, Washington, DC 20036 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- *With international search report.*
- *Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.*

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: INHIBITORS OF FACTOR Xa



(57) Abstract: Novel compounds, their salts and compositions related thereto having activity against mammalian factor Xa are disclosed. The compounds of formula (I) are useful *in vitro* or *in vivo* for preventing or treating coagulation disorders.



WO 01/12600 A1

## INHIBITORS OF FACTOR Xa

### Related Applications

This application claims benefit of priority under 35 USC § 119(e) to U.S. Provisional Application No. 60/202,202 filed on May 5, 2000 and U.S. Provisional Application No. 60/148,627 filed on August 12, 1999, which are both herein incorporated in their entirety by reference.

### Field of the Invention

This invention relates to novel compounds which are potent and highly selective inhibitors of isolated factor Xa or factor Xa when assembled in the prothrombinase complex. These compounds show selectivity for factor Xa versus other proteases of the coagulation (e.g. thrombin, fVIIa, fIXa) or the fibrinolytic cascades (e.g. plasminogen activators, plasmin). In another aspect, the present invention relates to novel monoamidino-containing compounds, their pharmaceutically acceptable salts, and pharmaceutically acceptable compositions thereof which are useful as potent and specific inhibitors of blood coagulation in mammals. In yet another aspect, the invention relates to methods for using these inhibitors as therapeutic agents for disease states in mammals characterized by coagulation disorders.

### Background of the Invention

Hemostasis, the control of bleeding, occurs by surgical means, or by the physiological properties of vasoconstriction and coagulation. This invention is particularly concerned with blood coagulation and ways in which it assists in maintaining the integrity of mammalian circulation after injury, inflammation, disease, congenital defect, dysfunction or other disruption. Although platelets and blood coagulation are both involved in thrombus formation, certain components of the coagulation cascade are primarily responsible for the amplification or acceleration of the processes involved in platelet aggregation and fibrin deposition.

Thrombin is a key enzyme in the coagulation cascade as well as in hemostasis. Thrombin plays a central role in thrombosis through its ability to catalyze the conversion of fibrinogen into fibrin and through its potent platelet activation activity. Direct or indirect



inhibition of thrombin activity has been the focus of a variety of recent anticoagulant strategies as reviewed by Claeson, G., "Synthetic Peptides and Peptidomimetics as Substrates and Inhibitors of Thrombin and Other Proteases in the Blood Coagulation System", *Blood Coag. Fibrinol.* 5, 411-436 (1994). Several classes of anticoagulants currently used in the clinic directly or indirectly affect thrombin (i.e. heparins, low-molecular weight heparins, heparin-like compounds and coumarins).

A prothrombinase complex, including Factor Xa (a serine protease, the activated form of its Factor X precursor and a member of the calcium ion binding, gamma carboxyglutamyl (Gla)-containing, vitamin K dependent, blood coagulation glycoprotein family), converts the zymogen prothrombin into the active procoagulant thrombin. Unlike thrombin, which acts on a variety of protein substrates as well as at a specific receptor, factor Xa appears to have a single physiologic substrate, namely prothrombin. Since one molecule of factor Xa may be able to generate up to 138 molecules of thrombin (Elodi et al., *Thromb. Res.* 15, 617-619 (1979)), direct inhibition of factor Xa as a way of indirectly inhibiting the formation of thrombin may be an efficient anticoagulant strategy. Therefore, it has been suggested that compounds which selectively inhibit factor Xa may be useful as *in vitro* diagnostic agents, or for therapeutic administration in certain thrombotic disorders, see e.g., WO 94/13693.

Polypeptides derived from hematophagous organisms have been reported which are highly potent and specific inhibitors of factor Xa. United States Patent 4,588,587 describes anticoagulant activity in the saliva of the Mexican leech, *Haementeria officinalis*. A principal component of this saliva was shown to be the polypeptide factor Xa inhibitor, antistasin (ATS), by Nutt, E. *et al.*, "The Amino Acid Sequence of Antistasin, a Potent Inhibitor of Factor Xa Reveals a Repeated Internal Structure", *J. Biol. Chem.*, 263, 10162-10167 (1988). Another potent and highly specific inhibitor of Factor Xa, called tick anticoagulant peptide (TAP), has been isolated from the whole body extract of the soft tick *Ornithodoros moubata*, as reported by Waxman, L., *et al.*, "Tick Anticoagulant Peptide (TAP) is a Novel Inhibitor of Blood Coagulation Factor Xa" *Science*, 248, 593-596 (1990).

Factor Xa inhibitory compounds which are not large polypeptide-type inhibitors have also been reported including: Tidwell, R.R. *et al.*, "Strategies for Anticoagulation With Synthetic Protease Inhibitors. Xa Inhibitors Versus Thrombin Inhibitors", *Thromb. Res.*, 19, 339-349 (1980); Turner, A.D. *et al.*, "p-Amidino Esters as Irreversible Inhibitors

of Factor IXa and Xa and Thrombin", *Biochemistry*, 25, 4929-4935 (1986); Hitomi, Y. *et al.*, "Inhibitory Effect of New Synthetic Protease Inhibitor (FUT-175) on the Coagulation System", *Haemostasis*, 15, 164-168 (1985); Sturzebecher, J. *et al.*, "Synthetic Inhibitors of Bovine Factor Xa and Thrombin. Comparison of Their Anticoagulant Efficiency", *Thromb. Res.*, 54, 245-252 (1989); Kam, C.M. *et al.*, "Mechanism Based Isocoumarin Inhibitors for Trypsin and Blood Coagulation Serine Proteases: New Anticoagulants", *Biochemistry*, 27, 2547-2557 (1988); Hauptmann, J. *et al.*, "Comparison of the Anticoagulant and Antithrombotic Effects of Synthetic Thrombin and Factor Xa Inhibitors", *Thromb. Haemost.*, 63, 220-223 (1990); and the like.

Others have reported Factor Xa inhibitors which are small molecule organic compounds, such as nitrogen containing heterocyclic compounds which have amidino substituent groups, wherein two functional groups of the compounds can bind to Factor Xa at two of its active sites. For example, WO 98/28269 describes pyrazole compounds having a terminal C(=NH)-NH<sub>2</sub> group; WO 97/21437 describes benzimidazole compounds substituted by a basic radical which are connected to a naphthyl group via a straight or branched chain alkylene, -C(=O) or -S(=O)<sub>2</sub> bridging group; WO 99/10316 describes compounds having a 4-phenyl-N-alkylamidino-piperidine and 4-phenoxy-N-alkylamidino-piperidine group connected to a 3-amidinophenyl group via a carboxamidealkyleneamino bridge; and EP 798295 describes compounds having a 4-phenoxy-N-alkylamidino-piperidine group connected to an amidinonaphthyl group via a substituted or unsubstituted sulfonamide or carboxamide bridging group.


There exists a need for effective therapeutic agents for the regulation of hemostasis, and for the prevention and treatment of thrombus formation and other pathological processes in the vasculature induced by thrombin such as restenosis and inflammation. In particular, there continues to be a need for compounds which selectively inhibit factor Xa or its precursors. Compounds that have different combinations of bridging groups and functional groups than compounds previously discovered are needed, particularly compounds which selectively or preferentially bind to Factor Xa. Compounds with a higher degree of binding to Factor Xa than to thrombin are desired, especially those compounds having good bioavailability and/or solubility.

### Summary of the Invention

The present invention relates to novel compounds which inhibit factor Xa, their pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives, and pharmaceutically acceptable compositions thereof which have particular biological properties and are useful as potent and specific inhibitors of blood coagulation in mammals. Pharmaceutical compositions of the invention may be used to prevent or treat a condition in a mammal characterized by undesired thrombosis. In another aspect, the invention relates to methods of using these inhibitors as diagnostic reagents or as therapeutic agents for disease states in mammals which have coagulation disorders, such as in the treatment or prevention of a condition in a mammal characterized by undesired thrombosis such as, for example, any thrombotically mediated acute coronary or cerebrovascular syndrome, any thrombotic syndrome occurring in the venous system, any coagulopathy, and any thrombotic complications associated with extracorporeal circulation or instrumentation, and for the inhibition of coagulation of biological samples and blood in biological samples.

In certain embodiments, this invention relates to novel compounds which are potent and highly selective inhibitors of isolated factor Xa or factor Xa when assembled in the prothrombinase complex. These compounds show selectivity for factor Xa versus other proteases of the coagulation cascade (e.g. thrombin, etc.) or the fibrinolytic cascade, and are useful as diagnostic reagents as well as antithrombotic agents.

In a preferred embodiment, the present invention provides a compound of the formula I:



wherein:

A is selected from:

- (a) C<sub>1</sub>-C<sub>6</sub>-alkyl;
- (b) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;
- (c) -N(-R<sup>2</sup>, -R<sup>3</sup>), R<sup>3</sup>-C(=N-R<sup>2</sup>)-, (-R<sup>2</sup>, -R<sup>3</sup>)N-C(=N-R<sup>2</sup>)-, (-R<sup>2</sup>, -R<sup>3</sup>)N-C(=N-R<sup>2</sup>)-N(-R-)-

- (d) phenyl, which is independently substituted with 0-2  $R^1$  substituents;
- (e) naphthyl, which is independently substituted with 0-2  $R^1$  substituents; and
- (f) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^1$  substituents;

$R^1$  is selected from:

Halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN, -NO<sub>2</sub>,  $R^2-C(=N-R^3)-$ ,  $(-R^2, -R^3)N-C(=N-R^2)-$ ,  $-(CH_2)_mNR^2R^3$ ,  $-C(=O)-N(-R^2, -R^3)$ ,  $-SO_2N(-R^2, -R^3)$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_1-C_4$ -alkyl, -CN  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl and -NO<sub>2</sub>;

$R^2$  and  $R^3$  are independently selected from the group consisting of:

H,  $-OR^2$ ,  $-N(-R^2, -R^3)$ ,  $-C_{1-4}$ alkyl,  $-C_{2-6}$ alkenyl,  $-C_{2-6}$ alkynyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkylphenyl and  $-C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $-C_{1-4}$ alkyl,  $-C_{2-6}$ alkenyl,  $-C_{2-6}$ alkynyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN, and -NO<sub>2</sub>;

or  $R^2$  and  $R^3$  taken together can form a 3-8 membered cycloalkyl or a heterocyclic ring system, wherein the heterocyclic ring system may have from 5 to 10 ring atoms, with 1 to 2 rings being in the ring system and contain from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the heterocyclic ring system may be independently replaced with a member selected from the group consisting of halo,  $C_1-C_4$ -alkyl, -CN  $-C_{1-4}$ alkyl,  $-C_{2-6}$ alkenyl,  $-C_{2-6}$ alkynyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl and -NO<sub>2</sub>;

m is an integer of 0-2;

Q is a member selected from the group consisting of:

a direct link,  $-CH_2-$ ,  $-C(=O)-$ ,  $-N(R^4)-$ ,  $-N(R^4)CH_2-$ ,  $-C=N(R^4)-$ ,  $-C(=O)-N(R^4)-$ ,  $-N(R^4)-C(=O)-$ ,  $-SO_2-$ ,  $-O-$ ,  $-SO_2-N(R^4)-$  and  $-N(R^4)-SO_2-$ ;



R<sup>4</sup> is selected from:

H, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylphenyl and -C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

D is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1a</sup> substituents; and
- (b) an aromatic six-membered heterocyclic ring having from 1-2 ring nitrogen atoms, and wherein the ring atoms may be substituted with 0-2 R<sup>1a</sup> substituents;

R<sup>1a</sup> is selected from:

Halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2a</sup>R<sup>3a</sup>, SO<sub>2</sub>NR<sup>2a</sup>R<sup>3a</sup>, SO<sub>2</sub>R<sup>2a</sup>, CF<sub>3</sub>, OR<sup>2a</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>2a</sup> and R<sup>3a</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

M is a member selected from the group consisting of:

-N(R<sup>16</sup>)-C(=O)-, -N(R<sup>16</sup>)-C(=S)-, -C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-, -C(-R<sup>19</sup>, -R<sup>19a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-, -C(-R<sup>17</sup>)-C(-R<sup>18</sup>)-C(=O)-, -C(-R<sup>17</sup>)-C(-R<sup>18</sup>)-C(=S)-, -C(-R<sup>17</sup>)-C(-R<sup>18</sup>)-, -O-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -O-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S(=O)<sub>2</sub>-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S(=O)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S(=O)<sub>2</sub>-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S(=O)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(=O)-C(=O)-, -N(R<sup>16</sup>)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -N(R<sup>16</sup>)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-,

-C(=S)-C(=S)-, -C(=S)-C(=O)-, -C(=O)-C(=S)-, -N=C(-R<sup>17</sup>)-C(=O)-, -N=C(-R<sup>17</sup>)-C(=S)-, -C(-R<sup>17</sup>)=N-, -N(-R<sup>16</sup>)-C(=O)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -O-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -N(-R<sup>16</sup>)-C(=S)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -S(=O)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -S(=O)<sub>2</sub>-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -C(=C(R<sup>17b</sup>, -R<sup>17c</sup>))-C(=O)-, -C(=C(R<sup>17b</sup>, -R<sup>17c</sup>))-C(=S)-, -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(=S)-, -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-(N(-H, -R<sup>18b</sup>)), -R<sup>17a</sup>)-C(=O)-; -N=C(-R<sup>17</sup>)- and -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-(N(-H, -R<sup>18b</sup>)), -R<sup>17a</sup>)-C(=S)-; wherein the first named atom of the chain is directly attached to D, and wherein D, M and the N atom attached to the last chain atom of M collectively form a bicyclic ring structure;

R<sup>16</sup>, R<sup>17</sup>, R<sup>17a</sup>, R<sup>18</sup>, R<sup>18a</sup>, R<sup>18b</sup>, R<sup>19</sup>, and R<sup>19a</sup> are each independently selected from the group consisting of:

hydrogen, halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>R<sup>2</sup>, CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, -CN, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

R<sup>17b</sup> and R<sup>17c</sup> are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy, -C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the cycloalkyl, the phenyl ring, or the aromatic heterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, -CN, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

E is a member selected from the group consisting of:

a direct link, -C(=O)-, -C(=O)-N(R<sup>5</sup>)-, -C(-R<sup>5a</sup>, -R<sup>6a</sup>)- and -C(-R<sup>5b</sup>, -R<sup>6b</sup>)-C(-R<sup>5c</sup>, -R<sup>6c</sup>)-;

wherein R<sup>5</sup>, R<sup>5a</sup>, R<sup>6a</sup>, R<sup>5b</sup>, R<sup>6b</sup>, R<sup>5c</sup> and R<sup>6c</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylheteroaryl, C<sub>1-4</sub>alkylCOOH and

C<sub>1-4</sub>alkylCOOC<sub>1-4</sub>alkyl, wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -OH, -O-C<sub>1-4</sub>alkyl, -SH, -S-C<sub>1-4</sub>alkyl, -CN and -NO<sub>2</sub>;

G is selected from:

a direct link, -C(R<sup>7</sup>,R<sup>8</sup>)-, -C(R<sup>7a</sup>,R<sup>8a</sup>)-C(R<sup>7b</sup>,R<sup>8b</sup>)- and -C(R<sup>7c</sup>)=C(R<sup>8c</sup>)-;

wherein R<sup>7</sup>, R<sup>8</sup>, R<sup>7a</sup>, R<sup>8a</sup>, R<sup>7b</sup>, R<sup>8b</sup>, R<sup>7c</sup> and R<sup>8c</sup> are independently a member selected from from the group consisting of:

hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)OR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)-<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, a naturally occurring or synthetic amino acid side chain, and C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the C<sub>0-4</sub>alkylheterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>9</sup> and R<sup>10</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>3-8</sub>cycloalkyl, and C<sub>1-4</sub>alkyl-O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl-COOH wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>, and wherein R<sup>9</sup> and R<sup>10</sup> taken together can form a 5-8 membered heterocyclic ring;

J is a member selected from the group consisting of:

a direct link, -O-, -O-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -S-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)<sub>2</sub>-(-R<sup>11</sup>, -R<sup>11a</sup>)-, -C(=O)-, -C(=O)-N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(=O)-, -N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(-R<sup>11</sup>, -R<sup>11a</sup>)- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>11</sup> substituents;

R<sup>11</sup>, R<sup>11a</sup>, R<sup>11b</sup>, and R<sup>11</sup> are a member independently selected from the group consisting of:

hydrogen, halo, -CF<sub>3</sub>, -CN, -NR<sup>9</sup>R<sup>10</sup>, -SO<sub>2</sub>Me, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>2-6</sub>alkenyl, -O-C<sub>2-6</sub>alkynyl, -O-C<sub>3-8</sub>cycloalkyl, -O-C<sub>1-4</sub>alkyl-O-C<sub>1-4</sub>alkyl, -O-C<sub>1-4</sub>alkyl-COOH, -O-C<sub>1-4</sub>alkyl-phenyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>2-6</sub>alkenyl, -C(=O)-O-C<sub>2-6</sub>alkynyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, C<sub>0-4</sub>alkylC(=O)OR<sup>9</sup>, C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl; wherein from 1-4 hydrogen atoms on the C<sub>0-4</sub>alkylheterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-phenyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents;
- (b) naphthyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents; and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1b</sup> substituents;

R<sup>1b</sup> is a member selected from the group consisting of:



halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>R<sup>2b</sup>, CF<sub>3</sub>, OR<sup>2b</sup>, O-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, O-CH<sub>2</sub>-COOR<sup>2b</sup>, N(R<sup>2b</sup>)-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, N(-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>)<sub>2</sub>, N(R<sup>2b</sup>)-C(=O)R<sup>3b</sup>, N(R<sup>2b</sup>)-SO<sub>2</sub>-R<sup>3b</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-phenyl, -CN and -NO<sub>2</sub>;

R<sup>2b</sup> and R<sup>3b</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

L is selected from:

H, -CN, C(=O)NR<sup>12</sup>R<sup>13</sup>, -(CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup>, C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, OR<sup>12</sup>, -NR<sup>12</sup>C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, and NR<sup>12</sup>C(=NR<sup>12</sup>)-R<sup>13</sup>;

n is an integer from 0 to 8;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

hydrogen, -OR<sup>14</sup>, -NR<sup>14</sup>R<sup>15</sup>, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, COOC<sub>1-4</sub>alkyl, COO-C<sub>0-4</sub>alkylphenyl and COO-C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH, -O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

R<sup>14</sup> and R<sup>15</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

In certain aspects of this invention, compounds are provided which are useful as diagnostic reagents. In another aspect, the present invention includes pharmaceutical compositions comprising a pharmaceutically effective amount of the compounds of this invention and a pharmaceutically acceptable carrier. In yet another aspect, the present invention includes methods comprising using the above compounds and pharmaceutical compositions for preventing or treating disease states characterized by disorders of the blood coagulation process in mammals, or for preventing coagulation in stored blood products and samples. Optionally, the methods of this invention comprise administering the pharmaceutical composition in combination with an additional therapeutic agent such as an antithrombotic and/or a thrombolytic agent and/or an anticoagulant.

The preferred compounds also include their pharmaceutically acceptable isomers, hydrates, solvates, salts and prodrug derivatives.

## Detailed Description of the Invention

### Definitions

In accordance with the present invention and as used herein, the following terms are defined with the following meanings, unless explicitly stated otherwise.

The term "alkenyl" refers to a trivalent straight chain or branched chain unsaturated aliphatic radical. The term "alkinyl" (or "alkynyl") refers to a straight or branched chain aliphatic radical that includes at least two carbons joined by a triple bond. If no number of carbons is specified alkenyl and alkinyl each refer to radicals having from 2-12 carbon atoms.

The term "alkyl" refers to saturated aliphatic groups including straight-chain, branched-chain and cyclic groups having the number of carbon atoms specified, or if no number is specified, having up to 12 carbon atoms. The term "cycloalkyl" as used herein refers to a mono-, bi-, or tricyclic aliphatic ring having 3 to 14 carbon atoms and preferably 3 to 7 carbon atoms.

As used herein, the terms "carbocyclic ring structure" and "C<sub>3-16</sub> carbocyclic mono, bicyclic or tricyclic ring structure" or the like are each intended to mean stable ring structures having only carbon atoms as ring atoms wherein the ring structure is a substituted or unsubstituted member selected from the group consisting of: a stable monocyclic ring which is aromatic ring ("aryl") having six ring atoms; a stable monocyclic

non-aromatic ring having from 3 to 7 ring atoms in the ring; a stable bicyclic ring structure having a total of from 7 to 12 ring atoms in the two rings wherein the bicyclic ring structure is selected from the group consisting of ring structures in which both of the rings are aromatic, ring structures in which one of the rings is aromatic and ring structures in which both of the rings are non-aromatic; and a stable tricyclic ring structure having a total of from 10 to 16 atoms in the three rings wherein the tricyclic ring structure is selected from the group consisting of: ring structures in which three of the rings are aromatic, ring structures in which two of the rings are aromatic and ring structures in which three of the rings are non-aromatic. In each case, the non-aromatic rings when present in the monocyclic, bicyclic or tricyclic ring structure may independently be saturated, partially saturated or fully saturated. Examples of such carbocyclic ring structures include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin). Moreover, the ring structures described herein may be attached to one or more indicated pendant groups via any carbon atom which results in a stable structure. The term "substituted" as used in conjunction with carbocyclic ring structures means that hydrogen atoms attached to the ring carbon atoms of ring structures described herein may be substituted by one or more of the substituents indicated for that structure if such substitution(s) would result in a stable compound.

The term "aryl" which is included with the term "carbocyclic ring structure" refers to an unsubstituted or substituted aromatic ring, substituted with one, two or three substituents selected from loweralkoxy, loweralkyl, loweralkylamino, hydroxy, halogen, cyano, hydroxyl, mercapto, nitro, thioalkoxy, carboxaldehyde, carboxyl, carboalkoxy and carboxamide, including but not limited to carbocyclic aryl, heterocyclic aryl, and biaryl groups and the like, all of which may be optionally substituted. Preferred aryl groups include phenyl, halophenyl, loweralkylphenyl, naphthyl, biphenyl, phenanthrenyl and naphthacenyl.

The term "arylalkyl" which is included with the term "carbocyclic aryl" refers to one, two, or three aryl groups having the number of carbon atoms designated, appended to an alkyl group having the number of carbon atoms designated. Suitable arylalkyl groups

include, but are not limited to, benzyl, picolyl, naphthylmethyl, phenethyl, benzyhydryl, trityl, and the like, all of which may be optionally substituted.

As used herein, the term "heterocyclic ring" or "heterocyclic ring system" is intended to mean a substituted or unsubstituted member selected from the group consisting of stable monocyclic ring having from 5-7 members in the ring itself and having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S; a stable bicyclic ring structure having a total of from 7 to 12 atoms in the two rings wherein at least one of the two rings has from 1 to 4 hetero atoms selected from N, O and S, including bicyclic ring structures wherein any of the described stable monocyclic heterocyclic rings is fused to a hexane or benzene ring; and a stable tricyclic heterocyclic ring structure having a total of from 10 to 16 atoms in the three rings wherein at least one of the three rings has from 1 to 4 hetero atoms selected from the group consisting of N, O and S. Any nitrogen and sulfur atoms present in a heterocyclic ring of such a heterocyclic ring structure may be oxidized. Unless indicated otherwise the terms "heterocyclic ring" or "heterocyclic ring system" include aromatic rings, as well as non-aromatic rings which can be saturated, partially saturated or fully saturated non-aromatic rings. Also, unless indicated otherwise the term "heterocyclic ring system" includes ring structures wherein all of the rings contain at least one hetero atom as well as structures having less than all of the rings in the ring structure containing at least one hetero atom, for example bicyclic ring structures wherein one ring is a benzene ring and one of the rings has one or more hetero atoms are included within the term "heterocyclic ring systems" as well as bicyclic ring structures wherein each of the two rings has at least one hetero atom. Moreover, the ring structures described herein may be attached to one or more indicated pendant groups via any hetero atom or carbon atom which results in a stable structure. Further, the term "substituted" means that one or more of the hydrogen atoms on the ring carbon atom(s) or nitrogen atom(s) of the each of the rings in the ring structures described herein may be replaced by one or more of the indicated substituents if such replacement(s) would result in a stable compound. Nitrogen atoms in a ring structure may be quaternized, but such compounds are specifically indicated or are included within the term "a pharmaceutically acceptable salt" for a particular compound. When the total number of O and S atoms in a single heterocyclic ring is greater than 1, it is preferred that such atoms not be adjacent to one another. Preferably, there are no more than 1 O or S ring atoms in the same ring of a given heterocyclic ring structure.



Examples of monocyclic and bicyclic heterocyclic ring systems, in alphabetical order, are acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalinyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolyl, imidazolyl, 1H-indazolyl, indolinyl, indoliziny, indolyl, 3H-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl (benzimidazolyl), isothiazolyl, isoxazolyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolidinyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyroazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridoxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2H-pyrrolyl, pyrrolyl, quinazolinyl, quinolinyl, 4H-quinoliziny, quinoxalinyl, quinuclidinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl and xanthenyl. Preferred heterocyclic ring structures include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrrolidinyl, imidazolyl, indolyl, benzimidazolyl, 1H-indazolyl, oxazolyl, or isatinoyl. Also included are fused ring and spiro compounds containing, for example, the above heterocyclic ring structures.

As used herein the term "aromatic heterocyclic ring system" has essentially the same definition as for the monocyclic and bicyclic ring systems except that at least one ring of the ring system is an aromatic heterocyclic ring or the bicyclic ring has an aromatic or non-aromatic heterocyclic ring fused to an aromatic carbocyclic ring structure.

The terms "halo" or "halogen" as used herein refer to Cl, Br, F or I substituents. The term "haloalkyl", and the like, refer to an aliphatic carbon radicals having at least one hydrogen atom replaced by a Cl, Br, F or I atom, including mixtures of different halo atoms. Trihaloalkyl includes trifluoromethyl and the like as preferred radicals, for example.

The term "methylene" refers to  $-\text{CH}_2-$ . The term "Bu" refers to "butyl" or  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ ; the term "Ph" refers to "phenyl"; the term "Me" refers to "methyl" or

—CH<sub>3</sub>; the term “Et” refers to “ethyl” or —CH<sub>2</sub>CH<sub>3</sub>; the term “Bu(t)” or “t-Bu” refers to “*tert*-butyl” or —C(CH<sub>3</sub>)<sub>4</sub>.

The term “amino acid side chain” includes any naturally occurring or synthetic side chain, *i.e.* the side chain “R” of an amino acid having the formula: NH<sub>2</sub>-CHR-COOH. Synthetic amino acids include both (R) and (S) enantiomers, as well as derivatives of the naturally-occurring amino acid side chains. The side chain may be any synthetic amino acid side chain known in the art, including but not limited to, those produced by recombinant DNA techniques, fermentation, or by solid phase synthesis techniques. Such synthetic amino acid side chains may contain one or more functional groups selected from the following: alkyl, aryl, alkenyl, alkynyl, thiols, primary and secondary amines, aldehydes, carboxylates, nitriles, aromatic amines, aromatic carboxylates, primary alcohols, and the like.

The term "pharmaceutically acceptable salts" includes salts of compounds derived from the combination of a compound and an organic or inorganic acid. These compounds are useful in both free base and salt form. In practice, the use of the salt form amounts to use of the base form; both acid and base addition salts are within the scope of the present invention.

“Pharmaceutically acceptable acid addition salt” refers to salts retaining the biological effectiveness and properties of the free bases and which are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

“Pharmaceutically acceptable base addition salts” include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium and magnesium salts. Salts derived from pharmaceutically acceptable organic nontoxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic


amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-diethylaminoethanol, trimethamine, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like. Particularly preferred organic nontoxic bases are isopropylamine, diethylamine, ethanolamine, trimethamine, dicyclohexylamine, choline, and caffeine.

"Biological property" for the purposes herein means an *in vivo* effector or antigenic function or activity that is directly or indirectly performed by a compound of this invention that are often shown by *in vitro* assays. Effector functions include receptor or ligand binding, any enzyme activity or enzyme modulatory activity, any carrier binding activity, any hormonal activity, any activity in promoting or inhibiting adhesion of cells to an extracellular matrix or cell surface molecules, or any structural role. Antigenic functions include possession of an epitope or antigenic site that is capable of reacting with antibodies raised against it.

In the compounds of this invention, carbon atoms bonded to four non-identical substituents are asymmetric. Accordingly, the compounds may exist as diastereoisomers, enantiomers or mixtures thereof. The syntheses described herein may employ racemates, enantiomers or diastereomers as starting materials or intermediates. Diastereomeric products resulting from such syntheses may be separated by chromatographic or crystallization methods, or by other methods known in the art. Likewise, enantiomeric product mixtures may be separated using the same techniques or by other methods known in the art. Each of the asymmetric carbon atoms, when present in the compounds of this invention, may be in one of two configurations (R or S) and both are within the scope of the present invention.

#### Preferred Embodiments

In a preferred embodiment, the present invention provides a compound according to the formula I:



wherein:

A is selected from:

- (a) C<sub>1</sub>-C<sub>6</sub>-alkyl;
- (b) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;
- (c) phenyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;
- (d) naphthyl, which is independently substituted with 0-2 R<sup>1</sup> substituents; and
- (e) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from the group consisting of N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1</sup> substituents;

R<sup>1</sup> is selected from:

halo, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, -C<sub>1-4</sub>alkyl, -CN, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

R<sup>2</sup> and R<sup>3</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

m is an integer of 0-2;

Q is a member selected from the group consisting of:



a direct link,  $-C(=O)-$ ,  $-N(R^4)-$ ,  $-C(=O)-N(R^4)-$ ,  $-N(R^4)-C(=O)-$ ,  $-SO_2-$ ,  $-O-$ ,  $-SO_2-N(R^4)-$  and  $-N(R^4)-SO_2-$ ;

$R^4$  is selected from:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$ , and  $-NO_2$ ;

D is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1a}$  substituents; and
- (b) an aromatic six-membered heterocyclic ring having from 1-2 ring nitrogen atoms, and wherein the ring atoms may be substituted with 0-2  $R^{1a}$  substituents;


$R^{1a}$  is selected from:

halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$ ,  $-NO_2$ ,  $(CH_2)_mNR^{2a}R^{3a}$ ,  $SO_2NR^{2a}R^{3a}$ ,  $SO_2R^{2a}$ ,  $CF_3$ ,  $OR^{2a}$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$  and  $-NO_2$ ;

$R^{2a}$  and  $R^{3a}$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$  and  $-NO_2$ ;

M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



and the like, wherein the aromatic carbocyclic ring corresponding to the D portion for each of the bicyclic rings may be replaced with an aromatic heterocyclic ring as defined above for D, and wherein 0 to 2 of the hydrogen atoms on the D portion of the bicyclic ring may be replaced by  $R^{1a}$  substituents as defined above;

$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN, -NO<sub>2</sub>,  $(CH_2)_mNR^2R^3$ , SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>R<sup>2</sup>, CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl, -CN,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl and -NO<sub>2</sub>;

$R^{17b}$  and  $R^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy, -C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the cycloalkyl, the phenyl ring, or the aromatic heterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, -CN, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

E is a member selected from the group consisting of:

a direct link, -C(=O)-, -C(=O)-N(R<sup>5</sup>)-, -C(-R<sup>5a</sup>, -R<sup>6a</sup>)- and  
-(-C(-R<sup>5b</sup>, -R<sup>6b</sup>)-C(-R<sup>5c</sup>, -R<sup>6c</sup>)-;

wherein R<sup>5</sup>, R<sup>5a</sup>, R<sup>6a</sup>, R<sup>5b</sup>, R<sup>6b</sup>, R<sup>5c</sup> and R<sup>6c</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylheteroaryl, C<sub>1-4</sub>alkylCOOH and C<sub>1-4</sub>alkylCOOC<sub>1-4</sub>alkyl, wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -OH, -O-C<sub>1-4</sub>alkyl, -SH, -S-C<sub>1-4</sub>alkyl, -CN and -NO<sub>2</sub>;

G is selected from:

a direct link, -C(R<sup>7</sup>, R<sup>8</sup>)-, -C(R<sup>7a</sup>, R<sup>8a</sup>)-C(R<sup>7b</sup>, R<sup>8b</sup>)- and -C(R<sup>7c</sup>)=C(R<sup>8c</sup>)-;

wherein R<sup>7</sup>, R<sup>8</sup>, R<sup>7a</sup>, R<sup>8a</sup>, R<sup>7b</sup>, R<sup>8b</sup>, R<sup>7c</sup> and R<sup>8c</sup> are independently a member selected from from the group consisting of:

hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)-<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, and a naturally occurring or synthetic amino acid side chain, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

$R^9$  and  $R^{10}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl, -CN and -NO<sub>2</sub>, and wherein  $R^9$  and  $R^{10}$  taken together can form a 5-8 membered heterocyclic ring;

J is a member selected from the group consisting of:

a direct link, -O-, -O-C( $R^{11}$ ,  $R^{11a}$ )-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -S-C( $R^{11}$ ,  $R^{11a}$ )-, -S(=O)-C( $R^{11}$ ,  $R^{11a}$ )-, -S(=O)<sub>2</sub>-( $R^{11}$ ,  $R^{11a}$ )-, -C(=O)-, -C(=O)-N( $R^{11b}$ )-, -N( $R^{11b}$ )-C(=O)-, -N( $R^{11b}$ )-, -N( $R^{11b}$ )-C( $R^{11}$ ,  $R^{11a}$ )- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{11c}$  substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$  and  $R^{11c}$  are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O- $C_{1-4}$ alkyl, -O- $C_{2-6}$ alkenyl, -O- $C_{2-6}$ alkynyl, -O- $C_{3-8}$ cycloalkyl, -COOH, -C(=O)-O- $C_{1-4}$ alkyl, -C(=O)-O- $C_{2-6}$ alkenyl, -C(=O)-O- $C_{2-6}$ alkynyl, -C(=O)-O- $C_{3-8}$ cycloalkyl,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $C_{0-4}$ alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S,  $CH_2COOC_{1-4}$ alkyl,  $CH_2COOC_{1-4}$ alkylphenyl and  $CH_2COOC_{1-4}$ alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1b}$  substituents;
- (b) naphthyl, which is independently substituted with 0-2  $R^{1b}$  substituents; and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{1b}$  substituents;

$R^{1b}$  is a member selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl, -CN, -NO<sub>2</sub>,  $NR^{2b}R^{3b}$ ,  $SO_2NR^{2b}R^{3b}$ ,  $SO_2R^{2b}$ ,  $CF_3$ ,  $OR^{2b}$ ,  $O-CH_2-CH_2-OR^{2b}$ ,  $O-CH_2-COOR^{2b}$ ,  $N(R^{2b})-CH_2-CH_2-OR^{2b}$ ,  $N(-CH_2-CH_2-OR^{2b})_2$ ,  $N(R^{2b})-C(=O)R^{3b}$ ,  $N(R^{2b})-$

SO<sub>2</sub>-R<sup>3b</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>2b</sup> and R<sup>3b</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

L is selected from:

H, -CN, C(=O)NR<sup>12</sup>R<sup>13</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup>, C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, OR<sup>12</sup>, -NR<sup>12</sup>C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, and NR<sup>12</sup>C(=NR<sup>12</sup>)-R<sup>13</sup>;

n is an integer from 0 to 8;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

hydrogen, -OR<sup>14</sup>, -NR<sup>14</sup>R<sup>15</sup>, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, COOC<sub>1-4</sub>alkyl, COO-C<sub>0-4</sub>alkylphenyl and COO-C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH, -O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;


R<sup>14</sup> and R<sup>15</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.



In a further preferred embodiment, the present invention provides a compound according to the formula I:



wherein:

A is selected from:

- (a)  $C_1$ - $C_6$ -alkyl;
- (b)  $C_3$ - $C_8$ -cycloalkyl;
- (c) phenyl, which is independently substituted with 0-2  $R^1$  substituents;
- (d) naphthyl, which is independently substituted with 0-2  $R^1$  substituents; and
- (e) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^1$  substituents;

$R^1$  is selected from:

halo,  $C_{1-4}$ alkyl, -CN, -NO<sub>2</sub>,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$ ,  $SO_2R^2$ , CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

$R^2$  and  $R^3$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;

m is an integer of 0-2;

B is a member selected from the group consisting of:

a direct link, -C(=O)-, -N( $R^4$ )-, -C(=O)-N( $R^4$ )-, -N( $R^4$ )-C(=O)-, -SO<sub>2</sub>-, -O-, -SO<sub>2</sub>-N( $R^4$ )- and -N( $R^4$ )-SO<sub>2</sub>-;

$R^4$  is selected from:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;

D is phenyl, which is independently substituted with 0-2  $R^{1a}$  substituents;


$R^{1a}$  is selected from:

halo,  $C_{1-4}$ alkyl,  $-CN$ ,  $-NO_2$ ,  $-(CH_2)_mNR^{2a}R^{3a}$ ,  $-SO_2NR^{2a}R^{3a}$ ,  $-SO_2R^{2a}$ ,  $CF_3$ ,  $-OR^{2a}$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

$R^{2a}$  and  $R^{3a}$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;

M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



wherein 0 to 2 of the hydrogen atoms on the D portion of the bicyclic ring may be replaced by  $R^{1a}$  substituents as defined above;

$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $-CN$ ,  $-NO_2$ ,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$ ,  $SO_2R^2$ ,  $CF_3$ ,  $OR^2$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

$R^{17b}$  and  $R^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy,  $-C_{1-4}alkyl$ ,  $-CN$ ,  $-NO_2$ ,  $-(CH_2)_mNR^2R^3$ ,  $-SO_2NR^2R^3$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S;

$m$  is an integer from 0-6;

$E$  is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-(-C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-)$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H,  $C_{1-4}alkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $C_{0-4}alkylheteroaryl$ ,  $C_{1-4}alkylCOOH$  and  $C_{1-4}alkylCOOC_{1-4}alkyl$ , wherein from 0-2 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}alkyl$ ,  $-OH$ ,  $-O-C_{1-4}alkyl$ ,  $-SH$ ,  $-S-C_{1-4}alkyl$ ,  $-CN$  and  $-NO_2$ ;

$G$  is selected from:

a direct link,  $-C(R^7, R^8)-$ ,  $-C(R^{7a}, R^{8a})-C(R^{7b}, R^{8b})-$  and  $-C(R^{7c})=C(R^{8c})-$ ;

wherein  $R^7$ ,  $R^8$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{7b}$ ,  $R^{8b}$ ,  $R^{7c}$  and  $R^{8c}$  are independently a member selected from from the group consisting of:

hydrogen, halogen,  $C_{1-4}alkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $-OR^9$ ,  $-N(R^9R^{10})$ ,  $-C_{0-4}alkylCOOR^9$ ,  $-C_{0-4}alkylC(=O)NR^9R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9-CH_2-CH_2-O-R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9(-CH_2-CH_2-O-R^{10})_2$ ,  $-N(R^9)COR^{10}$ ,  $-N(R^9)C(=O)R^{10}$ ,  $-N(R^9)SO_2R^{10}$ , and a naturally occurring or synthetic amino acid side chain;

$R^9$  and  $R^{10}$  are independently selected from:

H,  $C_{1-4}alkyl$ ,  $C_{0-4}alkylphenyl$  and  $C_{0-4}alkylnaphthyl$ ;

$J$  is a member selected from the group consisting of:

a direct link,  $-O-$ ,  $-O-C(-R^{11}, -R^{11a})-$ ,  $-S-$ ,  $-S(=O)_2-$ ,  $-S-C(-R^{11}, -R^{11a})-$ ,  $-S(=O)_2-(-R^{11}, -R^{11a})-$ ,  $-C(=O)-N(R^{11b})-$ ,  $-N(R^{11b})-$ ,  $-N(R^{11b})-C(-R^{11}, -R^{11a})-$  and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring

atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{11c}$  substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$  and  $R^{11c}$  are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>3-8</sub>cycloalkyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, and a C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1b}$  substituents;
- (b) naphthyl, which is independently substituted with 0-2  $R^{1b}$  substituents; and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{1b}$  substituents;

$R^{1b}$  is a member selected from the group consisting of:

halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>R<sup>2b</sup>, CF<sub>3</sub>, OR<sup>2b</sup>, O-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, O-CH<sub>2</sub>-COOR<sup>2b</sup>, N(R<sup>2b</sup>)-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, N(-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>)<sub>2</sub>, N(R<sup>2b</sup>)-C(=O)R<sup>3b</sup>, N(R<sup>2b</sup>)-SO<sub>2</sub>-R<sup>3b</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

$R^{2b}$  and  $R^{3b}$  are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

L is selected from:

H, -CN, C(=O)NR<sup>12</sup>R<sup>13</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup>, C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, OR<sup>12</sup>, -NR<sup>12</sup>C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, and NR<sup>12</sup>C(=NR<sup>12</sup>)-R<sup>13</sup>;

n is an integer from 0 to 6;

$R^{12}$  and  $R^{13}$  are independently selected from:


hydrogen,  $-OR^{14}$ ,  $-NR^{14}R^{15}$ ,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $COOC_{1-4}$ alkyl,  $COO-C_{0-4}$ alkylphenyl and  $COO-C_{0-4}$ alkylnaphthyl, wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $-OH$ ,  $-O-C_{1-4}$ alkyl,  $C_{1-4}$ alkyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$ , and  $-NO_2$ ;

$R^{14}$  and  $R^{15}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

In a still further preferred embodiment, the present invention provides a compound according to the formula I:



wherein:

A is selected from:

- (a) phenyl, which is independently substituted with 0-2  $R^1$  substituents; and
- (b) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^1$  substituents;

$R^1$  is selected from:

halo,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$  and  $SO_2R^2$ ;

$R^2$  and  $R^3$  are independently selected from the group consisting of:

H and  $C_{1-4}$ alkyl;

m is an integer of 0-2;

Q is a member selected from the group consisting of:




a direct link,  $-\text{C}(=\text{O})-$ ,  $-\text{SO}_2-$ , and  $-\text{O}-$ ;

D is phenyl, which is independently substituted with 0-2  $\text{R}^{1a}$  substituents;

$\text{R}^{1a}$  is selected from:

halo and  $\text{C}_{1-4}$ alkyl;

M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



$\text{R}^{16}$ ,  $\text{R}^{17}$ ,  $\text{R}^{17a}$ ,  $\text{R}^{18}$ ,  $\text{R}^{18a}$ ,  $\text{R}^{18b}$ ,  $\text{R}^{19}$  and  $\text{R}^{19a}$  are each independently selected from the group consisting of:

halo,  $\text{C}_{1-4}$ alkyl,  $-\text{CN}$ ,  $-\text{NO}_2$ ,  $(\text{CH}_2)_m\text{NR}^2\text{R}^3$ ,  $\text{SO}_2\text{NR}^2\text{R}^3$ ,  $\text{SO}_2\text{R}^2$ ,  $\text{CF}_3$  and  $\text{OR}^2$ ;

$\text{R}^{17b}$  and  $\text{R}^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy,  $-C_{1-4}alkyl$ ,  $-CN$ ,  $-NO_2$ ,  $-(CH_2)_mNR^2R^3$ ,  $-SO_2NR^2R^3$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-3 N atoms;

E is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-(-C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-)$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H,  $C_{1-4}alkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $C_{0-4}alkylheteroaryl$ ,  $C_{1-4}alkylCOOH$  and  $C_{1-4}alkylCOOC_{1-4}alkyl$ ;

G is selected from:

a direct link,  $-C(R^7, R^8)-$ ,  $-C(R^{7a}, R^{8a})-C(R^{7b}, R^{8b})-$  and  $-C(R^{7c})=C(R^{8c})-$ ;

wherein  $R^7$ ,  $R^8$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{7b}$ ,  $R^{8b}$ ,  $R^{7c}$  and  $R^{8c}$  are independently a member selected from from the group consisting of:

hydrogen, halogen,  $C_{1-4}alkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $-OR^9$ ,  $-N(R^9R^{10})$ ,  $-C_{0-4}alkylCOOR^9$ ,  $-C_{0-4}alkylC(=O)NR^9R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9-CH_2-CH_2-O-R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9(-CH_2-CH_2-O-R^{10})_2$ ,  $-N(R^9)COR^{10}$ ,  $-N(R^9)C(=O)R^{10}$ ,  $-N(R^9)SO_2R^{10}$ , and a naturally occurring or synthetic amino acid side chain;

$R^9$  and  $R^{10}$  are independently selected from:

H,  $C_{1-4}alkyl$ ,  $C_{0-4}alkylphenyl$  and  $C_{0-4}alkylnaphthyl$ ;

J is a member selected from the group consisting of:

a direct link,  $-O-$ ,  $-S-$ ,  $-C(=O)-N(R^{11b})-$ ,  $-N(R^{11b})-$ ,  $-N(R^{11b})-C(-R^{11}, -R^{11a})-$  and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{11c}$  substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$  and  $R^{11c}$  are a member independently selected from the group consisting of:

hydrogen, halo,  $-CN$ ,  $-NO_2$ ,  $-OH$ ,  $-O-C_{1-4}alkyl$ ,  $-O-C_{3-8}cycloalkyl$ ,  $-COOH$ ,  $-C(=O)-O-C_{1-4}alkyl$ ,  $-C(=O)-O-C_{3-8}cycloalkyl$ ,  $C_{1-4}alkyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ , and a  $C_{0-4}alkylheterocyclic$  ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S,  $CH_2COOC_{1-4}alkyl$ ,  $CH_2COOC_{1-4}alkylphenyl$  and  $CH_2COOC_{1-4}alkylnaphthyl$ ;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1b}$  substituents;
- (b) an aromatic heterocyclic ring having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring may be substituted with 0-2  $R^{1b}$  substituents;
- (c) a fused aromatic bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the bicyclic ring system may be substituted with 0-2  $R^{1b}$  substituents;

$R^{1b}$  is a member selected from the group consisting of:

halo,  $-C_{1-4}alkyl$ ,  $-OH$ ,  $-OBn$ ,  $-O-CH_2-CH_2-OH$ ,  $-O-CH_2-CH_2-OCH_3$ ,  
 $-O-CH_2-COOH$ ,  $-O-CH_2-C(=O)-O-CH_3$ ,  $-NH_2$ ,  $-NH-CH_2-CH_2-O-CH_3$ ,  
 $-NH-C(=O)-O-CH_3$  and  $-NH-SO_2-CH_3$ ;

L is selected from:

H,  $-C(=O)NR^{12}R^{13}$ ,  $-(CH_2)_nNR^{12}R^{13}$  and  $-C(=NR^{12})NR^{12}R^{13}$ ;


n is an integer from 0 to 6;

$R^{12}$  and  $R^{13}$  are independently selected from:

hydrogen and  $C_{1-4}alkyl$ ;


and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

In yet another preferred embodiment, the present invention provides a compound according to formula I:




wherein:

A is a member selected from the group consisting of:



D is a member selected from the group consisting of:



E is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-(-C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-)$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $C_{0-4}$ alkylheteroaryl,  $C_{1-4}$ alkylCOOH and  $C_{1-4}$ alkylCOOC $_{1-4}$ alkyl;

G is selected from:

a direct link,  $-C(R^7, R^8)-$ ,  $-C(R^{7a}, R^{8a})-C(R^{7b}, R^{8b})-$  and  $-C(R^{7c})=C(R^{8c})-$ ;

wherein  $R^7$ ,  $R^8$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{7b}$ ,  $R^{8b}$ ,  $R^{7c}$  and  $R^{8c}$  are independently a member selected from from the group consisting of:

hydrogen, halogen,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $-OR^9$ ,  $-N(R^9R^{10})$ ,  $-C_{0-4}$ alkylCOOR<sup>9</sup>,  $-C_{0-4}$ alkylC(=O)NR<sup>9</sup>R<sup>10</sup>,  $-C_{0-4}$ alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>,  $-C_{0-4}$ alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)-<sub>2</sub>,  $-N(R^9)COR^{10}$ ,  $-N(R^9)C(=O)R^{10}$ ,  $-N(R^9)SO_2R^{10}$ , and a naturally occurring or synthetic amino acid side chain;

$R^9$  and  $R^{10}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;

J is a member selected from the group consisting of:


a direct link, -O-, -S-,  $-C(=O)-N(R^{11b})-$ ,  $-N(R^{11b})-$ ,  $-N(R^{11b})-C(-R^{11}, -R^{11a})-$  and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{11c}$  substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$  and  $R^{11c}$  are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O- $C_{1-4}$ alkyl,  $-C_{1-4}$ alkyl, -COOH, phenyl, and benzyl wherein the aromatic ring of the phenyl or benzyl is substituted with 0-2 members independently selected from the group consisting of halo, -CN, -NO<sub>2</sub>, -OH, -O- $C_{1-4}$ alkyl,  $-C_{1-4}$ alkyl, -COOH and  $-C(=O)-O-C_{1-4}$ alkyl;



Y and L taken together are a member selected from the group consisting of:



M and Q are as defined elsewhere in the specification;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

The following non-limiting tables illustrate representative compounds of the present invention wherein the "Y-L" portions for each of the formula in each of the tables are taken together and are independently selected from the group consisting of:






Table 1



Formula II

$\text{R}^7$	$\text{R}^{11b}$	$\text{R}^7$	$\text{R}^{11b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 1a



Formula IIa

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 1b



R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et



Table 1c



Formula IIc


















R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 1d



Formula IIId

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 1e



Formula IIe

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 1f



Formula II f


















$\text{R}^7$	$\text{R}^{11b}$	$\text{R}^7$	$\text{R}^{11b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 1g




Formula IIg

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 1h



Formula IIIh

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 1i



Formula Iii

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 1j



Formula IIj

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 1k



Formula IIk


















$\text{R}^7$	$\text{R}^{11b}$	$\text{R}^7$	$\text{R}^{11b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 1(l)




Formula II(l)

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et




Table 1m



Formula IIIm

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 1n



Formula IIIn


















R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 1o



Formula IIo

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 1p



Formula IIp

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 1q




Formula IIq

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 2



Formula III

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2a



Formula IIIa

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2b



Formula IIIb

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2c



Formula IIIc

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2d



Formula IIIId

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl

Table 2e




Formula IIIe

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl




Table 2f



Formula IIIf

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2g



Formula IIIg

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2h



Formula IIIh

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2i



Formula IIIi

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2j



Formula IIIj

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 2k



Formula IIIk

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 2(l)




Formula III(l)

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl




Table 2m



Formula III m

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2n



Formula IIIIn

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 2o



Formula IIIo

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2p



Formula IIIp

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2q



Formula IIIq

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 3



Formula IV

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 3a




Formula VIa

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 3b



Formula VIb

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3c



Formula VIc

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3d



Formula IVd

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3e



Formula IVe

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3f



Formula IVf

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3g



Formula IVg

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 3h




Formula IVh

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 3i



Formula IVi

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 3j



Formula IVj


















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 3k



Formula IVk

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3(1)



Formula IV(1)

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3m



Formula IVm

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3n



Formula IVn

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 3o



Formula IVo

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 3p




Formula IVp

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et




Table 3q



Formula IVq

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 4



Formula V

$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 4a



Formula Va

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4b



Formula Vb


















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 4c



Formula Vc

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 4d



Formula Vd

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 4e



Formula Ve

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4f




Formula Vf

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 4g



Formula Vg

$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 4h



Formula Vh

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 4i



Formula Vi

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 4j



Formula Vj

$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4k



Formula Vk


















$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 4(1)



Formula V(l)


















$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 4m



Formula Vm

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4n




Formula Vn

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



Table 4o



Formula Vo


















$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 4p



Formula Vp


















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 4q



Formula Vq


















$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 5



Formula VI

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 5a



Formula VIa


















R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 5b



Formula VIb

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 5c



Formula VIc

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 5d




Formula VIId

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



Table 5e



Formula VIe


















R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 5f



Formula VI f

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 5g



Formula VIg

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 5h



Formula VIh

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 5i



Formula VII

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 5j



Formula VIj

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 5k



Formula VIk

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 5(1)




Formula VI(1)

$\text{R}^{7c}$	$\text{R}^{8c}$	$\text{R}^{7c}$	$\text{R}^{8c}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 5m



Formula VIIm

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 5n



Formula VIIn

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 5o



Formula VIo

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 5p



Formula VIp


















$\text{R}^{7c}$	$\text{R}^{8c}$	$\text{R}^{7c}$	$\text{R}^{8c}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 5q



Formula VIq

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6



Formula VII

$\text{R}^{7a}$	$\text{R}^{7b}$	$\text{R}^{7a}$	$\text{R}^{7b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6a



Formula VIIa

$\text{R}^{7a}$	$\text{R}^{7b}$	$\text{R}^{7a}$	$\text{R}^{7b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 6b




Formula VIIb

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et




Table 6c



Formula VIIc

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6d



Formula VIId

$\text{R}^{7a}$	$\text{R}^{7b}$	$\text{R}^{7a}$	$\text{R}^{7b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6e



Formula VIIe

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6f



Formula VIIf

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6g



Formula VIIg

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6h



Formula VIIh

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6i



Formula VIIi

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 6j




Formula VIIj

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 6k



Formula VIIk

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6(1)



Formula VII(1)

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6m



Formula VIIm

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6n



Formula VIIIn

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 6o



Formula VIIo

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6p



Formula VIIp

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6q



Formula VIIq

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 7




Formula VIII

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 7a



Formula VIIIa

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7b



Formula VIIIb

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7c



Formula VIIIc

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7d



Formula VIII d

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 7e



Formula VIIIe

$\text{R}^{7a}$	$\text{R}^{7b}$	$\text{R}^{7a}$	$\text{R}^{7b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7f



Formula VIII f

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7g



Formula VIIIg

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 7h




Formula VIIIh

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 7i



Formula VIIIi

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7j



Formula VIIIj

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7k



Formula VIIIk

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 7(1)



Formula VIII(1)

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7m



Formula VIII m

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7n



Formula VIII n

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7o



Formula VIIIo

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 7p




Formula VIIIp

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 7q



Formula VIIIq

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 8



Formula IX

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8a



Formula IXa

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8b



Formula IXb

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8c



Formula IXc

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8d



Formula IXd

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8e



Formula IXe

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 8f




Formula IXf

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl




Table 8g



Formula IXg

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8h



Formula IXh

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8i



Formula IXi

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8j



Formula IXj

$\text{R}^{6\text{c}}$	$\text{R}^{11\text{b}}$	$\text{R}^{6\text{c}}$	$\text{R}^{11\text{b}}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 8k



Formula IXk

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8(l)



Formula IX(l)

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8m



Formula IXm

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 8n




Formula IXn

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl




Table 8o



Formula IXo

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8p



Formula IXp

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8q



Formula IXq

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 9



Formula X

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 9a



Formula Xa

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 9b



Formula Xb

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 9c



Formula Xc

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9d




Formula Xd

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl




Table 9e



Formula Xe

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 9f



Formula Xf

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl

Table 9g



Formula Xg














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9h



Formula Xh














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 9i



Formula Xi

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9j



Formula Xj














$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 9k



Formula Xk

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9(l)




Formula X(l)

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl




Table 9m



Formula Xm

$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 9n



Formula Xn

$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl

Table 9o



Formula Xo














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9p



Formula Xp














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9q



Formula Xq














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10



Formula XI

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10a



Formula XIa

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10b




Formula XIb

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl



Table 10c



Formula XIc














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10d



Formula XIId














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10e



Formula XIe

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 10f



Formula XI f

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10g



Formula XIg

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl

Table 10h



Formula XIh














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10i



Formula XII

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10j



Formula XIj














$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl




Table 10k



Formula XIk

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10(I)



Formula XI(I)

$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 10m



Formula XIIm

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>2</sup> R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10n



Formula XIn














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10o



Formula XIo

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10p



Formula XIp














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10q



Formula XIq














$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11



Formula XII














$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl



Table 11a



Formula XIa














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11b



Formula XIb














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 11c



Formula XIc

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11d



Formula XIIId














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 11e



Formula XIIe

$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 11f



Formula XIIIf

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 11g



Formula XIIg

$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl

Table 11h




Formula XIIIh

$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl




Table 1 li



Formula XIII

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11j



Formula XIIj














$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 11k



Formula XIIk

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 11(l)



Formula XII(l)

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 11m



Formula XIIIm

$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl

Table 11n




























R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11q




Formula XIIq













R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

# **MISSING AT THE TIME OF PUBLICATION**




Table 11q




R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Also preferred are compounds according to Tables 1 through Table 11q, wherein



the biphenylene portions of their formulae:

are each replaced with the following ring structure:




5

wherein each of the A and D ring portions are as follows:

A	D	A	D

Even more preferred compounds are set forth in Tables 12-24, below.

**Table 12**



**Table 13**




Table 14




Table 15




Table 16





Table 17



5 wherein A—Q— is selected from the group consisting of:

t-Bu ; O-t-Bu ;  $-(CH_2)_{0-5}$ -amino ; OH ; carboxylic acid ester; carboxamide ;






Table 18




Table 19








Table 21







Table 22




;




;




;




;



;



;



and




Table 23




Table 24




Table 25




Table 26






Table 27




Wherein:

Q is a direct link, and A is a member selected from the group:



or Q is a -C(=NH)- group, and A is a member selected from the group:




G is a direct link;

J is a member selected from the group:



Y-L is a member selected from the group:



- 5 This invention also encompasses all pharmaceutically acceptable isomers, salts, hydrates and solvates of the compounds of formula I. In addition, the compounds of formula I can exist in various isomeric and tautomeric forms, and all such forms are meant to be included in the invention, along with pharmaceutically acceptable salts, hydrates and solvates of such isomers and tautomers.

The compounds of this invention may be isolated as the free acid or base or converted to salts of various inorganic and organic acids and bases. Such salts are within the scope of this invention. Non-toxic and physiologically compatible salts are particularly useful although other less desirable salts may have use in the processes of isolation and purification.

A number of methods are useful for the preparation of the salts described above and are known to those skilled in the art. For example, the free acid or free base form of a compound of one of the formulas above can be reacted with one or more molar equivalents of the desired acid or base in a solvent or solvent mixture in which the salt is insoluble, or in a solvent like water after which the solvent is removed by evaporation, distillation or freeze drying. Alternatively, the free acid or base form of the product may be passed over an ion exchange resin to form the desired salt or one salt form of the product may be converted to another using the same general process.

#### Prodrug Derivatives of Compounds

This invention also encompasses prodrug derivatives of the compounds contained herein. The term "prodrug" refers to a pharmacologically inactive derivative of a parent drug molecule that requires biotransformation, either spontaneous or enzymatic, within the organism to release the active drug. Prodrugs are variations or derivatives of the compounds of this invention which have groups cleavable under metabolic conditions. Prodrugs become the compounds of the invention which are pharmaceutically active *in vivo*, when they undergo solvolysis under physiological conditions or undergo enzymatic degradation. Prodrug compounds of this invention may be called single, double, triple etc., depending on the number of biotransformation steps required to release the active drug within the organism, and indicating the number of functionalities present in a precursor-type form. Prodrug forms often offer advantages of solubility, tissue compatibility, or delayed release in the mammalian organism (see, Bundgard, Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985 and Silverman, The Organic Chemistry of Drug Design and Drug Action, pp. 352-401, Academic Press, San Diego, CA, 1992). Prodrugs commonly known in the art include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acids with a suitable alcohol, or amides prepared by reaction of the parent acid compound with an amine, or basic groups reacted to form an acylated base derivative. Moreover, the



prodrug derivatives of this invention may be combined with other features herein taught to enhance bioavailability.

As mentioned above, the compounds of this invention find utility as therapeutic agents for disease states in mammals which have disorders of coagulation such as in the treatment or prevention of unstable angina, refractory angina, myocardial infarction, transient ischemic attacks, thrombotic stroke, embolic stroke, disseminated intravascular coagulation including the treatment of septic shock, deep venous thrombosis in the prevention of pulmonary embolism or the treatment of reocclusion or restenosis of reperfused coronary arteries. Further, these compounds are useful for the treatment or prophylaxis of those diseases which involve the production and/or action of factor Xa/prothrombinase complex. This includes a number of thrombotic and prothrombotic states in which the coagulation cascade is activated which include but are not limited to, deep venous thrombosis, pulmonary embolism, myocardial infarction, stroke, thromboembolic complications of surgery and peripheral arterial occlusion.

Accordingly, a method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprises administering to the mammal a therapeutically effective amount of a compound of this invention. In addition to the disease states noted above, other diseases treatable or preventable by the administration of compounds of this invention include, without limitation, occlusive coronary thrombus formation resulting from either thrombolytic therapy or percutaneous transluminal coronary angioplasty, thrombus formation in the venous vasculature, disseminated intravascular coagulopathy, a condition wherein there is rapid consumption of coagulation factors and systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the microvasculature leading to widespread organ failure, hemorrhagic stroke, renal dialysis, blood oxygenation, and cardiac catheterization.

The compounds of the invention also find utility in a method for inhibiting the coagulation of biological samples, (*e.g.* blood) which comprises the administration of a compound of the invention.

The compounds of the present invention may also be used in combination with other therapeutic or diagnostic agents. In certain preferred embodiments, the compounds of this invention may be coadministered along with other compounds typically prescribed

for these conditions according to generally accepted medical practice such as anticoagulant agents, thrombolytic agents, or other antithrombotics, including platelet aggregation inhibitors, tissue plasminogen activators, urokinase, prourokinase, streptokinase, heparin, aspirin, or warfarin. The compounds of the present invention may act in a synergistic fashion to prevent reocclusion following a successful thrombolytic therapy and/or reduce the time to reperfusion. These compounds may also allow for reduced doses of the thrombolytic agents to be used and therefore minimize potential hemorrhagic side-effects. The compounds of this invention can be utilized *in vivo*, ordinarily in mammals such as primates, (e.g. humans), sheep, horses, cattle, pigs, dogs, cats, rats and mice, or *in vitro*.

The biological properties of the compounds of the present invention can be readily characterized by methods that are well known in the art, for example by the *in vitro* protease activity assays and *in vivo* studies to evaluate antithrombotic efficacy, and effects on hemostasis and hematological parameters, such as are illustrated in the examples.

Diagnostic applications of the compounds of this invention will typically utilize formulations in the form of solutions or suspensions. In the management of thrombotic disorders the compounds of this invention may be utilized in compositions such as tablets, capsules or elixirs for oral administration, suppositories, sterile solutions or suspensions or injectable administration, and the like, or incorporated into shaped articles. Subjects in need of treatment (typically mammalian) using the compounds of this invention can be administered dosages that will provide optimal efficacy. The dose and method of administration will vary from subject to subject and be dependent upon such factors as the type of mammal being treated, its sex, weight, diet, concurrent medication, overall clinical condition, the particular compounds employed, the specific use for which these compounds are employed, and other factors which those skilled in the medical arts will recognize.

Formulations of the compounds of this invention are prepared for storage or administration by mixing the compound having a desired degree of purity with physiologically acceptable carriers, excipients, stabilizers etc., and may be provided in sustained release or timed release formulations. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical field, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co., (A.R. Gennaro edit.

1985). Such materials are nontoxic to the recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, acetate and other organic acid salts, antioxidants such as ascorbic acid, low molecular weight (less than about ten residues) peptides such as polyarginine, proteins, such as serum albumin, gelatin, or immunoglobulins, hydrophilic polymers such as polyvinylpyrrolidinone, amino acids such as glycine, glutamic acid, aspartic acid, or arginine, monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose or dextrans, chelating agents such as EDTA, sugar alcohols such as mannitol or sorbitol, counterions such as sodium and/or nonionic surfactants such as Tween, Pluronics or polyethyleneglycol.

Dosage formulations of the compounds of this invention to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile membranes such as 0.2 micron membranes, or by other conventional methods. Formulations typically will be stored in lyophilized form or as an aqueous solution. The pH of the preparations of this invention typically will be 3-11, more preferably 5-9 and most preferably 7-8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of cyclic polypeptide salts. While the preferred route of administration is by injection, other methods of administration are also anticipated such as orally, intravenously (bolus and/or infusion), subcutaneously, intramuscularly, colonically, rectally, nasally, transdermally or intraperitoneally, employing a variety of dosage forms such as suppositories, implanted pellets or small cylinders, aerosols, oral dosage formulations and topical formulations such as ointments, drops and dermal patches. The compounds of this invention are desirably incorporated into shaped articles such as implants which may employ inert materials such as biodegradable polymers or synthetic silicones, for example, Silastic, silicone rubber or other polymers commercially available.

The compounds of the invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of lipids, such as cholesterol, stearylamine or phosphatidylcholines.

The compounds of this invention may also be delivered by the use of antibodies,

antibody fragments, growth factors, hormones, or other targeting moieties, to which the compound molecules are coupled. The compounds of this invention may also be coupled with suitable polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidinone, pyran copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxyethyl-aspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, compounds of the invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels. Polymers and semipermeable polymer matrices may be formed into shaped articles, such as valves, stents, tubing, prostheses and the like.

Therapeutic compound liquid formulations generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by hypodermic injection needle.

Therapeutically effective dosages may be determined by either *in vitro* or *in vivo* methods. For each particular compound of the present invention, individual determinations may be made to determine the optimal dosage required. The range of therapeutically effective dosages will be influenced by the route of administration, the therapeutic objectives and the condition of the patient. For injection by hypodermic needle, it may be assumed the dosage is delivered into the body's fluids. For other routes of administration, the absorption efficiency must be individually determined for each compound by methods well known in pharmacology. Accordingly, it may be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. The determination of effective dosage levels, that is, the dosage levels necessary to achieve the desired result, will be readily determined by one skilled in the art. Typically, applications of compound are commenced at lower dosage levels, with dosage levels being increased until the desired effect is achieved.

The compounds of the invention can be administered orally or parenterally in an effective amount within the dosage range of about 0.1 to 100 mg/kg, preferably about 0.5



to 50 mg/kg and more preferably about 1 to 20 mg/kg on a regimen in a single or 2 to 4 divided daily doses and/or continuous infusion.

Typically, about 5 to 500 mg of a compound or mixture of compounds of this invention, as the free acid or base form or as a pharmaceutically acceptable salt, is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, dye, flavor etc., as called for by accepted pharmaceutical practice. The amount of active ingredient in these compositions is such that a suitable dosage in the range indicated is obtained.

10

Typical adjuvants which may be incorporated into tablets, capsules and the like are binders such as acacia, corn starch or gelatin, and excipients such as microcrystalline cellulose, disintegrating agents like corn starch or alginic acid, lubricants such as magnesium stearate, sweetening agents such as sucrose or lactose, or flavoring agents. When a dosage form is a capsule, in addition to the above materials it may also contain liquid carriers such as water, saline, or a fatty oil. Other materials of various types may be used as coatings or as modifiers of the physical form of the dosage unit. Sterile compositions for injection can be formulated according to conventional pharmaceutical practice. For example, dissolution or suspension of the active compound in a vehicle such as an oil or a synthetic fatty vehicle like ethyl oleate, or into a liposome may be desired. Buffers, preservatives, antioxidants and the like can be incorporated according to accepted pharmaceutical practice.

15

20

### Preparation of Compounds

25

The compounds of the present invention may be synthesized by either solid or liquid phase methods described and referenced in standard textbooks, or by a combination of both methods. These methods are well known in the art. See, Bodanszky, "The Principles of Peptide Synthesis", Hafner, *et al.*, Eds., Springer-Verlag, Berlin, 1984.

30

Starting materials used in any of these methods are commercially available from chemical vendors such as Aldrich, Sigma, Nova Biochemicals, Bachem Biosciences, and the like, or may be readily synthesized by known procedures.


Reactions are carried out in standard laboratory glassware and reaction vessels under reaction conditions of standard temperature and pressure, except where otherwise indicated.

5


During the synthesis of these compounds, the functional groups of the amino acid derivatives used in these methods are protected by blocking groups to prevent cross reaction during the coupling procedure. Examples of suitable blocking groups and their use are described in "The Peptides: Analysis, Synthesis, Biology", Academic Press, Vol. 3  
10 (Gross, *et al.*, Eds., 1981) and Vol. 9 (1987), the disclosures of which are incorporated herein by reference.

Non-limiting exemplary synthesis schemes are outlined directly below, and specific steps are described in the Examples. The reaction products are isolated and  
15 purified by conventional methods, typically by solvent extraction into a compatible solvent. The products may be further purified by column chromatography or other appropriate methods.

Scheme 1




Scheme 2







### Scheme 3




Scheme 4




Scheme 5




Scheme 6




Scheme 7




Scheme 8




Scheme 9



Scheme 10




Scheme 11







Scheme 12




Scheme 13




Scheme 14



Scheme 15



Scheme 16



### Compositions and Formulations

The compounds of this invention may be isolated as the free acid or base or converted to salts of various inorganic and organic acids and bases. Such salts are within the scope of this invention. Non-toxic and physiologically compatible salts are particularly  
5 useful although other less desirable salts may have use in the processes of isolation and purification.

A number of methods are useful for the preparation of the salts described above and are known to those skilled in the art. For example, reaction of the free acid or free base form of a compound of the structures recited above with one or more molar  
10 equivalents of the desired acid or base in a solvent or solvent mixture in which the salt is insoluble, or in a solvent like water after which the solvent is removed by evaporation, distillation or freeze drying. Alternatively, the free acid or base form of the product may be passed over an ion exchange resin to form the desired salt or one salt form of the product may be converted to another using the same general process.

15 Diagnostic applications of the compounds of this invention will typically utilize formulations such as solution or suspension. In the management of thrombotic disorders the compounds of this invention may be utilized in compositions such as tablets, capsules or elixirs for oral administration, suppositories, sterile solutions or suspensions or injectable administration, and the like, or incorporated into shaped articles. Subjects in  
20 need of treatment (typically mammalian) using the compounds of this invention can be administered dosages that will provide optimal efficacy. The dose and method of administration will vary from subject to subject and be dependent upon such factors as the type of mammal being treated, its sex, weight, diet, concurrent medication, overall clinical condition, the particular compounds employed, the specific use for which these  
25 compounds are employed, and other factors which those skilled in the medical arts will recognize.

Formulations of the compounds of this invention are prepared for storage or administration by mixing the compound having a desired degree of purity with physiologically acceptable carriers, excipients, stabilizers etc., and may be provided in sustained release or timed release formulations. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical field, and are described, for example, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., (A.R. Gennaro edit. 1985). Such materials are nontoxic to the recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, acetate and other organic acid salts, antioxidants such as ascorbic acid, low molecular weight (less than about ten residues) peptides such as polyarginine, proteins, such as serum albumin, gelatin, or immunoglobulins, hydrophilic polymers such as polyvinylpyrrolidinone, amino acids such as glycine, glutamic acid, aspartic acid, or arginine, monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose or dextrans, chelating agents such as EDTA, sugar alcohols such as mannitol or sorbitol, counterions such as sodium and/or nonionic surfactants such as Tween, Pluronic or polyethyleneglycol.

Dosage formulations of the compounds of this invention to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile membranes such as 0.2 micron membranes, or by other conventional methods. Formulations typically will be stored in lyophilized form or as an aqueous solution. The pH of the preparations of this invention typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of cyclic polypeptide salts. While the preferred route of administration is by injection, other methods of administration are also anticipated such as intravenously (bolus and/or infusion), subcutaneously, intramuscularly, colonically, rectally, nasally or intraperitoneally, employing a variety of dosage forms such as suppositories, implanted pellets or small cylinders, aerosols, oral dosage formulations and topical formulations such

as ointments, drops and dermal patches. The compounds of this invention are desirably incorporated into shaped articles such as implants which may employ inert materials such as biodegradable polymers or synthetic silicones, for example, Silastic, silicone rubber or other polymers commercially available.

5           The compounds of this invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of lipids, such as cholesterol, stearylamine or phosphatidylcholines.

          The compounds of this invention may also be delivered by the use of antibodies,  
10   antibody fragments, growth factors, hormones, or other targeting moieties, to which the compound molecules are coupled. The compounds of this invention may also be coupled with suitable polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxyethyl-aspartamide-phenol, or polyethyleneoxide-polylysine substituted with  
15   palmitoyl residues. Furthermore, the factor Xa inhibitors of this invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross linked or amphipathic block copolymers  
20   of hydrogels. Polymers and semipermeable polymer matrices may be formed into shaped articles, such as valves, stents, tubing, prostheses and the like.

          Therapeutic compound liquid formulations generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by hypodermic injection needle.

25           Therapeutically effective dosages may be determined by either *in vitro* or *in vivo* methods. For each particular compound of the present invention, individual determinations may be made to determine the optimal dosage required. The range of



therapeutically effective dosages will naturally be influenced by the route of administration, the therapeutic objectives, and the condition of the patient. For injection by hypodermic needle, it may be assumed the dosage is delivered into the body's fluids. For other routes of administration, the absorption efficiency must be individually  
5 determined for each inhibitor by methods well known in pharmacology. Accordingly, it may be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. The determination of effective dosage levels, that is, the dosage levels necessary to achieve the desired result, will be within the ambit of one skilled in the art. Typically, applications of compound are  
10 commenced at lower dosage levels, with dosage levels being increased until the desired effect is achieved.

A typical dosage might range from about 0.001 mg/kg to about 1000 mg/kg, preferably from about 0.01 mg/kg to about 100 mg/kg, and more preferably from about 0.10 mg/kg to about 20 mg/kg. Advantageously, the compounds of this invention may be  
15 administered several times daily, and other dosage regimens may also be useful.

Typically, about 0.5 to 500 mg of a compound or mixture of compounds of this invention, as the free acid or base form or as a pharmaceutically acceptable salt, is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, dye, flavor etc., as called for by accepted pharmaceutical practice.  
20 The amount of active ingredient in these compositions is such that a suitable dosage in the range indicated is obtained.

Typical adjuvants which may be incorporated into tablets, capsules and the like are a binder such as acacia, corn starch or gelatin, and excipient such as microcrystalline cellulose, a disintegrating agent like corn starch or alginic acid, a lubricant such as  
25 magnesium stearate, a sweetening agent such as sucrose or lactose, or a flavoring agent. When a dosage form is a capsule, in addition to the above materials it may also contain a liquid carrier such as water, saline, a fatty oil. Other materials of various types may be

used as coatings or as modifiers of the physical form of the dosage unit. Sterile compositions for injection can be formulated according to conventional pharmaceutical practice. For example, dissolution or suspension of the active compound in a vehicle such as an oil or a synthetic fatty vehicle like ethyl oleate, or into a liposome may be desired. Buffers, preservatives, antioxidants and the like can be incorporated according to accepted pharmaceutical practice.

In practicing the methods of this invention, the compounds of this invention may be used alone or in combination, or in combination with other therapeutic or diagnostic agents. In certain preferred embodiments, the compounds of this inventions may be coadministered along with other compounds typically prescribed for these conditions according to generally accepted medical practice, such as anticoagulant agents, thrombolytic agents, or other antithrombotics, including platelet aggregation inhibitors, tissue plasminogen activators, urokinase, prourokinase, streptokinase, heparin, aspirin, or warfarin. The compounds of this invention can be utilized *in vivo*, ordinarily in mammals such as primates, such as humans, sheep, horses, cattle, pigs, dogs, cats, rats and mice, or *in vitro*.

The preferred compounds of the present invention are characterized by their ability to inhibit thrombus formation with acceptable effects on classical measures of coagulation parameters, platelets and platelet function, and acceptable levels of bleeding complications associated with their use. Conditions characterized by undesired thrombosis would include those involving the arterial and venous vasculature.

With respect to the coronary arterial vasculature, abnormal thrombus formation characterizes the rupture of an established atherosclerotic plaque which is the major cause of acute myocardial infarction and unstable angina, as well as also characterizing the occlusive coronary thrombus formation resulting from either thrombolytic therapy or percutaneous transluminal coronary angioplasty (PTCA).


With respect to the venous vasculature, abnormal thrombus formation characterizes the condition observed in patients undergoing major surgery in the lower extremities or the abdominal area who often suffer from thrombus formation in the venous vasculature resulting in reduced blood flow to the affected extremity and a predisposition to pulmonary embolism. Abnormal thrombus formation further characterizes disseminated intravascular coagulopathy commonly occurs within both vascular systems during septic shock, certain viral infections and cancer, a condition wherein there is rapid consumption of coagulation factors and systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the microvasculature leading to widespread organ failure.

The compounds of this present invention, selected and used as disclosed herein, are believed to be useful for preventing or treating a condition characterized by undesired thrombosis, such as (a) the treatment or prevention of any thrombotically mediated acute coronary syndrome including myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, (b) the treatment or prevention of any thrombotically mediated cerebrovascular syndrome including embolic stroke, thrombotic stroke or transient ischemic attacks, (c) the treatment or prevention of any thrombotic syndrome occurring in the venous system including deep venous thrombosis or pulmonary embolus occurring either spontaneously or in the setting of malignancy, surgery or trauma, (d) the treatment or prevention of any coagulopathy including disseminated intravascular coagulation (including the setting of septic shock or other infection, surgery, pregnancy, trauma or malignancy and whether associated with multi-organ failure or not), thrombotic thrombocytopenic purpura, thromboangiitis obliterans, or thrombotic disease associated with heparin induced thrombocytopenia, (e) the treatment or prevention of thrombotic complications associated with extracorporeal circulation (e.g. renal dialysis, cardiopulmonary bypass or other oxygenation procedure, plasmapheresis), (f) the treatment or prevention of thrombotic complications associated with instrumentation (e.g.

cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve), and (g) those involved with the fitting of prosthetic devices.


Anticoagulant therapy is also useful to prevent coagulation of stored whole blood and to prevent coagulation in other biological samples for testing or storage. Thus the  
5 compounds of this invention can be added to or contacted with any medium containing or suspected to contain factor Xa and in which it is desired that blood coagulation be inhibited, e.g., when contacting the mammal's blood with material such as vascular grafts, stents, orthopedic prostheses, cardiac stents, valves and prostheses, extra corporeal circulation systems and the like.

10 Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. The following working examples therefore, specifically point out preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the  
15 disclosure.

EXAMPLESExample 1

Part 1. Into a chilled suspension of 4-amino-3-nitrobenzoic acid (3.6 g, 20 mmol) in dry methanol (45 mL) was bubbled gaseous HCl for 10 min. The yellow suspension was stirred at room temperature for 2 days. The reaction mixture was filtered, washed with cold methanol and dried under high vacuum to give the HCl salt of methyl 4-amino-3-nitrobenzoate (3.25 g, 70%) as a yellow solid.

Part 2. To a chilled suspension of methyl 4-amino-3-nitrobenzoate (2.56 g, 11 mmol) and 4-dimethylaminopyridine (68 mg, 0.56 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (40 mL) and N,N-diisopropylethylamine (1.9 mL, 11 mmol) was added a solution of di-tert-butylidicarbonate (2.2 g, 10 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) dropwise over 20 min. The suspension was stirred at room temperature for 2 hours. Another solution of di-tert-butylidicarbonate (1.2 g, 5.5 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was added, and the resulting solution was stirred for another hour. The reaction was then concentrated, diluted with ethyl acetate, washed with 5% citric acid, water and brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel chromatography using 5% ethyl acetate in CH<sub>2</sub>Cl<sub>2</sub> as eluent to yield methyl 4-[(tert-butoxy)carbonylamino]-3-nitrobenzoate (2.4 g, 74%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.54 (s, 9H); 3.93 (s, 3H); 8.19-8.22 (d, 1H); 8.66-8.68 (d, 1H); 8.86 (s, 1H).

Example 2


Part 1. To a solution of methyl 4-[(tert-butoxy)carbonylamino]-3-nitrobenzoate (0.44 g, 1.5 mmol) in methanol (2 mL), ethyl acetate (3 mL) and triethylamine (0.21 mL, 1.5

mmol) was added 10% Pd on carbon (159 mg, 0.15 mmol). The reaction mixture was hydrogenated under 1 atm H<sub>2</sub> for 3.5 hours, filtered, and concentrated *in vacuo* to yield methyl 3-amino-4-[(tert-butoxycarbonyl)amino]benzoate (0.40 g, 99%) as a white solid. ES-MS (M+H-C<sub>4</sub>H<sub>8</sub>)<sup>+</sup> = 211.1.

Part 2. To a chilled solution of methyl 3-amino-4-[(tert-butoxy)carbonylamino]benzoate (0.40 g, 1.5 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and triethylamine (0.21 mL, 1.5 mmol) was added a solution of triphosgene (161 mg, 0.54 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (6 mL) over 5 min. The reaction was stirred at room temperature for 2 hr, diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed with water and brine, dried over sodium sulfate and concentrated *in vacuo*. The crude product was purified by silica gel chromatography using 50% ethyl acetate in CH<sub>2</sub>Cl<sub>2</sub> as eluent to give methyl 1-[(tert-butyl)oxycarbonyl]-2-oxo-(3H)-benzimidazole-5-carboxylate (0.27 g, 61%).


<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.68 (s, 9H); 3.91 (s, 3H); 7.73 (s, 1H); 7.79-7.81 (d, 1H); 7.85-7.87 (dd, 1H); 8.9 (s, 1H). ES-MS (M+H-C<sub>4</sub>H<sub>8</sub>)<sup>+</sup> = 237.1.

### Example 3




Part 1. To a chilled solution of methyl 1-[(tert-butyl)oxycarbonyl]-2-oxo-(3H)-benzimidazole-5-carboxylate (0.37 g, 1.28 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (8 mL) and N,N-diisopropylethylamine (0.23 mL, 1.3 mmol) was added a solution of benzyl chloroformate (0.208 mL, 1.46 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL). The reaction was stirred at room temperature for 2.5 hr, diluted with ethyl acetate, washed with water and brine, dried over sodium sulfate and concentrated *in vacuo*. This residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (9 mL) and treated with neat trifluoroacetic acid (2 mL) at 0 °C for 40 min, diluted with ethyl acetate, washed with 5% NaHCO<sub>3</sub>, water and brine, dried over sodium sulfate and concentrated to yield methyl 2-oxo-3-[benzyloxycarbonyl]-(3H)-benzimidazole-5-carboxylate (0.37 g, 89%) as an off-white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 3.51 (s, 3H); 5.14 (s, 2H); 6.67-6.69 (d, 1H); 6.97-7.06 (m, 5H); 7.17-7.18 (d, 1H); 7.51-7.53 (d, 1H); 8.07 (s, 1H). ES-MS (M+Na)<sup>+</sup> = 349.0.



Example 4

To a chilled solution of methyl 2-oxo-3-(benzyloxycarbonyl)-(3H)-benzimidazole-5-carboxylate (110 mg, 0.337 mmol) in DMF (3 mL) was added cesium carbonate (0.275g, 0.84 mmol) followed by 7-(bromomethyl)naphthalene-2-carbonitrile (108 mg, 0.44 mmol). The reaction was stirred at room temperature for 1.5 hours, diluted with ethyl acetate, washed with water and brine, dried over sodium sulfate and concentrated *in vacuo*. This residue was redissolved in methanol (4 mL), CH<sub>3</sub>CN (3 mL) and 1N LiOH (0.7 mL). The reaction mixture was stirred for 1 hour, acidified with 1N HCl (0.7 mL), filtered and dried to yield methyl 1-[(7-cyano(2-naphthyl))methyl]-2-oxo-(3H)-benzimidazole-5-carboxylate (84 mg, 70%) as a white solid. ES-MS (M+H)<sup>+</sup>=358.1.


To a suspension of methyl 1-[(7-cyano-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxylate (82 mg, 0.23 mmol) in CH<sub>3</sub>CN (2.5 mL) and methanol (2 mL) was added 1N LiOH (2.3 mL, 10 eq.). The reaction mixture was stirred at room temperature for 3 days, acidified with 1N HCl (2.4 mL), filtered and dried to yield 1-[(7-cyano-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxylic acid (72 mg, 91%) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 5.16 (s, 2H); 6.77-6.79 (d, 1H); 7.50-7.54 (t, 2H); 7.62-7.64 (dd, 1H); 7.68 (s, 1H); 7.72 (s, 1H); 7.78-7.84 (dd, 2H); 8.11 (s, 1H); 10.8 (s, 1H). ES-MS (M+H)<sup>+</sup>=344.1.

Example 5

A solution of 1-[(7-cyano-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxylic acid (70 mg, 0.20 mmol) and hydroxylamine hydrochloride (28 mg, 0.40 mmol) in dry


ethanol (4 mL) and DIEA (0.11 mL, 0.61 mmol) was stirred at 55 °C for 8 hours. The reaction was concentrated *in vacuo*, and the resulting residue redissolved in glacial acetic acid (4 mL). To this solution was added acetic anhydride (0.038 mL, 0.40 mmol). The reaction mixture was stirred at room temperature for 1 hr, followed by addition of methanol (3 mL) and 10% Pd on carbon (24 mg, 0.023 mmol). The mixture was hydrogenated under 1 atm H<sub>2</sub> for 17 hours, filtered, and concentrated *in vacuo*. Purification on a Vydac C<sub>18</sub> HPLC column yielded 1-[(7-carboxamidino-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxylic acid (30 mg, 41%) as a white fluffy solid after lyophilization. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 5.29 (s, 2H); 7.17-7.19 (d, 1H); 7.56 (s, 1H); etc. ES-MS (M+H)<sup>+</sup>=361.1.

#### Example 6



To a suspension of 1-[(7-carboxamidino-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxylic acid (7 mg, 0.019 mmol) and BOP reagent (10 mg, 0.023 mmol) in DMF (0.8 mL) and N,N-diisopropylethylamine (10 μL, 0.057 mmol) was added neat pyrrolidine (2 μL, 0.024 mmol). The resulting solution was stirred at room temperature under argon for 4 hours. The reaction was concentrated and purified by HPLC to yield 7-{[2-oxo-5-(pyrrolidinylcarbonyl)-(3H)-benzimidazolyl]methyl}naphthalene-2-carboxamide (6 mg, 75%) as a white fluffy solid after lyophilization. ES-MS (M+H)<sup>+</sup>=414.1.


#### Example 7





Using a method similar method to that used in Example 6, 1-[(7-carboxamidino-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxamide was synthesized from the carboxylic acid via a BOP coupling with ammonium hydroxide. ES-MS (M+H)<sup>+</sup>=360.1.


#### 5 Example 8



Using a method similar to that used in Example 6, 7-[[5-(N-methylcarbamoyl)-2-oxo-(3H)-benzimidazolyl)methyl]naphthalene-2-carboxamidine was synthesized from the carboxylic acid via a BOP coupling with methylamine. ES-MS (M+H)<sup>+</sup>=374.1.

10


#### Example 9



Using a method similar to that used in Example 6, 7-[[5-(N,N-dimethylcarbamoyl)-2-oxo-(3H)-benzimidazolyl)methyl]naphthalene-2-carboxamidine was synthesized from the carboxylic acid via a BOP coupling with dimethylamine. ES-MS (M+H)<sup>+</sup>=388.1.

15

#### Example 10




20

Using a method similar to that used in Example 6, 3-{3-[2-oxo-5-(pyrrolidinylcarbonyl)-(3H)-benzimidazolyl]propoxy}benzamidinium was synthesized by alkylation with 3-(3-

bromopropoxy)benzenecarbonitrile, saponification, BOP coupling with pyrrolidine, and hydroxylamine method for conversion of nitrile to amidine. ES-MS  $(M+H)^+=408.1$ .

### Example 11




5

Using a method similar to that used in Example 6, 3-(2-methyl-5-{[2-oxo-5-(pyrrolidinylcarbonyl)-(3H)-benzimidazolyl]}methyl)-1,3-thiazol-4-yl)-benzamidinium was synthesized by alkylation with 3-[5-(chloromethyl)-2-methyl-1,3-thiazol-4-yl]benzenecarbonitrile, saponification, BOP coupling with pyrrolidine, and hydroxylamine method for conversion of nitrile to amidine. ES-MS  $(M+H)^+=461.0$ .

10


### Example 12



Using a method similar to that used in Example 6, 1-ethyl-2-{[2-oxo-5-(pyrrolidinylcarbonyl)-(3H)-benzimidazolyl]}methyl}indole-6-carboxamidinium was synthesized by alkylation with 2-(chloromethyl)-1-ethylindole-6-carbonitrile, saponification, BOP coupling with pyrrolidine, and hydroxylamine method for conversion of nitrile to amidine. ES-MS  $(M+H)^+=431.1$ .

15

### Example 13



20


Part 1. To a solution of 2*H*-1,4-Benzoxazin-3(4*H*)-one (1.86 g, 12.5 mmol, 1.0 equiv) in 25 mL of CHCl<sub>3</sub> at 0 °C was added Br<sub>2</sub> (2.0 g, 1.0 equiv) dropwise. After stirring at room temperature overnight, the solvent was evaporated and the residue was recrystallized from EtOH/H<sub>2</sub>O to give the product in 95% yield. LRMS found for C<sub>8</sub>H<sub>7</sub>BrNO<sub>2</sub> (M+H)<sup>+</sup>: 227.9.

Part 2. A solution of the product from Part 1 (103 mg, 0.4 mmol, 1.0 equiv), 2-(*t*-butylaminosulfonyl)phenylboronic acid (91.2 mg, 1.0 equiv), PdCl<sub>2</sub>(dppf) (32.6 mg, 0.1 equiv), and triethylamine (279 μL, 5.0 equiv) in 10 mL of DME was degassed with argon for 15 min, then heated to reflux overnight. After cooling to room temperature, the mixture was diluted with ethyl acetate, washed with water, dried over MgSO<sub>4</sub>, and concentrated. Flash chromatography on silica gel gave the product in 21% yield. LRMS found for C<sub>18</sub>H<sub>21</sub>N<sub>2</sub>O<sub>4</sub>S (M+H)<sup>+</sup>: 361.1.

Part 3: A solution of the product from Part 2 (36 mg, 0.1 mmol, 1.0 equiv) in 2 mL of DMF was treated with 2-bromomethyl-7-cyanonaphthalene (40 mg, 75%, 1.2 equiv) and Cs<sub>2</sub>CO<sub>3</sub> (65 mg, 2 equiv) for 0.5h. The mixture was diluted with ethyl acetate, washed with water and subjected to flash column chromatography on silica gel to give the desired product in 95% yield. LRMS found for C<sub>30</sub>H<sub>28</sub>N<sub>3</sub>O<sub>4</sub>S (M+H)<sup>+</sup>: 526.2.

Step 4: The compound obtained in Step 3 (50 mg, 0.1 mmol, 1.0 equiv) was dissolved in 5 mL of methanol. The reaction mixture was cooled to 0 °C and HCl gas was bubbled in until saturation, and the mixture was stirred at room temperature overnight. The solvent was evaporated and the resulting residue was treated with ammonium acetate and 10 mL of methanol at reflux temperature for 2 hr. The solvent was removed at reduced pressure and the crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give the desired salt in 80% yield. LRMS found for C<sub>26</sub>H<sub>23</sub>N<sub>4</sub>O<sub>4</sub>S (M+H)<sup>+</sup>: 487.1.


#### Example 14



Step 1: The product from Example 13, Step 2 (36 mg, 0.1 mmol, 1.0 equiv) in 2 mL of DMF was treated with 2-bromomethyl-2'-cyanobiphenyl (41 mg, 1.5 equiv) and  $\text{Cs}_2\text{CO}_3$  (65 mg, 2 equiv) for 0.5 hr. The mixture was diluted with ethyl acetate, washed with water and purified by flash column chromatography to give the desired product in 91% yield. LRMS found for  $\text{C}_{32}\text{H}_{30}\text{N}_3\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 552.2.

Step 2: The product from Step 1 (50 mg, 0.09 mmol, 1.0 equiv) was dissolved in 5 mL of methanol. The reaction mixture was cooled to 0 °C, HCl gas was bubbled in until saturation, and the mixture was stirred at room temperature overnight. The solvent was evaporated and the resulting residue was treated with ammonium acetate and 10 ml methanol at reflux temperature for 2 hr. The solvent was removed under reduced pressure and the crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  to give the desired salt in 77% yield. LRMS found for  $\text{C}_{28}\text{H}_{25}\text{N}_4\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 513.2.

#### Example 15 and 16



Part 1. A solution of 2-nitro-5-bromophenol (2.53 g, 18.2 mmol) in 15 mL of ethanol was treated with  $\text{SnCl}_2 \cdot \text{H}_2\text{O}$ , and the reaction was heated at reflux for 3h. The solvent was evaporated to give a white residue, which was used in the next step without further purification. LRMS:  $\text{C}_9\text{H}_{10}\text{NO}_3$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 180.1.


Part 2. A solution of 2-amino-5-bromophenol (2.53 g, 18.2 mmol, 1.0 equiv) in 15 mL of isobutylmethyl ketone and 15 mL of water was cooled to 0 °C,  $\text{NaHCO}_3$  (3.67 g, 2.4 equiv)

and then chloroacetyl chloride (2.36 g, 1.67 mL, 1.15 equiv) were added. The mixture was heated to reflux overnight, then cooled to room temperature. The mixture was diluted with ethyl acetate, washed with water, dried over  $\text{MgSO}_4$ , and concentrated *in vacuo*. Flash chromatography on silica gel gave 7-bromo-3,4-dihydro-2H-1,4-benzoxazin-3-one in 77% yield. LRMS found for  $\text{C}_9\text{H}_{10}\text{NO}_3$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 180.1.

Part 3. A solution of 7-bromo-3,4-dihydro-2H-1,4-benzoxazin-3-one (103 mg, 0.4 mmol, 1.0 equiv), 2-t-butylaminosulfonyl phenyl boronic acid (91.2 mg, 1.0 equiv),  $\text{PdCl}_2(\text{dppf})$  (32.6 mg, 0.1 equiv), and triethylamine (279  $\mu\text{L}$ , 5.0 equiv) in 10 mL of DME was degassed with argon for 15 min, then heated to reflux overnight. After cooling to room temperature, the mixture was diluted with ethyl acetate, washed with water, dried over  $\text{MgSO}_4$ , evaporated. Flash chromatography on silica gel gave the product in 21% yield. LRMS found for  $\text{C}_{18}\text{H}_{21}\text{N}_2\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 361.1.


Part 4. The product from Part 3 (36 mg, 0.1 mmol, 1.0 equiv) in 2 mL DMF was treated with 1-(3-cyanophenyl)-3-methyl-5-chloromethyl pyrazole (35 mg, 1.5 equiv) and  $\text{Cs}_2\text{CO}_3$  (65 mg, 2 equiv) for 0.5 hr. The mixture was diluted with ethyl acetate, washed with water and purified over silica gel to give the desired product in 91% yield. LRMS found for  $\text{C}_{30}\text{H}_{29}\text{N}_5\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 556.1.

Part 5. The compound from Part 4 (50 mg, 0.09 mmol, 1.0 equiv) was dissolved in 5 mL of methanol. The reaction mixture was cooled to 0 °C, HCl gas was bubbled in until saturation, and the mixture was stirred at room temperature overnight. The solvent was evaporated and the resulting residue was treated with ammonium acetate and 10 mL of methanol at reflux temperature for 2 hr. The reaction was concentrated *in vacuo* and the crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  to give the desired amidine (Example 15) in 77% yield. LRMS found for  $\text{C}_{26}\text{H}_{24}\text{N}_6\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 516.2. Example 16 was obtained as a byproduct in 15% yield. LRMS found for  $\text{C}_{26}\text{H}_{23}\text{N}_5\text{O}_5\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 517.1.

Example 17

Part 1: The product (36 mg, 0.1 mmol, 1.0 equiv) obtained in Part 3 of Example 15 in 2 mL DMF was treated with 1-(3-cyanophenyl)-3-methyl-5-chloromethyl pyrazole (41 mg, 1.5 equiv) and  $\text{Cs}_2\text{CO}_3$  65 mg, 2 equiv) for 0.5h. The mixture was diluted with ethyl acetate, washed with water and purified over silica gel to give the desired product in 91% yield. LRMS found for  $\text{C}_{32}\text{H}_{30}\text{N}_3\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 552.2.

Part 2: The compound obtained in Part 1 (50 mg, 0.09 mmol, 1.0 equiv) was dissolved in 5 mL of methanol. The reaction mixture was cooled to 0 °C, HCl gas was bubbled in until saturation, and the mixture was stirred at room temperature overnight. The solvent was evaporated and the resulting residue was treated with ammonium acetate and 10 ml of methanol at reflux temperature for 2 hr. The solvent was removed at reduced pressure and the crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  to give the desired salt in 77% yield. LRMS found for  $\text{C}_{28}\text{H}_{25}\text{N}_4\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 513.2.


Example 18

To a solution of tert-butylamine (41.4g, 566 mmol) and triethylamine (118 mL, 849 mmol) in DCM (1000 mL) in an ice bath, was added benzenesulfonyl chloride (100 g, 566 mmol) dropwise, and the mixture was stirred at room temperature overnight. Water was added to the mixture and organic layer was washed with water, sat. NaCl, dried over  $\text{Na}_2\text{SO}_4$ ,




filtered and filtrate evaporated *in vacuo* to give the title compound as light yellow solid (117.63 g, 97.6%). ES-MS (M+H)<sup>+</sup> = 214.5.

#### Example 19



A solution of the compound from example 18 (53.25 g, 250 mmol) in THF (600 mL) was cooled with an ice water bath, and a 2.5 M solution of n-butyllithium in hexane (200 mL, 500 mmol) was added dropwise. A thick precipitate was formed when the reaction mixture was warmed up to 10 °C. Triisopropylborate was added, keeping the temperature below 35 °C. After 1 hr., the mixture was cooled in an ice bath, 1N HCl (405 mL) was added, and the mixture was stirred overnight. The mixture was extracted with ether (3 x 100 mL), and the combined organic extracts were extracted with 1N NaOH (3 x 130 mL). The aqueous extracts were acidified to pH 1 with 12 N HCl, and then extracted with ether (3 x 140 mL). The combined ether extracts were dried over MgSO<sub>4</sub>, and solvents were evaporated *in vacuo*. Hexane and ether were added and a white precipitate formed. The solid was collected and washed with 10% ether/hexane to give the title compound. ES-MS (M+H)<sup>+</sup> = 257.8.


#### Example 20



To a solution of 5-bromoindole (1.96 g, 10 mmol) in DME (40 mL) and H<sub>2</sub>O (10 mL), was added the compound from Example 19 (3.85 g, 15 mmol), NaHCO<sub>3</sub> (1.68g, 20 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.58g, 0.5 mmol). The mixture was heated to reflux overnight, then cooled to room temperature and diluted with ethyl acetate. The organic layer was washed with water, dried with MgSO<sub>4</sub>, filtered and concentrated. The


residue was purified by silica gel column chromatography using 25% ethyl acetate in hexane as eluent to give the title compound (1.52g, 46%). ES-MS (M+Na)<sup>+</sup> 351.1.

### Example 21



To a solution of the compound from Example 20 (328 mg, 1 mmol) in DMF (10 mL) was added 7-cyano-2-bromomethylnaphthalene (296 mg, 1.2 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (1.3g, 4 mmol), and the mixture was stirred at room temperature overnight. The reaction mixture was partitioned between water and ethyl acetate, and the organic layer was washed with water, 1N HCl, sat. NaCl, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The residue was subjected to flash column chromatography on silica gel using 50% ethyl acetate in CH<sub>2</sub>Cl<sub>2</sub> followed by 100% CH<sub>2</sub>Cl<sub>2</sub> as eluent to give the title compound (237 mg, 48%). ES-MS (M+H)<sup>+</sup> = 494.2.

### Example 22




A solution of the compound from example 21 (115 mg, 0.23 mmol) in 20%MeOH/ethyl acetate (10 mL) was treated with a stream of HCl gas for 10 min. at 0 °C. The resulting solution was capped, stirred at room temperature overnight, then concentrated *in vacuo*. The residue was reconstituted in MeOH (10 mL) and the mixture was treated with ammonium acetate (350 mg, 4.6 mmol). The reaction mixture was heated at reflux for 2



hrs, then concentrated *in vacuo*. The residue was purified by prep HPLC to give the title compound as a white powder. ES-MS  $(M+H)^+ = 511.2$ .

### Example 23



Part 1. A solution of 4-bromo-2-nitroaniline (2.97 g, 13.7 mmol) in 100 mL of dichloromethane was treated with N,N-diisopropylethylamine (2.38 mL, 13.7 mmol) and 4-dimethylaminopyridine (167 mg, 1.37 mmol). Di-tert-butyl dicarbonate (3.88 g, 17.8 mmol) was then added in small portions. The mixture was stirred at room temperature overnight under argon, then diluted with 400 mL of dichloromethane and washed with water (x 2). The organic phase was dried over  $MgSO_4$ , concentrated *in vacuo* and purified by flash column chromatography. The desired product was obtained as a yellow solid in 48% yield (2.07 g).


Part 2. To a mixture of the product from Part 1 (1.02 g, 3.22 mmol), the boronic acid from Example 19 (827 mg, 3.22 mmol), and tetrakis(triphenylphosphine)palladium(0) (186 mg, 0.161 mmol) in 60 mL of benzene was added a solution of sodium hydroxide (515 mg, 12.9 mmol) in 5 mL of water. The mixture was degassed using an argon stream for 15 minutes and then heated at reflux for 24 hours. The solution was concentrated *in vacuo*, the residue was dissolved in ethyl acetate, passed through a bed of celite, and washed with water (x 2). The organic phase was dried over  $MgSO_4$ , concentrated *in vacuo*, then purified by flash column chromatography. The desired product was obtained as a solid in 60% yield (0.87 g).

Part 3. The product from Part 2 (72 mg, 0.16 mmol) and the bromide from Example 24 (80mg, 0.24 mmol) were dissolved in 4 mL of dry DMF, and cesium carbonate (156 mg, 0.48 mmol) was added. The resulting mixture was stirred overnight, then diluted with 100 mL of ethyl ether, washed with water (x 2), and dried over MgSO<sub>4</sub>. Filtration and concentration *in vacuo* gave a residue, which was subjected to flash column chromatography. The desired product was obtained in 43% yield (48 mg).

Part 4. The product from Part 3 (90 mg, 0.13 mmol) was dissolved in 5 mL of ethyl alcohol, and tin(II) chloride dihydrate (116 mg, 0.52 mmol) was added. The mixture was heated to reflux for 3 hours, and the ethanol was removed *in vacuo*. The residue was dissolved in ethyl acetate, washed with 1N aqueous NaOH and water. The organic phase was dried over MgSO<sub>4</sub>, concentrated *in vacuo* and pumped to dryness to give the desired product in quantitative yield (84 mg).

Part 5. The crude product from Part 4 (84 mg) was dissolved in 7 mL glacial acetic acid, and the mixture was stirred at 80 °C overnight. The mixture was concentrated with toluene to remove traces of acetic acid, and the residue was purified by preparative HPLC.


Part 6. The product from Part 5 (20 mg, 0.034 mmol) was dissolved in 5 mL anhydrous methanol, and a vacuum-distilling adapter, equipped with a rubber septum and with a balloon on its side-arm, was placed on the reaction flask. The solution was chilled in ice, and HCl gas was bubbled through the solution via a long needle immersed in the solution until the balloon expanded, and this reaction mixture was stirred overnight. The reaction was concentrated to dryness *in vacuo*, then dissolved in 5 mL of anhydrous methanol, and dry ammonium acetate (21 mg, 0.27 mmol) was added. The mixture was heated at reflux for 2 hours, then purified by preparative HPLC to afford the title compound. LRMS (M+H)<sup>+</sup> = 553.

Example 24

Part 1. A mixture of 1,1,1-trifluoro-2,4-pentanedione (5.43 mL, 44.7 mmol) and 3-bromophenylhydrazine HCl (10.0 g, 44.7 mmol) in 100 mL of ethanol was heated to reflux overnight. The reaction mixture was concentrated *in vacuo*, and the residue was dissolved in ethyl ether and washed with water (x 2). The organic phase was dried over MgSO<sub>4</sub>, concentrated and subjected to flash column chromatography, giving 12.83 g (94%) of the desired product as an oil, which was a mixture of the two possible regioisomers.

Part 2. The mixture of isomers from part 1 (12.83 g, 41.9 mmol), KCN (5.45 g, 83.8 mmol), CuI (798 mg, 4.19 mmol) and tetrakis(triphenylphosphine)palladium(0) (2.42 g, 2.09 mmol) in 100 mL of acetonitrile was degassed using an argon stream for 30 minutes. The mixture was heated at reflux for 15 hours, then passed through a silica plug on a Buchner funnel using ethyl acetate as eluent. The filtrate was concentrated and dissolved in dichloromethane, washed with water (x 2), and dried over MgSO<sub>4</sub>. Filtration and concentration gave a residue, which was subjected to column chromatography. The regioisomeric products were separable at this stage, giving 3.93 g (37%) of the 3-methyl and 4.0 g (38%) of the 5-methyl isomers.


Part 3. A mixture of the 5-methyl isomer (1.67 g, 6.63 mmol), NBS (1.41 g, 7.95 mmol), and AIBN (543 mg, 3.31 mmol) in 50 mL carbon tetrachloride was degassed using an argon stream for 15 minutes. The reaction was heated at reflux for 2 days, periodically adding additional small amounts of AIBN. The resulting mixture was diluted with dichloromethane, washed with water (x 2), and dried over MgSO<sub>4</sub>. Filtration and concentration gave a residue, which was subjected to flash column chromatography. The desired bromomethyl product was obtained in 1.06 g (48%), along with 1.06 g of recovered starting material.

Examples 25 and 26

Part 1. The nitro compound from Part 2 of Example 23 (2.70 g, 6.01 mmol) was dissolved in 60 mL of ethanol, and tin(II) chloride dihydrate (5.43 g, 24.0 mmol) was added. The mixture was heated at reflux for 1 hour, then concentrated on the rotovap. The residue was washed through a plug of silica gel and concentrated to give another residue, which was dissolved in 20 mL of formic acid (88%) and stirred at 55 °C for 1 hour. Traces of formic acid were removed by evaporation with toluene *in vacuo*, and the desired product was obtained by flash column chromatography to give 516 mg (26%) of the desired benzimidazole. LRMS (M+H)<sup>+</sup> = 330.

Part 2. The benzimidazole from Part 1 (62 mg, 0.188 mmol) was dissolved in 5 mL of DMF, followed by the addition of 2-bromomethyl-7-cyanonaphthalene (93 mg, 0.376 mmol) and cesium carbonate (184 mg, 0.564 mmol). The mixture was stirred overnight, then diluted with ethyl acetate and washed with water (x 2). The organic phase was dried over MgSO<sub>4</sub>, concentrated *in vacuo*, and the desired products were obtained as a mixture of two regioisomers by flash column chromatography.


Part 3. The mixture of isomers from Part 2 (90 mg, 0.18 mmol) was dissolved in 10 mL of dry methanol. A vacuum-distilling adapter, equipped with a rubber septum and with a balloon on its side-arm, was placed on the reaction flask. The solution was chilled in ice, and HCl gas was introduced via a long needle immersed in the solution until the balloon began to inflate. The mixture was stirred overnight and then concentrated to dryness *in vacuo*. The residue was dissolved in 5 mL anhydrous methanol, and dry ammonium acetate (139 mg, 1.8 mmol) was added. The mixture was heated at reflux for 2 hours, then purified by preparative HPLC to afford the two title compounds. LRMS (M+H)<sup>+</sup> = 457.

Example 27

Part 1. To a solution of 5-bromoindoline (1.0g, 5mmol) in 10ml of dioxane was added 5 mL of a 1N NaOH solution and 5 mL of water. This mixture was cooled with an ice bath, and di-tert-butyl dicarbonate (1.2g, 5.5mmol) was added in one portion. The reaction was allowed to warm to room temperature, stirred for 4 hours, and then concentrated. The residue was extracted with ethyl acetate (2 x 25 mL), and the combined organic phases were washed with water (2 x 25mL), saturated aqueous NaCl (2 x 25ml), then dried over MgSO<sub>4</sub>. Filtration and concentration *in vacuo* gave N-Boc-5-bromoindoline (1.33g, 89%) as a light brown powder after drying. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.541 (s, 9H); 3.03-3.08 (t, 2H); 3.92-3.96 (t, 2H); 7.23-7.27 (m, 3H).

Part 2. To a solution of N-Boc-5-bromoindoline (104 mg, 0.35 mmol) in 5ml of anhydrous dioxane were added the boronic acid from Example 19 (100 mg, 0.39 mmol), cesium carbonate (228mg, 0.7mmol) and tris(dibenzylideneacetone)dipalladium(0)-chloroform adduct (10 mg, 0.01 mmol). The reaction flask was thoroughly flushed with argon, and tri-*tert*-butylphosphine (0.006 mL, 0.025 mmol) was added via syringe. The reaction was heated to 80 °C and stirred overnight. The reaction was then cooled to room temperature, diluted with 25 mL of diethyl ether, flushed through a pad of Celite and concentrated *in vacuo* to give a brown residue. This residue was subjected to flash column chromatography on silica gel using 25% ethyl acetate in hexane to give the desired biphenylamine (150 mg, 100%) as a light brown oil after drying. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.02 (s, 9H); 1.22 (s, 9H); 3.10-3.15 (t, 2H); 4.02, (t, 2H); 7.25-8.14 (m, 7H).

Part 3. The product from Part 2 was treated with 10ml of 20% trifluoroacetic acid in dichloromethane for 10 minutes. The reaction mixture was concentrated *in vacuo*, redissolved in 25 mL of dichloromethane, washed with 1N NaOH (2x25ml), dried over magnesium sulfate, filtered and concentrated *in vacuo* to give the desired biphenylamine as a brown oil after drying. ES-MS (M+H<sup>+</sup>): 330.5

Example 28

Part 1. 3-Cyano-4-fluoronitrobenzene (9.50 g, 57 mmol) was dissolved in 250 mL of ethanol, 5% Pd/C (2.0 g) was added, and the black slurry was stirred under a hydrogen balloon for 2 days. The reaction mixture was filtered through a celite bed and concentrated *in vacuo* to give the amino compound in 95% yield (7.37 g). LRMS (M+H)<sup>+</sup> m/z 137. R<sub>f</sub> 0.62 (2:1 ethyl acetate/hexane).

Part 2. The amino compound from Part 1 (4.00 g, 29 mmol) was slurried in 28 mL of conc. HCl and chilled in an ice bath. Sodium nitrite (2.00 g, 29 mmol) was dissolved in 10 mL of water and chilled in ice bath, and this solution was added dropwise to the cold slurry of the amino compound. The reaction mixture was then stirred at 0 °C for 30 min. Tin chloride dihydrate (19.7 g, 29 mmol) was dissolved in 10 mL of conc. HCl and chilled in ice bath, and this solution was added dropwise to the previous reaction mixture. After completion, the mixture was chilled and the solid hydrazine product was isolated by filtration through a Buchner funnel. This material was washed with cold brine (60 mL) and cold hexane (60 mL) to give 5.85 g of a yellow solid, which was used in the next step without further purification.

Part 3. Ethyl 2,4-dioxovalerate (20.0 g, 127 mmol) was dissolved in 100 mL of ethanol, and methoxylamine hydrochloride (11.1 g, 133 mmol) was added. The mixture was heated at reflux for 20 hours, followed by concentration of the reaction mixture *in vacuo*. The residue was dissolved in ethyl acetate, and washed with water and brine. The separated




organic phase was dried over  $\text{MgSO}_4$  and evaporated *in vacuo* to give 19.3 g (81%) of the desired imine as an oil. LRMS  $(\text{M}+\text{H})^+$   $m/z$  188.


Part 4. A solution of the product from Part 2 (5.85 g) and the product from Part 3 (3.80 g, 20.3 mmol) in acetic acid (100 mL) and THF (50 mL) was heated at reflux overnight, and the reaction mixture was concentrated *in vacuo*. The residue was taken up in ethyl acetate and washed with water and brine. The organic phase was dried, filtered, and concentrated and the residue was subjected to flash column chromatography to yield the desired 5-carboethoxy isomer (4.45 g, 56% for two steps) and the undesired 3-carboethoxy isomer (1.00 g, 12% over two steps). LRMS  $(\text{M}+\text{H})^+$   $m/z$  274. 5-Carbomethoxy isomer  $R_f$  0.66 (1:1 ethyl acetate/hexane). 3-Carbomethoxy isomer  $R_f$  0.50 (1:1 ethyl acetate/hexane).

Part 5. The biphenylamine from Example 27, Part 3 (90 mg, 0.27 mmol) was dissolved in 2 mL of dichloromethane, and trimethylaluminum (0.7 mL of a 2.0 M solution in hexane, 1.35 mmol) was added at room temperature. The mixture was stirred for 30 minutes, and then a solution of the 5-carbomethoxy ester from Part 4 (77 mg, 0.28 mmol) in 3 mL of dichloromethane was added dropwise to the reaction. The resulting mixture was stirred under argon overnight. The reaction was then quenched by the careful addition of aqueous sat. Rochelle's salt, and the reaction was extracted with dichloromethane (x 3). The organic layer was concentrated and the residue was subjected to flash column chromatography to afford the desired coupling product (90 mg, 60%). LRMS  $(\text{M}+\text{H})^+$   $m/z$  558.  $R_f$  0.29 (1:1 ethyl acetate/hexane).

Part 6. The product from part 5 (83 mg, 0.15 mmol) was dissolved in 2 mL of ethanol, hydrazine monohydrate (40  $\mu\text{L}$ , 0.75 mmol) was added dropwise, and the reaction mixture was heated at reflux overnight. Concentration *in vacuo* gave a residue, which was dissolved in 2 mL of trifluoroacetic acid and heated at reflux for 30 minutes. The resulting mixture was concentrated and purified by prep HPLC [(C18 column, standard  $\text{H}_2\text{O}$  to  $\text{CH}_3\text{CN}$  gradient (0.1% TFA)] to give the desired compound. LRMS  $(\text{M}+\text{H})^+$   $m/z$  514.

Example 29

This compound was prepared from the 3-carbomethoxy intermediate from Part 4 of Example 28 in a manner similar to that in Parts 4 and 5, Example 28.

Example 30

Part 1. 3-Bromophenylhydrazine hydrochloride (6.80 g, 30.4 mmol) and the product from Example 28, Part 3 (2.84 g, 15.2 mmol) were dissolved in 60 mL of acetic acid and 30 mL of THF, and the reaction mixture was heated at reflux overnight. The solvent was removed *in vacuo*, the residue was dissolved in ethyl acetate, and the solution was washed with water and sat. NaCl. The organic layer was dried over MgSO<sub>4</sub>, evaporated *in vacuo*, and purified by flash column chromatography to give the desired pyrazole in quantitative yield. LRMS (M+H)<sup>+</sup> m/z 309, 311. R<sub>f</sub> 0.47 (1:4 ethyl acetate/hexane).

Part 2. The product from Part 1 was dissolved in 60 mL of acetonitrile, and KCN (2.91 g, 44.8 mmol), CuI (0.43 g, 2.2 mmol) and Pd(Ph<sub>3</sub>P)<sub>4</sub> (1.3 g, 1.1 mmol) were added. The mixture was degassed by bubbling argon through the solution for 30 minutes, and the reaction mixture was heated under argon at reflux overnight. The black mixture was then passed through a silica plug, and the filtrate was concentrated to give a residue, which was




taken up in ethyl acetate and washed with water (x 2). The organic layer was dried, concentrated and subjected to flash chromatography to yield the desired nitrile as a white solid (2.69 g, 10.5 mmol, 69% for two steps). LRMS (M+H)<sup>+</sup> m/z 256. R<sub>f</sub> 0.68 (1:1 ethyl acetate/hexane).

Part 3. The product from Part 2 (2.00 g, 7.8 mmol) was dissolved in 40 mL of methanol, and water (20 mL) and LiOH monohydrate (0.66 g, 15.7 mmol) were added. The mixture was stirred for 1 hour, then acidified with 5N HCl to a pH of 1. The methanol was removed *in vacuo*, and the mixture was extracted with ethyl acetate (x 3). The combined organic phases were dried and evaporated to give the desired acid as a white solid (1.38 g, 78%). LRMS (M+H)<sup>+</sup> m/z 228, (M+Na)<sup>+</sup> m/z 250.

Part 4. The acid from part 3 (200 mg, 0.88 mmol), the biphenyl amine from Example 25, Part 3 (290 mg, 0.88 mmol), and 4,4-dimethylaminopyridine (10 mg) were dissolved in 4 mL of pyridine and chilled in ice bath. To the mixture was added POCl<sub>3</sub> (0.25 mL, 2.6 mmol), and the reaction was stirred in the cold for 1 hour. The reaction was quenched with ice water, diluted with ethyl acetate, and the organic phase was washed with sat. NaCl (x 2). The organic layer was dried, concentrated, and subjected to flash column chromatography to yield the desired amide (240 mg, 51%). LRMS (M+H)<sup>+</sup> m/z 540. R<sub>f</sub> 0.37 (1:1 ethyl acetate/hexane).

Part 5. The product from Part 4 (140 mg, 0.26 mmol) was dissolved in 10 mL of anhydrous DMF and chilled in ice bath. To the reaction mixture was added NaBH<sub>4</sub> (80 mg, 2.08 mmol) and anhydrous CoCl<sub>2</sub> beads (68 mg, 0.52 mmol). The mixture was stirred for 30 minutes in the cold, then quenched with ice water and diluted with ethyl acetate. The mixture was filtered through a pad of celite, and the filtrate was washed with sat. NaCl (x 2). The organic phase was dried, and concentrated *in vacuo* to give a residue, which was dissolved in 2 mL of trifluoroacetic acid and stirred at 60 °C for 30 minutes. The mixture was purified by prep HPLC [(C18 column, standard H<sub>2</sub>O to CH<sub>3</sub>CN gradient (0.1% TFA)] to afford the desired aminomethyl compound. LRMS (M+H)<sup>+</sup> m/z 488.

Examples 31 and 32


Part 1. A solution of the product from Example 27, Part 2 (80 mg, 0.2 mmol) in 4 mL of CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 mL of trifluoroacetic acid, and the solution was stirred at room temperature for 10 min. The reaction mixture was then concentrated, and again concentrated from heptane to remove traces of trifluoroacetic acid. The residue was dissolved in 5 mL of DMF, and the bromomethyl compound from Example 24, Part 3 (80 mg, 0.24 mmol) was added, together with 195 mg of cesium carbonate, and the solution was stirred at room temperature for 1 hr. The reaction was diluted with ethyl acetate (25 mL), washed with water and sat. NaCl, and the organic layer was dried over MgSO<sub>4</sub>. Filtration and concentration gave a brown oil, which was used in the next reaction without further purification.

Part 2. The material from Part 1 (0.2 mmol) was taken up in 10 mL of absolute ethanol, and hydroxylamine hydrochloride (164 mg, 1.5 mmol) and triethylamine (0.2 mL, 1.5 mmol) were added. The solution was heated to 60 °C for 5 hours, then concentrated to give a residue, which was taken up in 5 mL of acetic acid, and acetic anhydride (0.2 mL) was added. The reaction mixture was stirred at room temperature for 30 min, then concentrated to dryness, followed by concentration with heptane to remove traces of acetic acid. The residue was used in the next reaction without further purification.

Part 3. The crude product from Part 2 was taken up in 5 mL of absolute ethanol, 1 drop of acetic acid was added, followed by 10% Pd/C (5 mg), and the reaction was placed under a balloon of hydrogen. After 4 hours, the starting material was gone by HPLC, so the reaction mixture was filtered and subjected directly to preparative HPLC [C18 column, standard H<sub>2</sub>O to CH<sub>3</sub>CN (0.1% TFA) gradient]. Two products were isolated, the expected


indoline product, and a product where the indoline ring had been oxidized to an indole ring. Indoline product: LRMS (M+H)<sup>+</sup> m/z 541.1. Indole product: LRMS (M+H)<sup>+</sup> m/z 539.1.

5 Example 33




A solution of the product from Example 30, Part 4 (75 mg, 0.19 mmol) was treated with methanol and HCl, followed by ammonium acetate in a manner similar to that in Example 23, Part 6. Preparative HPLC gave the desired product.


10

Example 34

Treatment of the product from Example 30, Part 4 with hydroxylamine hydrochloride and triethylamine gave the above amidoxime.


Example 35

A solution of 4-bromoaniline (15 g, 87 mmol) and  $K_2CO_3$  (181 g) in 170 mL of water was cooled to 5 °C. A solution of cinnamoyl chloride (18.2 g, 109 mmol) in acetone (87 mL) was added dropwise, resulting in the formation of a light brown solid. After 1 hour, the solid was recovered by filtration, washed with cold water, once with cold ether and dried overnight at 80 °C at 20 mm Hg to give the desired product (23.8 g, 90%). A portion of this amide (10 g, 33 mmol) was suspended in chlorobenzene (220 mL) and aluminum chloride (26 g, 195 mmol) was added portionwise. The solution was heated at reflux for 2 hr, the solution was concentrated *in vacuo* to one half of its original volume, and the solution was poured into 1L of ice water. The resulting solid was collected by filtration and recrystallized from methanol to give 6-bromoquinolin-2-one (5.0 g, 71%) as a brown solid.

Example 36


A solution of 6-bromohydroquinolin-2-one (1 g, 4.46 mmol), 2-bromomethyl-7-naphthonitrile (1.16 g, 4.46 mmol), and cesium carbonate (1.5 g, 5.35 mmol) in 15 mL of DMF was stirred at room temperature for 18 h. The reaction mixture was diluted with water and ethyl acetate and filtered through celite. The organic layer was washed with water (x 2), then with sat. NaCl, dried over  $\text{MgSO}_4$ , and concentrated *in vacuo*. The residue was subjected to flash column chromatography on silica gel, using 100%  $\text{CH}_2\text{Cl}_2$  as eluent to give 7-[(6-bromo-2-oxohydroquinolyl)methyl]naphthalene-2-carbonitrile (876 mg, 50%) as a white solid. ES-MS  $(\text{M}+\text{H})^+ = 389, 391$ .

### Example 37



To a solution of 7-[(6-bromo-2-oxohydroquinolyl)methyl]naphthalene-2-carbonitrile (30 mg, 0.077 mmol), 2-[[[(tert-butyl)amino]sulfonyl}phenyl]boronic acid (20 mg, 0.077 mmol),  $\text{Pd}_2(\text{dba})_3$  (1 mg, 1.5 mol%), and cesium carbonate (25 mg, 0.093 mmol) in dry dioxane (300  $\mu\text{L}$ ) was added tri-*t*-butylphosphine (0.7  $\mu\text{L}$ , 3.6 mol%). The reaction was heated to 75  $^\circ\text{C}$  for 11 h, cooled to room temperature, and diluted with  $\text{CH}_2\text{Cl}_2$  (2 mL). The solution was then filtered through celite, washed with 1N HCl and sat. NaCl, dried over  $\text{MgSO}_4$ , and concentrated *in vacuo* to afford 7-{[6-(2-[[[(tert-butyl)amino]sulfonyl}phenyl)-2-oxohydroquinolyl]methyl]naphthalene-2-carbonitrile (34 mg, 85%). ES-MS  $(\text{M}+\text{H})^+ = 522.2$ .


### Example 38



A solution of the nitrile from Example 37 (34 mg, 0.64 mmol) in ethyl acetate (5mL) containing MeOH (150  $\mu\text{L}$ ) was cooled to -78  $^\circ\text{C}$  and HCl gas was bubbled in until


saturation. The solution was stirred while warming to room temperature over 18 h. The solvent was removed *in vacuo*, the residue was taken up in methanol (2 mL), and dried.  $\text{NH}_4\text{OAc}$  (50 mg, 0.64 mmol) was added. The mixture was heated to 80 °C for 1h, cooled, and purified by preparative HPLC [(C18 column, water/ $\text{CH}_3\text{CN}$  gradient (0.1% TFA))]. The appropriate fractions were combined and lyophilized to give the desired amidine (5.6 mg, 19%) as a white powder. LRMS  $(\text{M}+\text{H})^+$   $m/z$  484.

#### Example 39



A solution of the nitrile from Example 37 (53 mg, 0.10 mmol) in methanol (0.5 mL) and DMF (0.25 mL) was treated with hydroxylamine hydrochloride (97 mg, 0.20 mmol), *N,N*-diisopropylethylamine (71  $\mu\text{L}$ , 0.20 mmol). The solution was heated to 40 °C for 18h, then concentrated *in vacuo*. The residue was stirred with TFA (2 mL) for 1h, then concentrated *in vacuo*, and this residue was purified by preparative HPLC [(C18 column, water/ $\text{CH}_3\text{CN}$  gradient (0.1% TFA))]. The appropriate fractions were combined and lyophilized to give the desired amidoxime (17 mg, 32%) as a white powder. LRMS  $(\text{M}+\text{H})^+$   $m/z$  499.

#### Example 40



Part 1. 7-Bromoisoquinoline

This compound was prepared as a 60:40 mixture with 5-bromoisoquinoline as in *J. Am. Chem. Soc.*, **1939**, *61*, 183.

5    Part 2. 7-Bromoisoquinoline N-oxide hydrochloride

This compound was prepared by a procedure analogous to that for 6-bromoisoquinoline N-oxide hydrochloride as in PCT WO 98/47876. A solution of 7.8 g (37.5 mmol) of a 60:40 mixture of 7-bromo and 5-bromoisoquinoline in 125 mL of CH<sub>2</sub>Cl<sub>2</sub> was treated portionwise with 9.7 g (~39.4 mmol) of 3-chloroperoxybenzoic acid (~70% purity). The  
10    solution, which was initially homogeneous, deposited a voluminous precipitate over 1 hr. Then 100 mL of methanol were added, and the reaction was concentrated to a volume of about 100 mL. Gaseous HCl was then bubbled through the solution for about 10 min, during which time the solution became warm and all of the precipitate dissolved. A few minutes later, another voluminous precipitate began to form. To this solution was added  
15    100 mL of ether, and the mixture was stirred in an ice-water bath for 20 minutes. The resulting product was isolated by filtration, washed thoroughly with ether, and air-dried to give 8.07 g (83%) of the desired compound as a white solid, which was still a 60:40 mixture of the 7- and 5-bromo isomers.

20    Part 3. 7-Bromo-1-chloroisoquinoline

This compound was prepared by a procedure analogous to that for 6-bromo-1-chloroisoquinoline as in PCT WO 98/47876. A solution of 8.07 g (31 mmol) of the mixture from Part B was taken up in 50 mL of POCl<sub>3</sub>, and the mixture was heated at 90 °C for 2 hr. The reaction mixture was concentrated to remove most of the POCl<sub>3</sub>, and the  
25    residue was taken up in 100 mL of CH<sub>2</sub>Cl<sub>2</sub>. The solution was carefully basified to pH 10 by the slow addition of 1N NaOH, and the organic layer was washed with 100 mL of H<sub>2</sub>O, 100 mL of sat. NaCl, and dried over MgSO<sub>4</sub>. Filtration and concentration gave a light yellow solid, which was subjected to flash column chromatography on silica gel first with 5% and then with 10% ethyl acetate in hexanes. A total of 3.62 g (48%) of the desired 7-  
30    bromo-1-chloroisoquinoline was isolated from this chromatography free of the 5-bromo isomer.



Part 4. 7-Bromo-1-phenoxyisoquinoline

A solution of 3.60 g (14.8 mmol) of 7-bromo-1-chloroisoquinoline and 1.5 g of solid KOH in 11.2 g of phenol was heated at 140 °C for 2 hr. The reaction was cooled to room temperature, then partitioned between 100 mL of CH<sub>2</sub>Cl<sub>2</sub> and 100 mL of 3N NaOH. The organic layer was washed with another 2 x 100 mL of 3N NaOH, then with 100 mL of H<sub>2</sub>O, and dried over MgSO<sub>4</sub>. Filtration and concentration gave a yellow oil, which was subjected to flash column chromatography on silica gel 30% CH<sub>2</sub>Cl<sub>2</sub> in hexanes, giving 3.42 g (77%) of the desired product as a light yellow solid.

Part 5. 1-Amino-7-bromoisoquinoline

A mixture of 3.40 g (11.3 mmol) of 1-amino-7-bromoisoquinoline and 7.65 g of ammonium acetate was heated at 150 °C for 15 hr. The reaction was cooled, and the residue was partitioned between 200 mL of ethyl acetate and 200 mL of 3N NaOH. The organic layer was extracted with 2 x 100 mL of 2N HCl, and the combined aqueous extracts were basified to pH 10 using 50% NaOH. This solution was extracted with 2 x 100 mL of ethyl acetate, and the organics were then washed with 100 mL of sat. NaCl and dried over MgSO<sub>4</sub>. Filtration and concentration gave 1.68 g (66%) of the desired amino compound as a yellow solid.

Part 6. 1-[Bis(t-butoxycarbonyl)amino]-7-bromoisoquinoline

A solution of 740 mg (3.32 mmol) of 1-amino-7-bromoisoquinoline in 50 mL of acetonitrile was treated with 1.4 mL of N,N-diisopropylethylamine and 100 mg of 4-(N,N-dimethylamino)pyridine, followed by 3.0 g (4.1 eq) of di-t-butyldicarbonate, and the reaction was stirred at 40 °C for 1 hr. By HPLC analysis, there was still some starting amino compound that remained, so another 1.0 g of di-t-butyldicarbonate were added, and the reaction was stirred at 40 °C for another 30 min. The reaction mixture was concentrated to give a dark oil, which was subjected to flash column chromatography on silica gel with 20% ethyl acetate in hexanes to give 736 mg of the desired product as a light yellow solid. Also isolated were 156 mg of product as a somewhat less pure light yellow solid, making the total yield 64%.



## Part 7. 1-[Bis(t-butoxycarbonyl)amino]isoquinoline-7-carboxaldehyde

A solution of 400 mg (0.95 mmol) of 1-[bis(t-butoxycarbonyl)amino]-7-bromoisquinoline in 50 mL of anhydrous THF was cooled with a liquid nitrogen/methanol slush bath (-98 °C), and 0.55 mL of a 2.43 M solution of n-BuLi in  
5 hexanes (1.3 eq) was added dropwise over 1 min. The solution was stirred in the cold for 5 min, then a solution of 5 mL of anhydrous DMF in 10 mL of anhydrous THF was added rapidly. The solution was allowed to warm to about 0 °C, then poured into 50 mL of 0.5 N HCl, and 50 mL of ethyl acetate were added. The aqueous layer was brought to pH 6 with 1N NaOH, 25 mL of sat. NaCl were added, and the layers were shaken and separated. The  
10 organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated to give an oily residue. This residue was subjected to flash column chromatography on silica gel with 20% ethyl acetate in hexanes to give 190 mg (54%) of the desired aldehyde as a yellow semisolid.

## Part 8. (2Z)-3-{[1-bis(t-butoxycarbonyl)amino]isoquinolin-7-yl}acrylic acid, 2-(trimethylsilyl)ethyl ester


A solution of 117 mg (0.29 mmol) of [bis(2,2,2-trifluoroethoxy)phosphinyl]acetic acid, 2-(trimethylsilyl)ethyl ester (*J. Org. Chem.*, 1991, 56, 4204) and 400 mg of 18-crown-6 in 25 mL of anhydrous THF was cooled with a dry ice-acetone bath under argon, and 0.75 mL of a 0.5 M solution of potassium bis(trimethylsilyl)amide in toluene were added dropwise  
20 over 2 min. The reaction was stirred in the cold for 15 min, then a solution of 100 mg (0.27 mmol) of 1-[bis(t-butoxycarbonyl)amino]isoquinoline-7-carboxaldehyde in 25 mL of anhydrous THF was added dropwise over 10 min. The reaction was then allowed to warm to room temperature overnight, then partitioned between 100 mL of CH<sub>2</sub>Cl<sub>2</sub> and 50 mL of H<sub>2</sub>O. The organics were washed with aqueous NaCl, and dried over MgSO<sub>4</sub>. Filtration  
25 and concentration gave an oily residue, which was subjected to flash column chromatography on silica gel with 25% ethyl acetate in hexanes to give 33 mg of the desired product as a clear, colorless oil.

Part 9. (2Z)-N-[5-(2{[(N-1,1-dimethylethyl)amino]sulfonyl}phenyl)-1-indoliny]-3-{[1-bis(t-butoxycarbonyl)amino]isoquinolin-7-yl}acrylamide


A solution of 498 mg (0.97 mmol) of (2Z)-3-{[1-bis(t-butoxycarbonyl)amino]isoquinolin-7-yl}acrylic acid, 2-(trimethylsilyl)ethyl ester in 5 mL of DMF was treated at room temperature with 1.2 mL (1.25 eq) of a 1.0 M tetrabutylammonium fluoride in THF, and the reaction was stirred for 5 hr. The reaction mixture was diluted with 100 mL of ether, and the solution was washed with 100 mL of water. The aqueous layer was again extracted with 100 mL of ether, and the combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration and concentration gave 350 mg of an off-white solid, which was used without further purification. A solution of 50 mg of this carboxylic acid and 44 mg of 2-indolin-5-ylbenzenesulfonamide from Example 27, Part 3 in 3 mL of DMF was treated with 100  $\mu$ L of N,N-diisopropylethylamine and 60 mg of HATU, and the reaction was stirred at room temperature overnight. The reaction mixture was diluted with 100 mL of ethyl acetate and washed with sat. NaHCO<sub>3</sub> (2 x 25 mL), and the organic layer was dried over MgSO<sub>4</sub>. Filtration and concentration gave an orange oil, which was subjected to flash column chromatography on silica gel, using 20% ethyl acetate in hexanes as eluent to give 62 mg of the desired product as a yellow oil.

Part 10. (2Z)-N-[4-(2{aminosulfonyl}phenyl)-1-indoliny]-3-{aminoisoquinolin-7-yl}acrylamide


A solution of the yellow oil from Part 9 in 2 mL of trifluoroacetic acid was stirred at room temperature overnight. The reaction mixture was concentrated with CH<sub>2</sub>Cl<sub>2</sub> to remove most of the TFA, then purified directly by prep HPLC to give 15 mg of the desired product was obtained as an off-white solid. LRMS (M + H)<sup>+</sup> = 470.

Example 41

This compound was prepared by a method similar to that used in Example 23, Part 6.

5 Example 42

This compound was prepared by a method similar to that used in Example 23, Part 6.

Example 43

10

This compound was prepared by catalytic hydrogenation of the compound from Example 42.

### BIOLOGICAL ACTIVITY EXAMPLES

Evaluation of the compounds of this invention is guided by in vitro protease activity assays (see below) and in vivo studies to evaluate antithrombotic efficacy, and effects on hemostasis and hematological parameters.

The compounds of the present invention are dissolved in buffer to give solutions containing concentrations such that assay concentrations range from 0 to 100  $\mu$ M. In the assays for thrombin, prothrombinase and factor Xa, a synthetic chromogenic substrate is added to a solution containing test compound and the enzyme of interest and the residual catalytic activity of that enzyme is determined spectrophotometrically. The  $IC_{50}$  of a compound is determined from the substrate turnover. The  $IC_{50}$  is the concentration of test compound giving 50% inhibition of the substrate turnover. The compounds of the present invention desirably have an  $IC_{50}$  of less than 500 nM in the factor Xa assay, preferably less than 200 nM, and more preferred compounds have an  $IC_{50}$  of about 100 nM or less in the factor Xa assay. The compounds of the present invention desirably have an  $IC_{50}$  of less than 4.0  $\mu$ M in the prothrombinase assay, preferably less than 200 nM, and more preferred compounds have an  $IC_{50}$  of about 10 nM or less in the prothrombinase assay. The compounds of the present invention desirably have an  $IC_{50}$  of greater than 1.0  $\mu$ M in the thrombin assay, preferably greater than 10.0  $\mu$ M, and more preferred compounds have an  $IC_{50}$  of greater than 100.0  $\mu$ M in the thrombin assay.

#### Amidolytic Assays for determining protease inhibition activity

The factor Xa and thrombin assays are performed at room temperature, in 0.02 M Tris·HCl buffer, pH 7.5, containing 0.15 M NaCl. The rates of hydrolysis of the para-nitroanilide substrate S-2765 (Chromogenix) for factor Xa, and the substrate Chromozym TH (Boehringer Mannheim) for thrombin following preincubation of the enzyme with inhibitor for 5 minutes at room temperature, and were determined using the Softmax 96-well plate reader (Molecular Devices), monitored at 405 nm to measure the time dependent appearance of p-nitroaniline.

The prothrombinase inhibition assay is performed in a plasma free system with modifications to the method described by Sinha, U. *et al.*, Thromb. Res., 75, 427-436 (1994). Specifically, the activity of the prothrombinase complex is determined by measuring the time course of thrombin generation using the p-nitroanilide substrate

5 Chromozym TH. The assay consists of preincubation ( 5 minutes) of selected compounds to be tested as inhibitors with the complex formed from factor Xa (0.5 nM), factor Va (2 nM), phosphatidyl serine:phosphatidyl choline (25:75, 20  $\mu$ M) in 20 mM Tris·HCl buffer, pH 7.5, containing 0.15 M NaCl, 5 mM CaCl<sub>2</sub> and 0.1% bovine serum albumin. Aliquots from the complex-inhibitor mixture are added to prothrombin (1 nM) and Chromozym TH  
10 (0.1 mM). The rate of substrate cleavage is monitored at 405 nm for two minutes. Eight different concentrations of inhibitor are assayed in duplicate. A standard curve of thrombin generation by an equivalent amount of untreated complex are used for determination of percent inhibition.

#### Antithrombotic Efficacy in a Rabbit Model of Venous Thrombosis

15 A rabbit deep vein thrombosis model as described by Hollenbach, S. et al., Thromb. Haemost. 71, 357-362 (1994), is used to determine the in-vivo antithrombotic activity of the test compounds. Rabbits are anesthetized with I.M. injections of Ketamine, Xylazine, and Acepromazine cocktail. A standardized protocol consists of insertion of a thrombogenic cotton thread and copper wire apparatus into the abdominal vena cava of the anesthetized rabbit. A  
20 non-occlusive thrombus is allowed to develop in the central venous circulation and inhibition of thrombus growth is used as a measure of the antithrombotic activity of the studied compounds. Test agents or control saline are administered through a marginal ear vein catheter. A femoral vein catheter is used for blood sampling prior to and during steady state infusion of test  
25 compound. Initiation of thrombus formation begins immediately after advancement of the cotton thread apparatus into the central venous circulation. Test compounds are administered from time = 30 min to time = 150 min at which the experiment is terminated. The rabbits are euthanized and the thrombus excised by surgical dissection and characterized by weight and

histology. Blood samples are analyzed for changes in hematological and coagulation parameters.

#### Effects of Compounds in Rabbit Venous Thrombosis model

Administration of compounds in the rabbit venous thrombosis model demonstrates  
5 antithrombotic efficacy at the higher doses evaluated. There are no significant effects of the compound on the aPTT and PT prolongation with the highest dose (100  $\mu\text{g/kg}$  + 2.57  $\mu\text{g/kg/min}$ ). Compounds have no significant effects on hematological parameters as compared to saline controls. All measurements are an average of all samples after steady state administration of vehicle or (D)-Arg-Gly-Arg-thiazole. Values are expressed as mean  $\pm$  SD.

10 Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods.

Unless defined otherwise, all technical and scientific terms used herein have the  
15 same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described.


20 It should be understood that the foregoing discussion and examples merely present a detailed description of certain preferred embodiments. It will be apparent to those of ordinary skill in the art that various modifications and equivalents can be made without departing from the spirit and scope of the invention. All the patents, journal articles and other documents discussed or cited above are herein incorporated by reference.

25



The claimed invention is:

1. A compound according to the formula:



wherein:

A is selected from:

- (a) C<sub>1</sub>-C<sub>6</sub>-alkyl;
- (b) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;
- (c) -N(-R<sup>2</sup>, -R<sup>3</sup>), R<sup>3</sup>-C(=N-R<sup>2</sup>)-, (-R<sup>2</sup>, -R<sup>3</sup>)N-C(=N-R<sup>2</sup>)-, (-R<sup>2</sup>, -R<sup>3</sup>)N-C(=N-R<sup>2</sup>)-N(-R-)-
- (d) phenyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;
- (e) naphthyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;  
and
- (f) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1</sup> substituents;

R<sup>1</sup> is selected from:

Halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, R<sup>2</sup>-C(=N-R<sup>3</sup>)-, (-R<sup>2</sup>, -R<sup>3</sup>)N-C(=N-R<sup>2</sup>)-, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -C(=O)-N(-R<sup>2</sup>, -R<sup>3</sup>), -SO<sub>2</sub>N(-R<sup>2</sup>, -R<sup>3</sup>), -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1</sub>-C<sub>4</sub>-alkyl, -CN C<sub>1</sub>.

<sub>4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

R<sup>2</sup> and R<sup>3</sup> are independently selected from the group consisting of:

H, -OR<sup>2</sup>, -N(-R<sup>2</sup>, -R<sup>3</sup>), -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylphenyl and -C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

or R<sup>2</sup> and R<sup>3</sup> taken together can form a 3-8 membered cycloalkyl or a heterocyclic ring system, wherein the heterocyclic ring system may have from 5 to 10 ring atoms, with 1 to 2 rings being in the ring system and contain from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the heterocyclic ring system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>-alkyl, -CN -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

m is an integer of 0-2;

Q is a member selected from the group consisting of:

a direct link, -CH<sub>2</sub>-, -C(=O)-, -N(R<sup>4</sup>)-, -N(R<sup>4</sup>)CH<sub>2</sub>-, -C=N(R<sup>4</sup>)-, -C(=O)-N(R<sup>4</sup>)-, -N(R<sup>4</sup>)-C(=O)-, -SO<sub>2</sub>-, -O-, -SO<sub>2</sub>-N(R<sup>4</sup>)- and -N(R<sup>4</sup>)-SO<sub>2</sub>-;

R<sup>4</sup> is selected from:

H, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylphenyl and -C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

D is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1a</sup> substituents;  
and



- (b) an aromatic six-membered heterocyclic ring having from 1-2 ring nitrogen atoms, and wherein the ring atoms may be substituted with 0-2 R<sup>1a</sup> substituents;

R<sup>1a</sup> is selected from:

Halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2a</sup>R<sup>3a</sup>, SO<sub>2</sub>NR<sup>2a</sup>R<sup>3a</sup>, SO<sub>2</sub>R<sup>2a</sup>, CF<sub>3</sub>, OR<sup>2a</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>2a</sup> and R<sup>3a</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

M is a member selected from the group consisting of:

-N(R<sup>16</sup>)-C(=O)-, -N(R<sup>16</sup>)-C(=S)-, -C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-, -C(-R<sup>19</sup>, -R<sup>19a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-, -C(-R<sup>17</sup>)-C(-R<sup>18</sup>)-C(=O)-, -C(-R<sup>17</sup>)-C(-R<sup>18</sup>)-C(=S)-, -C(-R<sup>17</sup>)-C(-R<sup>18</sup>)-, -O-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -O-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S(=O)<sub>2</sub>-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S(=O)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S(=O)<sub>2</sub>-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S(=O)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(=O)-C(=O)-, -N(R<sup>16</sup>)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -N(R<sup>16</sup>)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(=S)-C(=S)-, -C(=S)-C(=O)-, -C(=O)-C(=S)-, -N=C(-R<sup>17</sup>)-C(=O)-, -N=C(-R<sup>17</sup>)-C(=S)-, -C(-R<sup>17</sup>)=N-, -N(-R<sup>16</sup>)-C(=O)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -O-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -N(-R<sup>16</sup>)-C(=S)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -S(=O)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -S(=O)<sub>2</sub>-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -C(=C(R<sup>17b</sup>, -R<sup>17c</sup>))-C(=O)-, -C(=C(R<sup>17b</sup>, -R<sup>17c</sup>))-C(=S)-, -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(=S)-, -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-N(-H, -R<sup>18b</sup>), -R<sup>17a</sup>)-C(=O)-; -N=C(-R<sup>17</sup>)- and

$-N(-R^{16})-C(-R^{18}, -R^{18a})-C(-(N(-H, -R^{18b})), -R^{17a})-C(=S)-$ ; wherein the first named atom of the chain is directly attached to D, and wherein D, M and the N atom attached to the last chain atom of M collectively form a bicyclic ring structure;

$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$ , and  $R^{19a}$  are each independently selected from the group consisting of:

hydrogen, halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$ ,  $-NO_2$ ,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$ ,  $SO_2R^2$ ,  $CF_3$ ,  $OR^2$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_1-C_4$ -alkyl,  $-CN$ ,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl and  $-NO_2$ ;

$R^{17b}$  and  $R^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy,  $-C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $-C_{2-6}$ alkynyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl,  $-CN$ ,  $-NO_2$ ,  $-(CH_2)_mNR^2R^3$ ,  $-SO_2NR^2R^3$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the cycloalkyl, the phenyl ring, or the aromatic heterocyclic ring may be independently replaced with a member selected from the group consisting of halo,  $C_1-C_4$ -alkyl,  $-CN$ ,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl and  $-NO_2$ ;

E is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $C_{0-4}$ alkylheteroaryl,  $C_{1-4}$ alkylCOOH and  $C_{1-4}$ alkylCOOC $C_{1-4}$ alkyl, wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,

C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -OH, -O-C<sub>1-4</sub>alkyl, -SH, -S-C<sub>1-4</sub>alkyl, -CN and -NO<sub>2</sub>;

G is selected from:

a direct link, -C(R<sup>7</sup>,R<sup>8</sup>)-, -C(R<sup>7a</sup>,R<sup>8a</sup>)-C(R<sup>7b</sup>,R<sup>8b</sup>)- and -C(R<sup>7c</sup>)=C(R<sup>8c</sup>)-;

wherein R<sup>7</sup>, R<sup>8</sup>, R<sup>7a</sup>, R<sup>8a</sup>, R<sup>7b</sup>, R<sup>8b</sup>, R<sup>7c</sup> and R<sup>8c</sup> are independently a member selected from the group consisting of:

hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)OR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, a naturally occurring or synthetic amino acid side chain, and C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the C<sub>0-4</sub>alkylheterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>9</sup> and R<sup>10</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>3-8</sub>cycloalkyl, and C<sub>1-4</sub>alkyl-O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl-COOH wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>, and wherein R<sup>9</sup> and R<sup>10</sup> taken together can form a 5-8 membered heterocyclic ring;

J is a member selected from the group consisting of:

a direct link, -O-, -O-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -S-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)<sub>2</sub>-(-R<sup>11</sup>, -R<sup>11a</sup>)-, -C(=O)-, -C(=O)-N(R<sup>11b</sup>)-,

$-N(R^{11b})-C(=O)-$ ,  $-N(R^{11b})-$ ,  $-N(R^{11b})-C(-R^{11}, -R^{11a})-$  and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{11}$  substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$ , and  $R^{11}$  are a member independently selected from the group consisting of:

hydrogen, halo,  $-CF_3$ ,  $-CN$ ,  $-NR^9R^{10}$ ,  $-SO_2Me$ ,  $-NO_2$ ,  $-OH$ ,  $-O-C_{1-4}alkyl$ ,  $-O-C_{2-6}alkenyl$ ,  $-O-C_{2-6}alkynyl$ ,  $-O-C_{3-8}cycloalkyl$ ,  $-O-C_{1-4}alkyl-O-C_{1-4}alkyl$ ,  $-O-C_{1-4}alkyl-COOH$ ,  $-O-C_{1-4}alkyl-phenyl$ ,  $-COOH$ ,  $-C(=O)-O-C_{1-4}alkyl$ ,  $-C(=O)-O-C_{2-6}alkenyl$ ,  $-C(=O)-O-C_{2-6}alkynyl$ ,  $-C(=O)-O-C_{3-8}cycloalkyl$ ,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $C_{0-4}alkylC(=O)NR^9R^{10}$ ,  $C_{0-4}alkylC(=O)OR^9$ ,  $C_{0-4}alkylheterocyclic$  ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S,  $CH_2COOC_{1-4}alkyl$ ,  $CH_2COOC_{1-4}alkylphenyl$  and  $CH_2COOC_{1-4}alkylnaphthyl$ ; wherein from 1-4 hydrogen atoms on the  $C_{0-4}alkylheterocyclic$  ring may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkyl-phenyl$ ,  $C_{0-4}alkylC_{3-8}cycloalkyl$ ,  $-CN$  and  $-NO_2$ ;

wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $-OR^9$ ,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ ,  $-CN$  and  $-NO_2$ ;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1b}$  substituents;
- (b) naphthyl, which is independently substituted with 0-2  $R^{1b}$  substituents;  
and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{1b}$  substituents;

$R^{1b}$  is a member selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN, -NO<sub>2</sub>,  $NR^{2b}R^{3b}$ ,  $SO_2NR^{2b}R^{3b}$ ,  $SO_2R^{2b}$ ,  $CF_3$ ,  $OR^{2b}$ ,  $O-CH_2-CH_2-OR^{2b}$ ,  $O-CH_2-COOR^{2b}$ ,  $N(R^{2b})-CH_2-CH_2-OR^{2b}$ ,  $N(-CH_2-CH_2-OR^{2b})_2$ ,  $N(R^{2b})-C(=O)R^{3b}$ ,  $N(R^{2b})-SO_2-R^{3b}$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl-phenyl, -CN and -NO<sub>2</sub>;

$R^{2b}$  and  $R^{3b}$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN and -NO<sub>2</sub>;

L is selected from:

H, -CN,  $C(=O)NR^{12}R^{13}$ ,  $-(CH_2)_nNR^{12}R^{13}$ ,  $C(=NR^{12})NR^{12}R^{13}$ ,  $OR^{12}$ ,  $-NR^{12}C(=NR^{12})NR^{12}R^{13}$ , and  $NR^{12}C(=NR^{12})-R^{13}$ ;

n is an integer from 0 to 8;

$R^{12}$  and  $R^{13}$  are independently selected from:

hydrogen, -OR<sup>14</sup>, -NR<sup>14</sup>R<sup>15</sup>,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $COOC_{1-4}$ alkyl,  $COO-C_{0-4}$ alkylphenyl and  $COO-C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH, -O- $C_{1-4}$ alkyl,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN, and -NO<sub>2</sub>;

$R^{14}$  and  $R^{15}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on



the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

2. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 1.
3. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 1.
4. The method of claim 3, wherein the condition is selected from the group consisting of:  
acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.
5. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 1.

6. A compound of claim 1,

wherein:

A is selected from:

- (a) C<sub>1</sub>-C<sub>6</sub>-alkyl;
- (b) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;
- (c) phenyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;
- (d) naphthyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;  
and
- (e) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from the group consisting of N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1</sup> substituents;

R<sup>1</sup> is selected from:

halo, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, -C<sub>1-4</sub>alkyl, -CN, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

R<sup>2</sup> and R<sup>3</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

m is an integer of 0-2;

Q is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-N(R^4)-$ ,  $-C(=O)-N(R^4)-$ ,  $-N(R^4)-C(=O)-$ ,  $-SO_2-$ ,  $-O-$ ,  $-SO_2-N(R^4)-$  and  $-N(R^4)-SO_2-$ ;

$R^4$  is selected from:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$ , and  $-NO_2$ ;

D is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1a}$  substituents; and
- (b) an aromatic six-membered heterocyclic ring having from 1-2 ring nitrogen atoms, and wherein the ring atoms may be substituted with 0-2  $R^{1a}$  substituents;

$R^{1a}$  is selected from:


halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$ ,  $-NO_2$ ,  $(CH_2)_mNR^{2a}R^{3a}$ ,  $SO_2NR^{2a}R^{3a}$ ,  $SO_2R^{2a}$ ,  $CF_3$ ,  $OR^{2a}$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$  and  $-NO_2$ ;

$R^{2a}$  and  $R^{3a}$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$  and  $-NO_2$ ;



M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



and the like, wherein the aromatic carbocyclic ring corresponding to the D portion for each of the bicyclic rings may be replaced with an aromatic heterocyclic ring as defined above for D, and wherein 0 to 2 of the hydrogen atoms on the D portion of the bicyclic ring may be replaced by  $R^{1a}$  substituents as defined above;

$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN, -NO<sub>2</sub>,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$ ,  $SO_2R^2$ , CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl, -CN,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl and -NO<sub>2</sub>;

$R^{17b}$  and  $R^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy,  $-C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $-C_{2-6}alkynyl$ ,  $-C_{3-8}cycloalkyl$ ,  $-C_{0-4}alkyl-C_{3-8}cycloalkyl$ , -CN,  $-NO_2$ ,  $-(CH_2)_mNR^2R^3$ ,  $-SO_2NR^2R^3$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the cycloalkyl, the phenyl ring, or the aromatic heterocyclic ring may be independently replaced with a member selected from the group consisting of halo,  $C_1-C_4-alkyl$ , -CN,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylC_{3-8}cycloalkyl$  and  $-NO_2$ ;

E is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-(-C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-)$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylC_{3-8}cycloalkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $C_{0-4}alkylheteroaryl$ ,  $C_{1-4}alkylCOOH$  and  $C_{1-4}alkylCOOC_{1-4}alkyl$ , wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylC_{3-8}cycloalkyl$ , -OH,  $-O-C_{1-4}alkyl$ , -SH,  $-S-C_{1-4}alkyl$ , -CN and  $-NO_2$ ;

G is selected from:

a direct link,  $-C(R^7, R^8)-$ ,  $-C(R^{7a}, R^{8a})-C(R^{7b}, R^{8b})-$  and  $-C(R^{7c})=C(R^{8c})-$ ;

wherein  $R^7$ ,  $R^8$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{7b}$ ,  $R^{8b}$ ,  $R^{7c}$  and  $R^{8c}$  are independently a member selected from the group consisting of:

hydrogen, halogen,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $-OR^9$ ,  $-N(R^9R^{10})$ ,  $-C_{0-4}alkylCOOR^9$ ,  $-C_{0-4}alkylC(=O)NR^9R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9-CH_2-CH_2-O-R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9(-CH_2-CH_2-O-R^{10})_2$ ,  $-N(R^9)COR^{10}$ ,  $-N(R^9)C(=O)R^{10}$ ,  $-N(R^9)SO_2R^{10}$ , and a naturally occurring or synthetic amino acid side chain, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $-OR^9$ ,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ , -CN and  $-NO_2$ ;

$R^9$  and  $R^{10}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl, -CN and -NO<sub>2</sub>, and wherein  $R^9$  and  $R^{10}$  taken together can form a 5-8 membered heterocyclic ring;

J is a member selected from the group consisting of:

a direct link, -O-, -O-C( $R^{11}$ ,  $R^{11a}$ )-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -S-C( $R^{11}$ ,  $R^{11a}$ )-, -S(=O)-C( $R^{11}$ ,  $R^{11a}$ )-, -S(=O)<sub>2</sub>-( $R^{11}$ ,  $R^{11a}$ )-, -C(=O)-, -C(=O)-N( $R^{11b}$ )-, -N( $R^{11b}$ )-C(=O)-, -N( $R^{11b}$ )-, -N( $R^{11b}$ )-C( $R^{11}$ ,  $R^{11a}$ )- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{11c}$  substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$  and  $R^{11c}$  are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O- $C_{1-4}$ alkyl, -O- $C_{2-6}$ alkenyl, -O- $C_{2-6}$ alkynyl, -O- $C_{3-8}$ cycloalkyl, -COOH, -C(=O)-O- $C_{1-4}$ alkyl, -C(=O)-O- $C_{2-6}$ alkenyl, -C(=O)-O- $C_{2-6}$ alkynyl, -C(=O)-O- $C_{3-8}$ cycloalkyl,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $C_{0-4}$ alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC $C_{1-4}$ alkyl, CH<sub>2</sub>COOC $C_{1-4}$ alkylphenyl and CH<sub>2</sub>COOC $C_{1-4}$ alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1b}$  substituents;
- (b) naphthyl, which is independently substituted with 0-2  $R^{1b}$  substituents;  
and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{1b}$  substituents;

$R^{1b}$  is a member selected from the group consisting of:

halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>R<sup>2b</sup>, CF<sub>3</sub>, OR<sup>2b</sup>, O-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, O-CH<sub>2</sub>-COOR<sup>2b</sup>, N(R<sup>2b</sup>)-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, N(-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>)<sub>2</sub>, N(R<sup>2b</sup>)-C(=O)R<sup>3b</sup>, N(R<sup>2b</sup>)-SO<sub>2</sub>-R<sup>3b</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>2b</sup> and R<sup>3b</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

L is selected from:

H, -CN, C(=O)NR<sup>12</sup>R<sup>13</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup>, C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, OR<sup>12</sup>, -NR<sup>12</sup>C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, and NR<sup>12</sup>C(=NR<sup>12</sup>)-R<sup>13</sup>;

n is an integer from 0 to 8;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

hydrogen, -OR<sup>14</sup>, -NR<sup>14</sup>R<sup>15</sup>, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, COOC<sub>1-4</sub>alkyl, COO-C<sub>0-4</sub>alkylphenyl and COO-C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH, -O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

R<sup>14</sup> and R<sup>15</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl,

C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

7. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 6.
8. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 6.
9. The method of claim 8, wherein the condition is selected from the group consisting of:  
acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.
10. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 6.

11. A compound of claim 1,

wherein:

A is selected from:

- (a) C<sub>1</sub>-C<sub>6</sub>-alkyl;
- (b) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;
- (c) phenyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;
- (d) naphthyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;  
and
- (e) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1</sup> substituents;

R<sup>1</sup> is selected from:

halo, C<sub>1-4</sub>alkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>R<sup>2</sup>, CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

$R^2$  and  $R^3$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;

m is an integer of 0-2;

B is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-N(R^4)-$ ,  $-C(=O)-N(R^4)-$ ,  $-N(R^4)-C(=O)-$ ,  $-SO_2-$ ,  $-O-$ ,  $-SO_2-N(R^4)-$  and  $-N(R^4)-SO_2-$ ;

$R^4$  is selected from:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;

D is phenyl, which is independently substituted with 0-2  $R^{1a}$  substituents;

$R^{1a}$  is selected from:


halo,  $C_{1-4}$ alkyl,  $-CN$ ,  $-NO_2$ ,  $-(CH_2)_mNR^{2a}R^{3a}$ ,  $-SO_2NR^{2a}R^{3a}$ ,  $-SO_2R^{2a}$ ,  $CF_3$ ,  $-OR^{2a}$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

$R^{2a}$  and  $R^{3a}$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;



M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



wherein 0 to 2 of the hydrogen atoms on the D portion of the bicyclic ring may be replaced by  $R^{1a}$  substituents as defined above;

$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $-CN$ ,  $-NO_2$ ,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$ ,  $SO_2R^2$ ,  $CF_3$ ,  $OR^2$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;



$R^{17b}$  and  $R^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy,  $-C_{1-4}alkyl$ ,  $-CN$ ,  $-NO_2$ ,  $-(CH_2)_mNR^2R^3$ ,  $-SO_2NR^2R^3$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S;

$m$  is an integer from 0-6;

$E$  is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-(-C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-);$

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H,  $C_{1-4}alkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $C_{0-4}alkylheteroaryl$ ,  $C_{1-4}alkylCOOH$  and  $C_{1-4}alkylCOOC_{1-4}alkyl$ , wherein from 0-2 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}alkyl$ ,  $-OH$ ,  $-O-C_{1-4}alkyl$ ,  $-SH$ ,  $-S-C_{1-4}alkyl$ ,  $-CN$  and  $-NO_2$ ;

$G$  is selected from:

a direct link,  $-C(R^7, R^8)-$ ,  $-C(R^{7a}, R^{8a})-C(R^{7b}, R^{8b})-$  and  $-C(R^{7c})=C(R^{8c})-$ ;

wherein  $R^7$ ,  $R^8$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{7b}$ ,  $R^{8b}$ ,  $R^{7c}$  and  $R^{8c}$  are independently a member selected from from the group consisting of:

hydrogen, halogen,  $C_{1-4}alkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $-OR^9$ ,  $-N(R^9R^{10})$ ,  $-C_{0-4}alkylCOOR^9$ ,  $-C_{0-4}alkylC(=O)NR^9R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9-CH_2-CH_2-O-R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9(-CH_2-CH_2-O-R^{10})_2$ ,  $-N(R^9)COR^{10}$ ,  $-N(R^9)C(=O)R^{10}$ ,  $-N(R^9)SO_2R^{10}$ , and a naturally occurring or synthetic amino acid side chain;

$R^9$  and  $R^{10}$  are independently selected from:

H,  $C_{1-4}alkyl$ ,  $C_{0-4}alkylphenyl$  and  $C_{0-4}alkylnaphthyl$ ;

$J$  is a member selected from the group consisting of:

a direct link,  $-O-$ ,  $-O-C(-R^{11}, -R^{11a})-$ ,  $-S-$ ,  $-S(=O)_2-$ ,  $-S-C(-R^{11}, -R^{11a})-$ ,  $-S(=O)_2-(-R^{11}, -R^{11a})-$ ,  $-C(=O)-N(R^{11b})-$ ,  $-N(R^{11b})-$ ,  $-N(R^{11b})-C(-R^{11}, -R^{11a})-$  and a

monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{11c}$  substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$  and  $R^{11c}$  are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>3-8</sub>cycloalkyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, and a C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1b}$  substituents;
- (b) naphthyl, which is independently substituted with 0-2  $R^{1b}$  substituents; and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{1b}$  substituents;

$R^{1b}$  is a member selected from the group consisting of:

halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>R<sup>2b</sup>, CF<sub>3</sub>, OR<sup>2b</sup>, O-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, O-CH<sub>2</sub>-COOR<sup>2b</sup>, N(R<sup>2b</sup>)-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, N(-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>)<sub>2</sub>, N(R<sup>2b</sup>)-C(=O)R<sup>3b</sup>, N(R<sup>2b</sup>)-SO<sub>2</sub>-R<sup>3b</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

$R^{2b}$  and  $R^{3b}$  are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

L is selected from:

H, -CN, C(=O)NR<sup>12</sup>R<sup>13</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup>, C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, OR<sup>12</sup>, -NR<sup>12</sup>C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, and NR<sup>12</sup>C(=NR<sup>12</sup>)-R<sup>13</sup>;

n is an integer from 0 to 6;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

hydrogen, -OR<sup>14</sup>, -NR<sup>14</sup>R<sup>15</sup>, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, COOC<sub>1-4</sub>alkyl, COO-C<sub>0-4</sub>alkylphenyl and COO-C<sub>0-4</sub>alkylnaphthyl, wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH, -O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

R<sup>14</sup> and R<sup>15</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

12. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 11.
13. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 11.
14. The method of claim 13, wherein the condition is selected from the group consisting of:

acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.

15. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 11.

16. A compound of claim 1,

wherein:

A is selected from:

- (a) phenyl, which is independently substituted with 0-2  $R^1$  substituents; and
- (b) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^1$  substituents;

$R^1$  is selected from:

halo,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$  and  $SO_2R^2$ ;

$R^2$  and  $R^3$  are independently selected from the group consisting of:

H and  $C_{1-4}$ alkyl;

m is an integer of 0-2;

Q is a member selected from the group consisting of:


a direct link,  $-C(=O)-$ ,  $-SO_2-$ , and  $-O-$ ;

D is phenyl, which is independently substituted with 0-2  $R^{1a}$  substituents;

$R^{1a}$  is selected from:

halo and  $C_{1-4}$ alkyl;

M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:

halo, C<sub>1-4</sub>alkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>R<sup>2</sup>, CF<sub>3</sub> and OR<sup>2</sup>;

R<sup>17b</sup> and R<sup>17c</sup> are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy, -C<sub>1-4</sub>alkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-3 N atoms;

E is a member selected from the group consisting of:

a direct link, -C(=O)-, -C(=O)-N(R<sup>5</sup>)-, -C(-R<sup>5a</sup>, -R<sup>6a</sup>)- and -(-C(-R<sup>5b</sup>, -R<sup>6b</sup>)-C(-R<sup>5c</sup>, -R<sup>6c</sup>)-;

wherein R<sup>5</sup>, R<sup>5a</sup>, R<sup>6a</sup>, R<sup>5b</sup>, R<sup>6b</sup>, R<sup>5c</sup> and R<sup>6c</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylheteroaryl, C<sub>1-4</sub>alkylCOOH and C<sub>1-4</sub>alkylCOOC<sub>1-4</sub>alkyl;

G is selected from:

a direct link, -C(R<sup>7</sup>, R<sup>8</sup>)-, -C(R<sup>7a</sup>, R<sup>8a</sup>)-C(R<sup>7b</sup>, R<sup>8b</sup>)- and -C(R<sup>7c</sup>)=C(R<sup>8c</sup>)-;

wherein R<sup>7</sup>, R<sup>8</sup>, R<sup>7a</sup>, R<sup>8a</sup>, R<sup>7b</sup>, R<sup>8b</sup>, R<sup>7c</sup> and R<sup>8c</sup> are independently a member selected from the group consisting of:

hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)-<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, and a naturally occurring or synthetic amino acid side chain;

R<sup>9</sup> and R<sup>10</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

J is a member selected from the group consisting of:

a direct link, -O-, -S-, -C(=O)-N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(-R<sup>11</sup>, -R<sup>11a</sup>)- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>11c</sup> substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$  and  $R^{11c}$  are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>3-8</sub>cycloalkyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, and a C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1b}$  substituents;
- (b) an aromatic heterocyclic ring having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring may be substituted with 0-2  $R^{1b}$  substituents;
- (c) a fused aromatic bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the bicyclic ring system may be substituted with 0-2  $R^{1b}$  substituents;

$R^{1b}$  is a member selected from the group consisting of:

halo, -C<sub>1-4</sub>alkyl, -OH, -OBn, -O-CH<sub>2</sub>-CH<sub>2</sub>-OH, -O-CH<sub>2</sub>-CH<sub>2</sub>-OCH<sub>3</sub>, -O-CH<sub>2</sub>-COOH, -O-CH<sub>2</sub>-C(=O)-O-CH<sub>3</sub>, -NH<sub>2</sub>, -NH-CH<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>3</sub>, -NH-C(=O)-O-CH<sub>3</sub> and -NH-SO<sub>2</sub>-CH<sub>3</sub>;

L is selected from:

H, -C(=O)NR<sup>12</sup>R<sup>13</sup>, -(CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup> and -C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>;

n is an integer from 0 to 6;

$R^{12}$  and  $R^{13}$  are independently selected from:

hydrogen and C<sub>1-4</sub>alkyl;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug




derivatives thereof.


17. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 16.
18. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 16.
19. The method of claim 18, wherein the condition is selected from the group consisting of:  
acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.
20. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 16.



21. A compound of claim 1, wherein A is a member selected from the group consisting of:



Q is a member selected from the group consisting of:



E is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-(-C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-)$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $C_{0-4}$ alkylheteroaryl,  $C_{1-4}$ alkylCOOH and  $C_{1-4}$ alkylCOOC $C_{1-4}$ alkyl;

G is selected from:

a direct link,  $-C(R^7, R^8)-$ ,  $-C(R^{7a}, R^{8a})-C(R^{7b}, R^{8b})-$  and  $-C(R^{7c})=C(R^{8c})-$ ;

wherein  $R^7$ ,  $R^8$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{7b}$ ,  $R^{8b}$ ,  $R^{7c}$  and  $R^{8c}$  are independently a member selected from from the group consisting of:

hydrogen, halogen,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $-OR^9$ ,  $-N(R^9R^{10})$ ,  $-C_{0-4}$ alkylCOOR<sup>9</sup>,  $-C_{0-4}$ alkylC(=O)NR<sup>9</sup>R<sup>10</sup>,  $-C_{0-4}$ alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>,  $-C_{0-4}$ alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)-<sub>2</sub>,  $-N(R^9)COR^{10}$ ,  $-N(R^9)C(=O)R^{10}$ ,  $-N(R^9)SO_2R^{10}$ , and a naturally occurring or synthetic amino acid side chain;

$R^9$  and  $R^{10}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;


J is a member selected from the group consisting of:

a direct link,  $-O-$ ,  $-S-$ ,  $-C(=O)-N(R^{11b})-$ ,  $-N(R^{11b})-$ ,  $-N(R^{11b})-C(-R^{11}, -R^{11a})-$  and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{11c}$  substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$  and  $R^{11c}$  are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -C<sub>1-4</sub>alkyl, -COOH, phenyl, and benzyl wherein the aromatic ring of the phenyl or benzyl is substituted with 0-2 members independently selected from the group consisting of halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -C<sub>1-4</sub>alkyl, -COOH and -C(=O)-O-C<sub>1-4</sub>alkyl;


Y and L taken together are a member selected from the group consisting of:




and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

22. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 16.
23. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 16.
24. The method of claim 18, wherein the condition is selected from the group consisting of:  
acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.
25. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 16.


26. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{11b}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^7$  is selected from the group consisting of:


H; Me; Et; Ph;




27. A compound of claim 1, having the formula:



wherein the following portion of said formula




is selected from the group consisting of:






wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{11c1}$  is selected from the group consisting of:

H; Me; Et; Ph; OMe; CF<sub>3</sub>; F; OH; Br; NH<sub>2</sub>; SO<sub>2</sub>Me; OCH<sub>2</sub>Ph; OCH<sub>2</sub>CH<sub>2</sub>OMe;




and wherein  $R^{11c2}$  is selected from the group consisting of:


H; Me; Et; Cl; Br; F; Ph;




28. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{1c1}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^{1c2}$  is selected from the group consisting of:


H; Me; Et; Ph;




29. A compound of claim 1, having the formula:




wherein the following portion of said formula:



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{1c1}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^{1c2}$  is selected from the group consisting of:


H; Me; Et; Ph;



30. A compound of claim 1, having the formula:




wherein the following portion of said formula






is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{7c}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^{8c}$  is selected from the group consisting of:


H; Me; Et; Ph;




31. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{7a}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^{7b}$  is selected from the group consisting of:


H; Me; Et; Ph;




32. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:






wherein  $R^{7a}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^{7b}$  is selected from the group consisting of:


H; Me; Et; Ph;




33. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{6c}$  is selected from the group consisting of:

H; Me; Et; Ph; OMe; F; OH; Br;  $NH_2$ ;  $SO_2Me$ ;  $OCH_2Ph$ ;  $OCH_2CH_2OMe$ ;




and wherein  $R^{11b}$  is selected from the group consisting of:


H; Me; Et; Cl; Br; F; Ph;




34. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{11c1}$  is selected from the group consisting of:

H; Me; Et; Ph; OMe; F; OH; Br; NH<sub>2</sub>; SO<sub>2</sub>Me; OCH<sub>2</sub>Ph; OCH<sub>2</sub>CH<sub>2</sub>OMe;




and wherein  $R^{11c2}$  is selected from the group consisting of:


H; Me; Et; Cl; Br; F; Ph;




35. A compound of claim 1, having the formula:



wherein the following portion of said formula




is selected from the group consisting of:






wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{11c1}$  is selected from the group consisting of:

H; Me; Et; Ph; OMe; F; OH; Br; NH<sub>2</sub>; SO<sub>2</sub>Me; OCH<sub>2</sub>Ph; OCH<sub>2</sub>CH<sub>2</sub>OMe;




and wherein  $R^{11c2}$  is selected from the group consisting of:


H; Me; Et; Cl; Br; F; Ph;




36. A compound of claim 1, having the formula:



wherein the following portion of said formula




is selected from the group consisting of:






wherein  $R^{1c1}$  is selected from the group consisting of:

H; Me; Et; Ph; OMe; F; OH; Br; NH<sub>2</sub>; SO<sub>2</sub>Me; OCH<sub>2</sub>Ph; OCH<sub>2</sub>CH<sub>2</sub>OMe;




and wherein  $R^{1c2}$  is selected from the group consisting of:


H; Me; Et; Cl; Br; F; Ph;




37. A compound of claim 1, selected from the group consisting of:




38. A compound of claim 1, selected from the group consisting of:




39. A compound of claim 1, selected from the group consisting of:




40. A compound of claim 1, selected from the group consisting of:



41. A compound of claim 1, selected from the group consisting of:






42. A compound according to the formula:




wherein A—Q— is selected from the group consisting of:

t-Bu ; O-t-Bu ;  $-(CH_2)_{0-5}$ -amino ; OH ; carboxylic acid ester; carboxamide ;







43. A compound of claim 1, selected from the group consisting of:




44. A compound of claim 1, selected from the group consisting of:




45. A compound of claim 1, selected from the group consisting of:




46. A compound of claim 1, selected from the group consisting of:




47. A compound of claim 1, selected from the group consisting of:




48. A compound of claim 1, selected from the group consisting of:




49. A compound of claim 1, selected from the group consisting of:



50. A compound of claim 1, selected from the group consisting of:




51. A compound of claim 1, selected from the group consisting of:







52. A compound, selected from the group consisting of:



wherein Q is a direct link, and A is a member selected from the group:



or Q is a  $-C(=NH)-$  group, and A is a member selected from the group:




G is a direct link;

J is a member selected from the group:



Y-L is a member selected from the group:



# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/21742

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07D 209/04, 215/00, 235/04, 241/36, 265/36, 403/02, 498/02

US CL : 544/105, 354; 546/165; 548/305.4, 364.7, 490

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 544/105, 354; 546/165; 548/305.4, 364.7, 490

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
EAST, Registry Online, CA Online

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 129:330730,	1-36, 52
---	RIES, U. 'Preparation of disubstituted bicyclic heterocyclic antithrombotics and	-----
Y	anticoagulants,' abstract, DE 19718181, 05 November 1998.	37-51
X	Database CA on STN, Chemical Abstracts (Columbus, OH, USA), No. 128:16433,	1-36, 52
---	SUZUKI, N. 'Preparation of thienopyridininones as GNRH agonists and antagonists,'	-----
Y	abstract, WO 97/40846, November 1997.	37-51
X	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 127:135807,	1-36, 52
---	SUZUKI, N. 'Preparation of condensed bicyclic compounds as prolactin production	-----
Y	inhibitors,' abstract, EP 781774, July 1997.	37-51
X	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 127:50652,	1-36, 52
---	FRECHETTE, R. 'Preparation of benzoxazine antimicrobial agents,' abstract, WO	-----
Y	97/17333, May 1997.	37-51
X	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 126:343504,	1-36, 52
---	FURUYA, S. 'Preparation of quinolin-4-ones as GnRH antagonists,' abstract WO	-----
Y	97/14682, April 1997.	37-51



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

06 November 2000 (06.11.2000)

Date of mailing of the international search report

25 JAN 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

*Dorothea Lawrence*  
D. Margaret Seaman

Telephone No. 703-308-1235

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/21742

## C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No 125:167737, KATRITZKY, A.R. 'A new indole synthesis via [3+3] annulation of 2-(benzotriazol-1-ylmethyl)pyrroles with .alpha...beta.-unsaturated aldehydes and ketones,' abstract Tetrahedron Lett, Vol.37, No. 32, pages 5641-5644, 1996.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 124:317157, SUMMERS, J.B. 'Platelet activating factor antagonists: imidazopyridine indoles,' abstract, US patent #5,486,525, January 1996.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 124:175907, SHEPPARD, G.S. 'Synthesis and evaluation of water soluble indole pyrrolothiazole PAF antagonists,' abstract, Bioorg. Med. Chem. Lett. Vol. 5, No. 23, pages 2913-1918, 1995.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 123:340121, SUMMERS, J.B. 'Preparation of 3-[(imidazopyridylalkyl)benzoyl]indoles and analogs as platelet activating factor antagonists,' abstract, WO 95/16687, June 1995.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 123:32955, BACH, N.J. 'Preparation of 1H-Indole-3-Acetic acid hydrazides as sPLA2 inhibitors,' abstract, EP 620214, October 1994.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA) No. 122:290852, BOYD, D. B. 'Preparation of arylindoles, benzimidazoles, and -indazoles as angiotensin II angatonists,' abstract, EP 574174, December 1993.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 119:49384, SUMMERS, J.B. 'Preparation of 7-indol-3-yl carbonyl pyrrolo 1,2-c thiazoles and related compounds as platelet activating factor antagonists,' abstract, WO 93/01813, February 1992.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA) No. 99:157, KOLASA, K. 'Preliminary pharmacological studies of the central action of phenyl and piperidinomethyl derivatives of 2-benzoxazolone,' abstract, Ann. Univ. mariae Curie-Sklodowska, Sect. D, Vol. 36, pages 73-81, 1981.	1-36, 52 ----- 37-51

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number  
**WO 01/012600 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 209/04**,  
215/00, 235/04, 241/36, 265/36, 403/02, 498/02

CA 94115 (US). **ZHANG, Penglie** [CN/US]; 224 Serrano  
Drive, San Francisco, CA 94132 (US).

(21) International Application Number: PCT/US00/21742

(74) Agent: **WEIMAR, Elizabeth, C.**; Morgan, Lewis &  
Bockius LLP, 1800 M Street NW, Washington, DC 20036  
(US).

(22) International Filing Date: 10 August 2000 (10.08.2000)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AI, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:  
60/148,627 12 August 1999 (12.08.1999) US  
60/202,202 5 May 2000 (05.05.2000) US

(71) Applicant (*for all designated States except US*): **COR  
THERAPEUTICS, INC.** [US/US]; 256 East Grand  
Avenue, South San Francisco, CA 94080 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ZHU, Bing-Yan**  
[CA/US]; 3325 Adelaide Way, Belmont, CA 94002  
(US). **SCARBOROUGH, Robert, M.** [US/US]; 22  
Greenbrier Court, Half Moon Bay, CA 94019 (US).  
**CLIZBE, Lane** [US/US]; 392 Newcastle Drive, Red-  
wood City, CA 94061 (US). **DOUGHAN, Brandon**  
[US/US]; 650 Guerrero Street, San Francisco, CA 94110  
(US). **JIA, Zhaozhong-Jon** [CN/US]; 849 West Or-  
ange Avenue #2007, San Francisco, CA 94110 (US).  
**KANE-MAGUIRE, Kim** [US/US]; 3425 Lodge Drive,  
Belmont, CA 94002 (US). **MARLOWE, Charles**  
[US/US]; 636 California Way, Redwood City, CA 94080  
(US). **SONG, Yonghong** [CA/US]; 1144 Nimitz Lane,  
Foster City, CA 94404 (US). **SU, Ting** [US/US]; 3222  
Upper Lock Avenue, Belmont, CA 94002 (US). **TENG,  
Willy** [US/US]; 2355 Scott Street #103, San Francisco,

**Published:**

— *with international search report*

(48) Date of publication of this corrected version:


12 September 2002

(15) Information about Correction:

see PCT Gazette No. 37/2002 of 12 September 2002, Sec-  
tion II

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: INHIBITORS OF FACTOR Xa



(57) Abstract: Novel compounds, their salts and compositions related thereto having activity against mammalian factor Xa are disclosed. The compounds of formula (I) are useful *in vitro* or *in vivo* for preventing or treating coagulation disorders.

WO 01/012600 A1

## INHIBITORS OF FACTOR Xa

Related Applications

This application claims benefit of priority under 35 USC § 119(e) to U.S. Provisional Application No. 60/202,202 filed on May 5, 2000 and U.S. Provisional Application No. 60/148,627 filed on August 12, 1999, which are both herein incorporated in their entirety by reference.

Field of the Invention

This invention relates to novel compounds which are potent and highly selective inhibitors of isolated factor Xa or factor Xa when assembled in the prothrombinase complex. These compounds show selectivity for factor Xa versus other proteases of the coagulation (e.g. thrombin, fVIIa, fIXa) or the fibrinolytic cascades (e.g. plasminogen activators, plasmin). In another aspect, the present invention relates to novel monoamidino-containing compounds, their pharmaceutically acceptable salts, and pharmaceutically acceptable compositions thereof which are useful as potent and specific inhibitors of blood coagulation in mammals. In yet another aspect, the invention relates to methods for using these inhibitors as therapeutic agents for disease states in mammals characterized by coagulation disorders.

Background of the Invention

Hemostasis, the control of bleeding, occurs by surgical means, or by the physiological properties of vasoconstriction and coagulation. This invention is particularly concerned with blood coagulation and ways in which it assists in maintaining the integrity of mammalian circulation after injury, inflammation, disease, congenital defect, dysfunction or other disruption. Although platelets and blood coagulation are both involved in thrombus formation, certain components of the coagulation cascade are primarily responsible for the amplification or acceleration of the processes involved in platelet aggregation and fibrin deposition.

Thrombin is a key enzyme in the coagulation cascade as well as in hemostasis. Thrombin plays a central role in thrombosis through its ability to catalyze the conversion of fibrinogen into fibrin and through its potent platelet activation activity. Direct or indirect inhibition of thrombin activity has been the focus of a variety of recent anticoagulant strategies as reviewed by Claeson, G., "Synthetic Peptides and Peptidomimetics as Substrates and Inhibitors of Thrombin and Other Proteases in the Blood Coagulation System", Blood Coag. Fibrinol. 5, 411-436 (1994). Several classes of anticoagulants currently used in the clinic directly or indirectly affect thrombin (i.e. heparins, low-molecular weight heparins, heparin-like compounds and coumarins).



A prothrombinase complex, including Factor Xa (a serine protease, the activated form of its Factor X precursor and a member of the calcium ion binding, gamma carboxyglutamyl (Gla)-containing, vitamin K dependent, blood coagulation glycoprotein family), converts the zymogen prothrombin into the active procoagulant thrombin. Unlike thrombin, which acts on a variety of protein substrates as well as at a specific receptor, factor Xa appears to have a single physiologic substrate, namely prothrombin. Since one molecule of factor Xa may be able to generate up to 138 molecules of thrombin (Elodi et al., *Thromb. Res.* 15, 617-619 (1979)), direct inhibition of factor Xa as a way of indirectly inhibiting the formation of thrombin may be an efficient anticoagulant strategy. Therefore, it has been suggested that compounds which selectively inhibit factor Xa may be useful as *in vitro* diagnostic agents, or for therapeutic administration in certain thrombotic disorders, see *e.g.*, WO 94/13693.

Polypeptides derived from hematophagous organisms have been reported which are highly potent and specific inhibitors of factor Xa. United States Patent 4,588,587 describes anticoagulant activity in the saliva of the Mexican leech, *Haementeria officinalis*. A principal component of this saliva was shown to be the polypeptide factor Xa inhibitor, antistasin (ATS), by Nutt, E. *et al.*, "The Amino Acid Sequence of Antistasin, a Potent Inhibitor of Factor Xa Reveals a Repeated Internal Structure", *J. Biol. Chem.*, 263, 10162-10167 (1988). Another potent and highly specific inhibitor of Factor Xa, called tick anticoagulant peptide (TAP), has been isolated from the whole body extract of the soft tick *Ornithodoros moubata*, as reported by Waxman, L., *et al.*, "Tick Anticoagulant Peptide (TAP) is a Novel Inhibitor of Blood Coagulation Factor Xa" *Science*, 248, 593-596 (1990).

Factor Xa inhibitory compounds which are not large polypeptide-type inhibitors have also been reported including: Tidwell, R.R. *et al.*, "Strategies for Anticoagulation With Synthetic Protease Inhibitors. Xa Inhibitors Versus Thrombin Inhibitors", *Thromb. Res.*, 19, 339-349 (1980); Turner, A.D. *et al.*, "p-Amidino Esters as Irreversible Inhibitors of Factor IXa and Xa and Thrombin", *Biochemistry*, 25, 4929-4935 (1986); Hitomi, Y. *et al.*, "Inhibitory Effect of New Synthetic Protease Inhibitor (FUT-175) on the Coagulation System", *Haemostasis*, 15, 164-168 (1985); Sturzebecher, J. *et al.*, "Synthetic Inhibitors of Bovine Factor Xa and Thrombin. Comparison of Their Anticoagulant Efficiency", *Thromb. Res.*, 54, 245-252 (1989); Kam, C.M. *et al.*, "Mechanism Based Isocoumarin Inhibitors for Trypsin and Blood Coagulation Serine Proteases: New Anticoagulants", *Biochemistry*, 27, 2547-2557 (1988); Hauptmann, J. *et al.*, "Comparison of the Anticoagulant and Antithrombotic Effects of Synthetic Thrombin and Factor Xa Inhibitors", *Thromb. Haemost.*, 63, 220-223 (1990); and the like.

Others have reported Factor Xa inhibitors which are small molecule organic compounds, such as nitrogen containing heterocyclic compounds which have amidino

substituent groups, wherein two functional groups of the compounds can bind to Factor Xa at two of its active sites. For example, WO 98/28269 describes pyrazole compounds having a terminal C(=NH)-NH<sub>2</sub> group; WO 97/21437 describes benzimidazole compounds substituted by a basic radical which are connected to a naphthyl group via a straight or branched chain alkylene, -C(=O) or -S(=O)<sub>2</sub> bridging group; WO 99/10316 describes compounds having a 4-phenyl-N-alkylamidino-piperidine and 4-phenoxy-N-alkylamidino-piperidine group connected to a 3-amidinophenyl group via a carboxamidealkyleneamino bridge; and EP 798295 describes compounds having a 4-phenoxy-N-alkylamidino-piperidine group connected to an amidinonaphthyl group via a substituted or unsubstituted sulfonamide or carboxamide bridging group.

There exists a need for effective therapeutic agents for the regulation of hemostasis, and for the prevention and treatment of thrombus formation and other pathological processes in the vasculature induced by thrombin such as restenosis and inflammation. In particular, there continues to be a need for compounds which selectively inhibit factor Xa or its precursors. Compounds that have different combinations of bridging groups and functional groups than compounds previously discovered are needed, particularly compounds which selectively or preferentially bind to Factor Xa. Compounds with a higher degree of binding to Factor Xa than to thrombin are desired, especially those compounds having good bioavailability and/or solubility.


### Summary of the Invention

The present invention relates to novel compounds which inhibit factor Xa, their pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives, and pharmaceutically acceptable compositions thereof which have particular biological properties and are useful as potent and specific inhibitors of blood coagulation in mammals. Pharmaceutical compositions of the invention may be used to prevent or treat a condition in a mammal characterized by undesired thrombosis. In another aspect, the invention relates to methods of using these inhibitors as diagnostic reagents or as therapeutic agents for disease states in mammals which have coagulation disorders, such as in the treatment or prevention of a condition in a mammal characterized by undesired thrombosis such as, for example, any thrombotically mediated acute coronary or cerebrovascular syndrome, any thrombotic syndrome occurring in the venous system, any coagulopathy, and any thrombotic complications associated with extracorporeal circulation or instrumentation, and for the inhibition of coagulation of biological samples and blood in biological samples.

In certain embodiments, this invention relates to novel compounds which are potent and highly selective inhibitors of isolated factor Xa or factor Xa when assembled in the

prothrombinase complex. These compounds show selectivity for factor Xa versus other proteases of the coagulation cascade (e.g. thrombin, etc.) or the fibrinolytic cascade, and are useful as diagnostic reagents as well as antithrombotic agents.

In a preferred embodiment, the present invention provides a compound of the formula I:



wherein:

A is selected from:

- (a)  $C_1$ - $C_6$ -alkyl;
- (b)  $C_3$ - $C_8$ -cycloalkyl;
- (c)  $-N(-R^2, -R^3)$ ,  $R^3-C(=N-R^2)-$ ,  $(-R^2, -R^3)N-C(=N-R^2)-$ ,  $(-R^2, -R^3)N-C(=N-R^2)-N(-R)-$
- (d) phenyl, which is independently substituted with 0-2  $R^1$  substituents;
- (e) naphthyl, which is independently substituted with 0-2  $R^1$  substituents; and
- (f) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^1$  substituents;

$R^1$  is selected from:

Halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$ ,  $-NO_2$ ,  $R^2-C(=N-R^3)-$ ,  $(-R^2, -R^3)N-C(=N-R^2)-$ ,  $-(CH_2)_mNR^2R^3$ ,  $-C(=O)-N(-R^2, -R^3)$ ,  $-SO_2N(-R^2, -R^3)$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl,  $-CN$ ,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl and  $-NO_2$ ;

$R^2$  and  $R^3$  are independently selected from the group consisting of:

H,  $-OR^2$ ,  $-N(-R^2, -R^3)$ ,  $-C_{1-4}$ alkyl,  $-C_{2-6}$ alkenyl,  $-C_{2-6}$ alkynyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkylphenyl and  $-C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be



independently replaced with a member selected from the group consisting of halo,  $-C_{1-4}$ alkyl,  $-C_{2-6}$ alkenyl,  $-C_{2-6}$ alkynyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$ , and  $-NO_2$ ;

or  $R^2$  and  $R^3$  taken together can form a 3-8 membered cycloalkyl or a heterocyclic ring system, wherein the heterocyclic ring system may have from 5 to 10 ring atoms, with 1 to 2 rings being in the ring system and contain from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the heterocyclic ring system may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $-CN$ ,  $-C_{1-4}$ alkyl,  $-C_{2-6}$ alkenyl,  $-C_{2-6}$ alkynyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl and  $-NO_2$ ;

m is an integer of 0-2;

Q is a member selected from the group consisting of:

a direct link,  $-CH_2-$ ,  $-C(=O)-$ ,  $-N(R^4)-$ ,  $-N(R^4)CH_2-$ ,  $-C=N(R^4)-$ ,  $-C(=O)-N(R^4)-$ ,  $-N(R^4)-C(=O)-$ ,  $-SO_2-$ ,  $-O-$ ,  $-SO_2-N(R^4)-$  and  $-N(R^4)-SO_2-$ ;

$R^4$  is selected from:

H,  $-C_{1-4}$ alkyl,  $-C_{2-6}$ alkenyl,  $-C_{2-6}$ alkynyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkylphenyl and  $-C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $-C_{1-4}$ alkyl,  $-C_{2-6}$ alkenyl,  $-C_{2-6}$ alkynyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$ , and  $-NO_2$ ;

D is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1a}$  substituents; and
- (b) an aromatic six-membered heterocyclic ring having from 1-2 ring nitrogen atoms, and wherein the ring atoms may be substituted with 0-2  $R^{1a}$  substituents;

$R^{1a}$  is selected from:

Halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$ ,  $-NO_2$ ,  $(CH_2)_mNR^{2a}R^{3a}$ ,  $SO_2NR^{2a}R^{3a}$ ,  $SO_2R^{2a}$ ,  $CF_3$ ,  $OR^{2a}$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $-CN$  and  $-NO_2$ ;

$R^{2a}$  and  $R^{3a}$  are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>.

M is a member selected from the group consisting of:

-N(R<sup>16</sup>)-C(=O)-, -N(R<sup>16</sup>)-C(=S)-, -C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-, -C(-R<sup>19</sup>, -R<sup>19a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-, -C(-R<sup>17</sup>)=C(-R<sup>18</sup>)-C(=O)-, -C(-R<sup>17</sup>)=C(-R<sup>18</sup>)-C(=S)-, -C(-R<sup>17</sup>)=C(-R<sup>18</sup>)-, -O-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -O-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S(=O)<sub>2</sub>-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S(=O)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S(=O)<sub>2</sub>-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S(=O)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(=O)-C(=O)-, -N(R<sup>16</sup>)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -N(R<sup>16</sup>)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(=S)-C(=S)-, -C(=S)-C(=O)-, -C(=O)-C(=S)-, -N=C(-R<sup>17</sup>)-C(=O)-, -N=C(-R<sup>17</sup>)-C(=S)-, -C(-R<sup>17</sup>)=N-, -N(-R<sup>16</sup>)-C(=O)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -O-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -N(-R<sup>16</sup>)-C(=S)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -S(=O)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -S(=O)<sub>2</sub>-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -C(=C(R<sup>17b</sup>, -R<sup>17c</sup>))-C(=O)-, -C(=C(R<sup>17b</sup>, -R<sup>17c</sup>))-C(=S)-, -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(=S)-, -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-(N(-H, -R<sup>18b</sup>)), -R<sup>17a</sup>)-C(=O)-; -N=C(-R<sup>17</sup>)- and -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-(N(-H, -R<sup>18b</sup>)), -R<sup>17a</sup>)-C(=S)-; wherein the first named atom of the chain is directly attached to D, and wherein D, M and the N atom attached to the last chain atom of M collectively form a bicyclic ring structure;

R<sup>16</sup>, R<sup>17</sup>, R<sup>17a</sup>, R<sup>18</sup>, R<sup>18a</sup>, R<sup>18b</sup>, R<sup>19</sup>, and R<sup>19a</sup> are each independently selected from the group consisting of:

hydrogen, halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>R<sup>2</sup>, CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, -CN, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

R<sup>17b</sup> and R<sup>17c</sup> are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy, -C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>,

phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the cycloalkyl, the phenyl ring, or the aromatic heterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>-alkyl, -CN, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

E is a member selected from the group consisting of:

a direct link, -C(=O)-, -C(=O)-N(R<sup>5</sup>)-, -C(-R<sup>5a</sup>, -R<sup>6a</sup>)- and -C(-R<sup>5b</sup>, -R<sup>6b</sup>)-C(-R<sup>5c</sup>, -R<sup>6c</sup>)-;

wherein R<sup>5</sup>, R<sup>5a</sup>, R<sup>6a</sup>, R<sup>5b</sup>, R<sup>6b</sup>, R<sup>5c</sup> and R<sup>6c</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylheteroaryl, C<sub>1-4</sub>alkylCOOH and C<sub>1-4</sub>alkylCOOC<sub>1-4</sub>alkyl, wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -OH, -O-C<sub>1-4</sub>alkyl, -SH, -S-C<sub>1-4</sub>alkyl, -CN and -NO<sub>2</sub>;

G is selected from:

a direct link, -C(R<sup>7</sup>, R<sup>8</sup>)-, -C(R<sup>7a</sup>, R<sup>8a</sup>)-C(R<sup>7b</sup>, R<sup>8b</sup>)- and -C(R<sup>7c</sup>)=C(R<sup>8c</sup>)-;

wherein R<sup>7</sup>, R<sup>8</sup>, R<sup>7a</sup>, R<sup>8a</sup>, R<sup>7b</sup>, R<sup>8b</sup>, R<sup>7c</sup> and R<sup>8c</sup> are independently a member selected from from the group consisting of:

hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)OR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)-<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, a naturally occurring or synthetic amino acid side chain, and C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the C<sub>0-4</sub>alkylheterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>9</sup> and R<sup>10</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>3-8</sub>cycloalkyl, and C<sub>1-4</sub>alkyl-O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl-COOH wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>, and wherein R<sup>9</sup> and R<sup>10</sup> taken together can form a 5-8 membered heterocyclic ring;

J is a member selected from the group consisting of:

a direct link, -O-, -O-C(R<sup>11</sup>, -R<sup>11a</sup>)-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -S-C(R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)-C(R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)<sub>2</sub>-(R<sup>11</sup>, -R<sup>11a</sup>)-, -C(=O)-, -C(=O)-N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(=O)-, -N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(R<sup>11</sup>, -R<sup>11a</sup>)- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>11</sup> substituents;

R<sup>11</sup>, R<sup>11a</sup>, R<sup>11b</sup>, and R<sup>11</sup> are a member independently selected from the group consisting of:

hydrogen, halo, -CF<sub>3</sub>, -CN, -NR<sup>9</sup>R<sup>10</sup>, -SO<sub>2</sub>Me, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>2-6</sub>alkenyl, -O-C<sub>2-6</sub>alkynyl, -O-C<sub>3-8</sub>cycloalkyl, -O-C<sub>1-4</sub>alkyl-O-C<sub>1-4</sub>alkyl, -O-C<sub>1-4</sub>alkyl-COOH, -O-C<sub>1-4</sub>alkyl-phenyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>2-6</sub>alkenyl, -C(=O)-O-C<sub>2-6</sub>alkynyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, C<sub>0-4</sub>alkylC(=O)OR<sup>9</sup>, C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl; wherein from 1-4 hydrogen atoms on the C<sub>0-4</sub>alkylheterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-phenyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents;
- (b) naphthyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents; and



- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{1b}$  substituents;

$R^{1b}$  is a member selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN, -NO<sub>2</sub>,  $NR^{2b}R^{3b}$ ,  $SO_2NR^{2b}R^{3b}$ ,  $SO_2R^{2b}$ ,  $CF_3$ ,  $OR^{2b}$ ,  $O-CH_2-CH_2-OR^{2b}$ ,  $O-CH_2-COOR^{2b}$ ,  $N(R^{2b})-CH_2-CH_2-OR^{2b}$ ,  $N(-CH_2-CH_2-OR^{2b})_2$ ,  $N(R^{2b})-C(=O)R^{3b}$ ,  $N(R^{2b})-SO_2-R^{3b}$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl-phenyl, -CN and -NO<sub>2</sub>;

$R^{2b}$  and  $R^{3b}$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN and -NO<sub>2</sub>;

L is selected from:

H, -CN,  $C(=O)NR^{12}R^{13}$ ,  $-(CH_2)_nNR^{12}R^{13}$ ,  $C(=NR^{12})NR^{12}R^{13}$ ,  $OR^{12}$ ,  $-NR^{12}C(=NR^{12})NR^{12}R^{13}$ , and  $NR^{12}C(=NR^{12})-R^{13}$ ;

n is an integer from 0 to 8;

$R^{12}$  and  $R^{13}$  are independently selected from:

hydrogen, -OR<sup>14</sup>, -NR<sup>14</sup>R<sup>15</sup>,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $COOC_{1-4}$ alkyl,  $COO-C_{0-4}$ alkylphenyl and  $COO-C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH, -O- $C_{1-4}$ alkyl,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN, and -NO<sub>2</sub>;

$R^{14}$  and  $R^{15}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring

atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

In certain aspects of this invention, compounds are provided which are useful as diagnostic reagents. In another aspect, the present invention includes pharmaceutical compositions comprising a pharmaceutically effective amount of the compounds of this invention and a pharmaceutically acceptable carrier. In yet another aspect, the present invention includes methods comprising using the above compounds and pharmaceutical compositions for preventing or treating disease states characterized by disorders of the blood coagulation process in mammals, or for preventing coagulation in stored blood products and samples. Optionally, the methods of this invention comprise administering the pharmaceutical composition in combination with an additional therapeutic agent such as an antithrombotic and/or a thrombolytic agent and/or an anticoagulant.

The preferred compounds also include their pharmaceutically acceptable isomers, hydrates, solvates, salts and prodrug derivatives.

### Detailed Description of the Invention

#### Definitions

In accordance with the present invention and as used herein, the following terms are defined with the following meanings, unless explicitly stated otherwise.

The term "alkenyl" refers to a trivalent straight chain or branched chain unsaturated aliphatic radical. The term "alkinyl" (or "alkynyl") refers to a straight or branched chain aliphatic radical that includes at least two carbons joined by a triple bond. If no number of carbons is specified alkenyl and alkinyl each refer to radicals having from 2-12 carbon atoms.

The term "alkyl" refers to saturated aliphatic groups including straight-chain, branched-chain and cyclic groups having the number of carbon atoms specified, or if no number is specified, having up to 12 carbon atoms. The term "cycloalkyl" as used herein refers to a mono-, bi-, or tricyclic aliphatic ring having 3 to 14 carbon atoms and preferably 3 to 7 carbon atoms.

As used herein, the terms "carbocyclic ring structure" and "C<sub>3-16</sub> carbocyclic mono, bicyclic or tricyclic ring structure" or the like are each intended to mean stable ring structures

having only carbon atoms as ring atoms wherein the ring structure is a substituted or unsubstituted member selected from the group consisting of: a stable monocyclic ring which is aromatic ring ("aryl") having six ring atoms; a stable monocyclic non-aromatic ring having from 3 to 7 ring atoms in the ring; a stable bicyclic ring structure having a total of from 7 to 12 ring atoms in the two rings wherein the bicyclic ring structure is selected from the group consisting of ring structures in which both of the rings are aromatic, ring structures in which one of the rings is aromatic and ring structures in which both of the rings are non-aromatic; and a stable tricyclic ring structure having a total of from 10 to 16 atoms in the three rings wherein the tricyclic ring structure is selected from the group consisting of: ring structures in which three of the rings are aromatic, ring structures in which two of the rings are aromatic and ring structures in which three of the rings are non-aromatic. In each case, the non-aromatic rings when present in the monocyclic, bicyclic or tricyclic ring structure may independently be saturated, partially saturated or fully saturated. Examples of such carbocyclic ring structures include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), 2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin). Moreover, the ring structures described herein may be attached to one or more indicated pendant groups via any carbon atom which results in a stable structure. The term "substituted" as used in conjunction with carbocyclic ring structures means that hydrogen atoms attached to the ring carbon atoms of ring structures described herein may be substituted by one or more of the substituents indicated for that structure if such substitution(s) would result in a stable compound.

The term "aryl" which is included with the term "carbocyclic ring structure" refers to an unsubstituted or substituted aromatic ring, substituted with one, two or three substituents selected from loweralkoxy, loweralkyl, loweralkylamino, hydroxy, halogen, cyano, hydroxyl, mercapto, nitro, thioalkoxy, carboxaldehyde, carboxyl, carboalkoxy and carboxamide, including but not limited to carbocyclic aryl, heterocyclic aryl, and biaryl groups and the like, all of which may be optionally substituted. Preferred aryl groups include phenyl, halophenyl, loweralkylphenyl, naphthyl, biphenyl, phenanthrenyl and naphthacenyl.

The term "arylalkyl" which is included with the term "carbocyclic aryl" refers to one, two, or three aryl groups having the number of carbon atoms designated, appended to an alkyl group having the number of carbon atoms designated. Suitable arylalkyl groups include, but are not limited to, benzyl, picolyl, naphthylmethyl, phenethyl, benzyhydryl, trityl, and the like, all of which may be optionally substituted.

As used herein, the term "heterocyclic ring" or "heterocyclic ring system" is intended to mean a substituted or unsubstituted member selected from the group consisting of stable

monocyclic ring having from 5-7 members in the ring itself and having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S; a stable bicyclic ring structure having a total of from 7 to 12 atoms in the two rings wherein at least one of the two rings has from 1 to 4 hetero atoms selected from N, O and S, including bicyclic ring structures wherein any of the described stable monocyclic heterocyclic rings is fused to a hexane or benzene ring; and a stable tricyclic heterocyclic ring structure having a total of from 10 to 16 atoms in the three rings wherein at least one of the three rings has from 1 to 4 hetero atoms selected from the group consisting of N, O and S. Any nitrogen and sulfur atoms present in a heterocyclic ring of such a heterocyclic ring structure may be oxidized. Unless indicated otherwise the terms "heterocyclic ring" or "heterocyclic ring system" include aromatic rings, as well as non-aromatic rings which can be saturated, partially saturated or fully saturated non-aromatic rings. Also, unless indicated otherwise the term "heterocyclic ring system" includes ring structures wherein all of the rings contain at least one hetero atom as well as structures having less than all of the rings in the ring structure containing at least one hetero atom, for example bicyclic ring structures wherein one ring is a benzene ring and one of the rings has one or more hetero atoms are included within the term "heterocyclic ring systems" as well as bicyclic ring structures wherein each of the two rings has at least one hetero atom. Moreover, the ring structures described herein may be attached to one or more indicated pendant groups via any hetero atom or carbon atom which results in a stable structure. Further, the term "substituted" means that one or more of the hydrogen atoms on the ring carbon atom(s) or nitrogen atom(s) of the each of the rings in the ring structures described herein may be replaced by one or more of the indicated substituents if such replacement(s) would result in a stable compound. Nitrogen atoms in a ring structure may be quaternized, but such compounds are specifically indicated or are included within the term "a pharmaceutically acceptable salt" for a particular compound. When the total number of O and S atoms in a single heterocyclic ring is greater than 1, it is preferred that such atoms not be adjacent to one another. Preferably, there are no more than 1 O or S ring atoms in the same ring of a given heterocyclic ring structure.

Examples of monocyclic and bicyclic heterocyclic ring systems, in alphabetical order, are acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalinyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolyl, imidazolyl, 1H-indazolyl, indolinyl, indoliziny, indolyl, 3H-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl (benzimidazolyl), isothiazolyl, isoxazolyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolidinyl, pyrimidinyl, phenanthridinyl,



phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyroazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridoxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2H-pyrrolyl, pyrrolyl, quinazolinyl, quinolinyl, 4H-quinoliziny, quinoxaliny, quinuclidinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl and xanthenyl. Preferred heterocyclic ring structures include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrrolidinyl, imidazolyl, indolyl, benzimidazolyl, 1H-indazolyl, oxazoliny, or isatinoyl. Also included are fused ring and spiro compounds containing, for example, the above heterocyclic ring structures.

As used herein the term "aromatic heterocyclic ring system" has essentially the same definition as for the monocyclic and bicyclic ring systems except that at least one ring of the ring system is an aromatic heterocyclic ring or the bicyclic ring has an aromatic or non-aromatic heterocyclic ring fused to an aromatic carbocyclic ring structure.

The terms "halo" or "halogen" as used herein refer to Cl, Br, F or I substituents. The term "haloalkyl", and the like, refer to an aliphatic carbon radicals having at least one hydrogen atom replaced by a Cl, Br, F or I atom, including mixtures of different halo atoms. Trihaloalkyl includes trifluoromethyl and the like as preferred radicals, for example.

The term "methylene" refers to  $-\text{CH}_2-$ . The term "Bu" refers to "butyl" or  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ ; the term "Ph" refers to "phenyl"; the term "Me" refers to "methyl" or  $-\text{CH}_3$ ; the term "Et" refers to "ethyl" or  $-\text{CH}_2\text{CH}_3$ ; the term "Bu(t)" or "t-Bu" refers to "tert-butyl" or  $-\text{C}(\text{CH}_3)_3$ .

The term "amino acid side chain" includes any naturally occurring or synthetic side chain, *i.e.* the side chain "R" of an amino acid having the formula:  $\text{NH}_2-\text{CHR}-\text{COOH}$ . Synthetic amino acids include both (R) and (S) enantiomers, as well as derivatives of the naturally-occurring amino acid side chains. The side chain may be any synthetic amino acid side chain known in the art, including but not limited to, those produced by recombinant DNA techniques, fermentation, or by solid phase synthesis techniques. Such synthetic amino acid side chains may contain one or more functional groups selected from the following: alkyl, aryl, alkenyl, alkynyl, thiols, primary and secondary amines, aldehydes, carboxylates, nitriles, aromatic amines, aromatic carboxylates, primary alcohols, and the like.

The term "pharmaceutically acceptable salts" includes salts of compounds derived from the combination of a compound and an organic or inorganic acid. These compounds are useful

in both free base and salt form. In practice, the use of the salt form amounts to use of the base form; both acid and base addition salts are within the scope of the present invention.

"Pharmaceutically acceptable acid addition salt" refers to salts retaining the biological effectiveness and properties of the free bases and which are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

"Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium and magnesium salts. Salts derived from pharmaceutically acceptable organic nontoxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-diethylaminoethanol, trimethamine, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like. Particularly preferred organic nontoxic bases are isopropylamine, diethylamine, ethanolamine, trimethamine, dicyclohexylamine, choline, and caffeine.


"Biological property" for the purposes herein means an *in vivo* effector or antigenic function or activity that is directly or indirectly performed by a compound of this invention that are often shown by *in vitro* assays. Effector functions include receptor or ligand binding, any enzyme activity or enzyme modulatory activity, any carrier binding activity, any hormonal activity, any activity in promoting or inhibiting adhesion of cells to an extracellular matrix or cell surface molecules, or any structural role. Antigenic functions include possession of an epitope or antigenic site that is capable of reacting with antibodies raised against it.

In the compounds of this invention, carbon atoms bonded to four non-identical substituents are asymmetric. Accordingly, the compounds may exist as diastereoisomers, enantiomers or mixtures thereof. The syntheses described herein may employ racemates, enantiomers or diastereomers as starting materials or intermediates. Diastereomeric products resulting from such syntheses may be separated by chromatographic or crystallization methods, or by other methods known in the art. Likewise, enantiomeric product mixtures may be separated using the same techniques or by other methods known in the art. Each of the

asymmetric carbon atoms, when present in the compounds of this invention, may be in one of two configurations (R or S) and both are within the scope of the present invention.

### Preferred Embodiments

In a preferred embodiment, the present invention provides a compound according to the formula I:



wherein:

A is selected from:

- (a) C<sub>1</sub>-C<sub>6</sub>-alkyl;
- (b) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;
- (c) phenyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;
- (d) naphthyl, which is independently substituted with 0-2 R<sup>1</sup> substituents; and
- (e) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from the group consisting of N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1</sup> substituents;

R<sup>1</sup> is selected from:

halo, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, -C<sub>1-4</sub>alkyl, -CN, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

R<sup>2</sup> and R<sup>3</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring

atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

m is an integer of 0-2;

Q is a member selected from the group consisting of:

a direct link, -C(=O)-, -N(R<sup>4</sup>)-, -C(=O)-N(R<sup>4</sup>)-, -N(R<sup>4</sup>)-C(=O)-, -SO<sub>2</sub>-, -O-, -SO<sub>2</sub>-N(R<sup>4</sup>)- and -N(R<sup>4</sup>)-SO<sub>2</sub>-;

R<sup>4</sup> is selected from:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

D is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1a</sup> substituents; and
- (b) an aromatic six-membered heterocyclic ring having from 1-2 ring nitrogen atoms, and wherein the ring atoms may be substituted with 0-2 R<sup>1a</sup> substituents;

R<sup>1a</sup> is selected from:


halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2a</sup>R<sup>3a</sup>, SO<sub>2</sub>NR<sup>2a</sup>R<sup>3a</sup>, SO<sub>2</sub>R<sup>2a</sup>, CF<sub>3</sub>, OR<sup>2a</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>2a</sup> and R<sup>3a</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

M, D and N collectively form a bicyclic ring structure selected from the group consisting of:





and the like, wherein the aromatic carbocyclic ring corresponding to the D portion for each of the bicyclic rings may be replaced with an aromatic heterocyclic ring as defined above for D, and wherein 0 to 2 of the hydrogen atoms on the D portion of the bicyclic ring may be replaced by  $R^{1a}$  substituents as defined above;

$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN, -NO<sub>2</sub>,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$ ,  $SO_2R^2$ , CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl, -CN,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl and -NO<sub>2</sub>;

$R^{17b}$  and  $R^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy,  $-C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $-C_{2-6}$ alkynyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl, -CN,  $-NO_2$ ,  $-(CH_2)_mNR^2R^3$ ,  $-SO_2NR^2R^3$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the cycloalkyl, the phenyl ring, or the aromatic heterocyclic ring may be independently replaced with a member selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl, -CN,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl and  $-NO_2$ ;

E is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-(C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-)$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $C_{0-4}$ alkylheteroaryl,  $C_{1-4}$ alkylCOOH and  $C_{1-4}$ alkylCOOC $C_{1-4}$ alkyl, wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl, -OH,  $-O-C_{1-4}$ alkyl, -SH,  $-S-C_{1-4}$ alkyl, -CN and  $-NO_2$ ;

G is selected from:

a direct link,  $-C(R^7, R^8)-$ ,  $-C(R^{7a}, R^{8a})-C(R^{7b}, R^{8b})-$  and  $-C(R^{7c})=C(R^{8c})-$ ;

wherein  $R^7$ ,  $R^8$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{7b}$ ,  $R^{8b}$ ,  $R^{7c}$  and  $R^{8c}$  are independently a member selected from from the group consisting of:

hydrogen, halogen,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $-OR^9$ ,  $-N(R^9R^{10})$ ,  $-C_{0-4}$ alkylCOOR<sup>9</sup>,  $-C_{0-4}$ alkylC(=O)NR<sup>9</sup>R<sup>10</sup>,  $-C_{0-4}$ alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>,  $-C_{0-4}$ alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)<sub>2</sub>,  $-N(R^9)COR^{10}$ ,  $-N(R^9)C(=O)R^{10}$ ,  $-N(R^9)SO_2R^{10}$ , and a naturally occurring or synthetic amino acid side chain, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $-OR^9$ ,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl, -CN and  $-NO_2$ ;

$R^9$  and  $R^{10}$  are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>, and wherein R<sup>9</sup> and R<sup>10</sup> taken together can form a 5-8 membered heterocyclic ring;

J is a member selected from the group consisting of:

a direct link, -O-, -O-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -S-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)<sub>2</sub>-(-R<sup>11</sup>, -R<sup>11a</sup>)-, -C(=O)-, -C(=O)-N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(=O)-, -N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(-R<sup>11</sup>, -R<sup>11a</sup>)- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>11c</sup> substituents;

R<sup>11</sup>, R<sup>11a</sup>, R<sup>11b</sup> and R<sup>11c</sup> are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>2-6</sub>alkenyl, -O-C<sub>2-6</sub>alkynyl, -O-C<sub>3-8</sub>cycloalkyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>2-6</sub>alkenyl, -C(=O)-O-C<sub>2-6</sub>alkynyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents;
- (b) naphthyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents; and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1b</sup> substituents;

R<sup>1b</sup> is a member selected from the group consisting of:

halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>R<sup>2b</sup>, CF<sub>3</sub>, OR<sup>2b</sup>, O-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, O-CH<sub>2</sub>-COOR<sup>2b</sup>, N(R<sup>2b</sup>)-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, N(-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>)<sub>2</sub>, N(R<sup>2b</sup>)-C(=O)R<sup>3b</sup>, N(R<sup>2b</sup>)-SO<sub>2</sub>-R<sup>3b</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic

system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>2b</sup> and R<sup>3b</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

L is selected from:

H, -CN, C(=O)NR<sup>12</sup>R<sup>13</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup>, C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, OR<sup>12</sup>, -NR<sup>12</sup>C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, and NR<sup>12</sup>C(=NR<sup>12</sup>)-R<sup>13</sup>;

n is an integer from 0 to 8;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

hydrogen, -OR<sup>14</sup>, -NR<sup>14</sup>R<sup>15</sup>, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, COOC<sub>1-4</sub>alkyl, COO-C<sub>0-4</sub>alkylphenyl and COO-C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH, -O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;


R<sup>14</sup> and R<sup>15</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

In a further preferred embodiment, the present invention provides a compound according to the formula I:





wherein:

A is selected from:

- (a) C<sub>1</sub>-C<sub>6</sub>-alkyl;
- (b) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;
- (c) phenyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;
- (d) naphthyl, which is independently substituted with 0-2 R<sup>1</sup> substituents; and
- (e) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1</sup> substituents;

R<sup>1</sup> is selected from:

halo, C<sub>1-4</sub>alkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>R<sup>2</sup>, CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

R<sup>2</sup> and R<sup>3</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

m is an integer of 0-2;

B is a member selected from the group consisting of:

a direct link, -C(=O)-, -N(R<sup>4</sup>)-, -C(=O)-N(R<sup>4</sup>)-, -N(R<sup>4</sup>)-C(=O)-, -SO<sub>2</sub>-, -O-, -SO<sub>2</sub>-N(R<sup>4</sup>)- and -N(R<sup>4</sup>)-SO<sub>2</sub>-;

R<sup>4</sup> is selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

D is phenyl, which is independently substituted with 0-2 R<sup>1a</sup> substituents;


R<sup>1a</sup> is selected from:

halo, C<sub>1-4</sub>alkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2a</sup>R<sup>3a</sup>, -SO<sub>2</sub>NR<sup>2a</sup>R<sup>3a</sup>, -SO<sub>2</sub>R<sup>2a</sup>, CF<sub>3</sub>, -OR<sup>2a</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

$R^{2a}$  and  $R^{3a}$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;

M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



wherein 0 to 2 of the hydrogen atoms on the D portion of the bicyclic ring may be replaced by  $R^{1a}$  substituents as defined above;

$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $-CN$ ,  $-NO_2$ ,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$ ,  $SO_2R^2$ ,  $CF_3$ ,  $OR^2$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

$R^{17b}$  and  $R^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy, -C<sub>1-4</sub>alkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S;

m is an integer from 0-6;

E is a member selected from the group consisting of:

a direct link, -C(=O)-, -C(=O)-N(R<sup>5</sup>)-, -C(-R<sup>5a</sup>, -R<sup>6a</sup>)- and -(-C(-R<sup>5b</sup>, -R<sup>6b</sup>)-C(-R<sup>5c</sup>, -R<sup>6c</sup>)-;

wherein R<sup>5</sup>, R<sup>5a</sup>, R<sup>6a</sup>, R<sup>5b</sup>, R<sup>6b</sup>, R<sup>5c</sup> and R<sup>6c</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylheteroaryl, C<sub>1-4</sub>alkylCOOH and C<sub>1-4</sub>alkylCOOC<sub>1-4</sub>alkyl, wherein from 0-2 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, -OH, -O-C<sub>1-4</sub>alkyl, -SH, -S-C<sub>1-4</sub>alkyl, -CN and -NO<sub>2</sub>;

G is selected from:

a direct link, -C(R<sup>7</sup>, R<sup>8</sup>)-, -C(R<sup>7a</sup>, R<sup>8a</sup>)-C(R<sup>7b</sup>, R<sup>8b</sup>)- and -C(R<sup>7c</sup>)=C(R<sup>8c</sup>)-;

wherein R<sup>7</sup>, R<sup>8</sup>, R<sup>7a</sup>, R<sup>8a</sup>, R<sup>7b</sup>, R<sup>8b</sup>, R<sup>7c</sup> and R<sup>8c</sup> are independently a member selected from from the group consisting of:

hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)-<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, and a naturally occurring or synthetic amino acid side chain;

R<sup>9</sup> and R<sup>10</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

J is a member selected from the group consisting of:

a direct link, -O-, -O-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S-, -S(=O)<sub>2</sub>-, -S-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)<sub>2</sub>-(-R<sup>11</sup>, -R<sup>11a</sup>)-, -C(=O)-N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(-R<sup>11</sup>, -R<sup>11a</sup>)- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>11c</sup> substituents;

R<sup>11</sup>, R<sup>11a</sup>, R<sup>11b</sup> and R<sup>11c</sup> are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>3-8</sub>cycloalkyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, and a C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents;
- (b) naphthyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents; and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1b</sup> substituents;

R<sup>1b</sup> is a member selected from the group consisting of:

halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>R<sup>2b</sup>, CF<sub>3</sub>, OR<sup>2b</sup>, O-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, O-CH<sub>2</sub>-COOR<sup>2b</sup>, N(R<sup>2b</sup>)-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, N(-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>)<sub>2</sub>, N(R<sup>2b</sup>)-C(=O)R<sup>3b</sup>, N(R<sup>2b</sup>)-SO<sub>2</sub>-R<sup>3b</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>2b</sup> and R<sup>3b</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

L is selected from:

H, -CN, C(=O)NR<sup>12</sup>R<sup>13</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup>, C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, OR<sup>12</sup>, -NR<sup>12</sup>C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, and NR<sup>12</sup>C(=NR<sup>12</sup>)-R<sup>13</sup>;

n is an integer from 0 to 6;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

hydrogen, -OR<sup>14</sup>, -NR<sup>14</sup>R<sup>15</sup>, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, COOC<sub>1-4</sub>alkyl, COO-C<sub>0-4</sub>alkylphenyl and COO-C<sub>0-4</sub>alkylnaphthyl, wherein from 0-4 hydrogen atoms


on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH, -O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

R<sup>14</sup> and R<sup>15</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

In a still further preferred embodiment, the present invention provides a compound according to the formula I:



wherein:

A is selected from:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1</sup> substituents; and
- (b) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1</sup> substituents;

R<sup>1</sup> is selected from:

halo, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup> and SO<sub>2</sub>R<sup>2</sup>;

R<sup>2</sup> and R<sup>3</sup> are independently selected from the group consisting of:

H and C<sub>1-4</sub>alkyl;

m is an integer of 0-2;

Q is a member selected from the group consisting of:


a direct link, -C(=O)-, -SO<sub>2</sub>-, and -O-;

D is phenyl, which is independently substituted with 0-2 R<sup>1a</sup> substituents;

R<sup>1a</sup> is selected from:

halo and C<sub>1-4</sub>alkyl;

M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $-CN$ ,  $-NO_2$ ,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$ ,  $SO_2R^2$ ,  $CF_3$  and  $OR^2$ ;

$R^{17b}$  and  $R^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy,  $-C_{1-4}$ alkyl,  $-CN$ ,  $-NO_2$ ,  $-(CH_2)_mNR^2R^3$ ,  $-SO_2NR^2R^3$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-3 N atoms;

E is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-(C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-)$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:



H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylheteroaryl, C<sub>1-4</sub>alkylCOOH and C<sub>1-4</sub>alkylCOOC<sub>1-4</sub>alkyl;

G is selected from:

a direct link, -C(R<sup>7</sup>,R<sup>8</sup>)-, -C(R<sup>7a</sup>,R<sup>8a</sup>)-C(R<sup>7b</sup>,R<sup>8b</sup>)- and -C(R<sup>7c</sup>)=C(R<sup>8c</sup>)-;

wherein R<sup>7</sup>, R<sup>8</sup>, R<sup>7a</sup>, R<sup>8a</sup>, R<sup>7b</sup>, R<sup>8b</sup>, R<sup>7c</sup> and R<sup>8c</sup> are independently a member selected from the group consisting of:

hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, and a naturally occurring or synthetic amino acid side chain;

R<sup>9</sup> and R<sup>10</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

J is a member selected from the group consisting of:

a direct link, -O-, -S-, -C(=O)-N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(-R<sup>11</sup>, -R<sup>11a</sup>)- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>11c</sup> substituents;

R<sup>11</sup>, R<sup>11a</sup>, R<sup>11b</sup> and R<sup>11c</sup> are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>3-8</sub>cycloalkyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, and a C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents;
- (b) an aromatic heterocyclic ring having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring may be substituted with 0-2 R<sup>1b</sup> substituents;
- (c) a fused aromatic bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S,

and wherein the bicyclic ring system may be substituted with 0-2  $R^{1b}$  substituents;

$R^{1b}$  is a member selected from the group consisting of:

halo,  $-C_{1-4}$ alkyl,  $-OH$ ,  $-OBn$ ,  $-O-CH_2-CH_2-OH$ ,  $-O-CH_2-CH_2-OCH_3$ ,  $-O-CH_2-COOH$ ,  $-O-CH_2-C(=O)-O-CH_3$ ,  $-NH_2$ ,  $-NH-CH_2-CH_2-O-CH_3$ ,  $-NH-C(=O)-O-CH_3$  and  $-NH-SO_2-CH_3$ ;

L is selected from:

H,  $-C(=O)NR^{12}R^{13}$ ,  $-(CH_2)_nNR^{12}R^{13}$  and  $-C(=NR^{12})NR^{12}R^{13}$ ;


n is an integer from 0 to 6;

$R^{12}$  and  $R^{13}$  are independently selected from:

hydrogen and  $C_{1-4}$ alkyl;


and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

In yet another preferred embodiment, the present invention provides a compound according to formula I:




wherein:

A is a member selected from the group consisting of:



D is a member selected from the group consisting of:





E is a member selected from the group consisting of:

a direct link,  $-\text{C}(=\text{O})-$ ,  $-\text{C}(=\text{O})-\text{N}(\text{R}^5)-$ ,  $-\text{C}(\text{R}^{5a}, \text{R}^{6a})-$  and  $-(\text{C}(\text{R}^{5b}, \text{R}^{6b})-\text{C}(\text{R}^{5c}, \text{R}^{6c})-$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylheteroaryl, C<sub>1-4</sub>alkylCOOH and C<sub>1-4</sub>alkylCOOC<sub>1-4</sub>alkyl;

G is selected from:

a direct link,  $-C(R^7, R^8)-$ ,  $-C(R^{7a}, R^{8a})-C(R^{7b}, R^{8b})-$  and  $-C(R^{7c})=C(R^{8c})-$ ;

wherein  $R^7$ ,  $R^8$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{7b}$ ,  $R^{8b}$ ,  $R^{7c}$  and  $R^{8c}$  are independently a member selected from from the group consisting of:

hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, and a naturally occurring or synthetic amino acid side chain;

$R^9$  and  $R^{10}$  are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;


J is a member selected from the group consisting of:

a direct link, -O-, -S-, -C(=O)-N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(-R<sup>11</sup>, -R<sup>11a</sup>)- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>11c</sup> substituents;

R<sup>11</sup>, R<sup>11a</sup>, R<sup>11b</sup> and R<sup>11c</sup> are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -C<sub>1-4</sub>alkyl, -COOH, phenyl, and benzyl wherein the aromatic ring of the phenyl or benzyl is substituted with 0-2 members independently selected from the group consisting of halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -C<sub>1-4</sub>alkyl, -COOH and -C(=O)-O-C<sub>1-4</sub>alkyl;

Y and L taken together are a member selected from the group consisting of:



M and Q are as defined elsewhere in the specification;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

The following non-limiting tables illustrate representative compounds of the present invention wherein the "Y-L" portions for each of the formula in each of the tables are taken together and are independently selected from the group consisting of:






Table 1



Formula II

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 1a



Formula IIa


R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 1b



R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 1c




Formula IIc

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 1d



Formula IIId

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 1e



Formula IIe

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 1f



Formula II f

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 1g



Formula IIg



































R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 1h




Formula IIh

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 1j



Formula IIj

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 1k



Formula IIk


















R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et




Table 1(I)



Formula II(I)

$\text{R}^7$	$\text{R}^{11b}$	$\text{R}^7$	$\text{R}^{11b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 1m



Formula IIIm

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 1n



Formula IIIn


















R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 1o



Formula IIo


















R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 1p



Formula Iip

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 1q



Formula IIq

R <sup>7</sup>	R <sup>11b</sup>	R <sup>7</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 2



Formula III

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 2a




Formula IIIa

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl




Table 2b




R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 2c



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2d



Formula IIIId


R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 2e



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 2f



Formula IIIf














$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 2g



Formula IIIg














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2h



Formula IIIh

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 2i



Formula IIIi














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl



Table 2j



Formula IIIj














$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2k



Formula IIIk

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2(l)



Formula III(l)

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2m



Formula IIIIm

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 2n
















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 2o



Formula IIIo














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2p



Formula IIIp

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 2q



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl




Table 3



Formula IV

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3a



Formula VIa


R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 3b



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3c



Formula VIc

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3d



Formula IVd

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 3e



Formula IVe


















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3f



Formula IVf

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 3g




Formula IVg

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 3h




Formula IVh

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 3j



Formula IVj


















$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3k



Formula IVk

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3(l)



Formula IV(l)

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 3m



Formula IVm

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 3n




















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 3o




Formula IVo

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



Table 3p



Formula IVp


















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 3q



Formula IVq


















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 4



Formula V

$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4a



Formula Va


















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 4b




















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 4c



Formula Vc


















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4d



Formula Vd


















$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 4e



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et




Table 4f



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4g



Formula Vg



















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 4h



Formula Vh

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4j



Formula Vj


















$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 4k



Formula Vk

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 4(l)



Formula V(l)

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4m




Formula Vm

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



Table 4n



Formula Vn


















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 4o



Formula Vo


R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4p



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 4q




















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 5



Formula VI

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 5a



Formula VIa


















$\text{R}^{7c}$	$\text{R}^{8c}$	$\text{R}^{7c}$	$\text{R}^{8c}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 5b



Formula VIb


















R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 5c




Formula VIc

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



Table 5d



Formula VIId


















$\text{R}^{7c}$	$\text{R}^{8c}$	$\text{R}^{7c}$	$\text{R}^{8c}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 5e



Formula VIe


















R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 5f



Formula VI f


















R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 5g



Formula VIg


R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 5h



Formula VIh

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



Formula VII


















$\text{R}^{7c}$	$\text{R}^{8c}$	$\text{R}^{7c}$	$\text{R}^{8c}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 5j



Formula VIj

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 5k




Formula VIk

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



Table 5(l)



Formula VI(l)


















$\text{R}^{7c}$	$\text{R}^{8c}$	$\text{R}^{7c}$	$\text{R}^{8c}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 5m



Formula VIIm

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 5n




















R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et

Table 5o



Formula VIo


















$\text{R}^{7c}$	$\text{R}^{8c}$	$\text{R}^{7c}$	$\text{R}^{8c}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 5p



Formula VIp


















R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 5q



Formula VIq

R <sup>7c</sup>	R <sup>8c</sup>	R <sup>7c</sup>	R <sup>8c</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6



Formula VII

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 6a




Formula VIIa

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 6b



Formula VIIb

$\text{R}^{7a}$	$\text{R}^{7b}$	$\text{R}^{7a}$	$\text{R}^{7b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6c



Formula VIIc

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6d



Formula VIId

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6e



Formula VIIe

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6f



Formula VIIf

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 6g



Formula VIIg


R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 6h



Formula VIIh

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Formula VIIi

$R^{7a}$	$R^{7b}$	$R^{7a}$	$R^{7b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 6j



Formula VIIj

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6k



Formula VIIk

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6(l)



Formula VII(l)

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6m



Formula VIIm

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6n



Formula VIIIn

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6o



Formula VIIo

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 6p



Formula VIIp

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 6q




Formula VIIq

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et




Table 7



Formula VIII

$\text{R}^{7a}$	$\text{R}^{7b}$	$\text{R}^{7a}$	$\text{R}^{7b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7a



Formula VIIIa


R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 7b




Formula VIIIb

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7d



Formula VIIIId

$\text{R}^{7a}$	$\text{R}^{7b}$	$\text{R}^{7a}$	$\text{R}^{7b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7e



Formula VIIIe

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7f



Formula VIII f

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 7g




Formula VIIIg

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 7h




Formula VIIIh

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et



R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7j



Formula VIIIj

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7k



Formula VIIIk

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7(I)



Formula VIII(I)

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH <sub>2</sub>	H
		OCH <sub>2</sub> Ph	H
		OCH <sub>2</sub> CH <sub>2</sub> OMe	H
		H	Et
		Me	Et


Table 7m



Formula VIII m

$\text{R}^{7a}$	$\text{R}^{7b}$	$\text{R}^{7a}$	$\text{R}^{7b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7n



Formula VIIIIn

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et

Table 7o




Formula VIIIo

$\text{R}^{7a}$	$\text{R}^{7b}$	$\text{R}^{7a}$	$\text{R}^{7b}$
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et




Table 7p



Formula VIIIp

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 7q



Formula VIIIq

R <sup>7a</sup>	R <sup>7b</sup>	R <sup>7a</sup>	R <sup>7b</sup>
H	H	OMe	H
Me	Me	OMe	Me
		F	H
		-OH	Me
		Br	Me
		-NH2	H
		OCH2Ph	H
		OCH2CH2OMe	H
		H	Et
		Me	Et


Table 8



Formula IX

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8a



Formula IXa

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8b



Formula IXb

$\text{R}^{6c}$	$\text{R}^{11b}$	$\text{R}^{6c}$	$\text{R}^{11b}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 8c



Formula IXc

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8d



Formula IXd

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 8e




Formula IXe

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl




Table 8f



Formula IXf

$\text{R}^{6\text{c}}$	$\text{R}^{11\text{b}}$	$\text{R}^{6\text{c}}$	$\text{R}^{11\text{b}}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 8g



Formula IXg


R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 8h




Formula IXh

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl



R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8j



Formula IXj

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 8k



Formula IXk

$\text{R}^{6c}$	$\text{R}^{11b}$	$\text{R}^{6c}$	$\text{R}^{11b}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 8(I)



Formula IX(I)

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 8m




Formula IXm

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl




Table 8n



Formula IXn

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 8o



Formula IXo

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 8p



Formula IXp

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 8q



Formula IXq

R <sup>6c</sup>	R <sup>11b</sup>	R <sup>6c</sup>	R <sup>11b</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl


Table 9



Formula X

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 9a



Formula Xa


R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9b



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9c




Formula Xc

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl




Table 9d



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9e



Formula Xe














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9f
















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9g



Formula Xg















$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9h




Formula Xh

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9j



Formula Xj














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 9k



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl



Table 9(l)



Formula X(l)















$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl

Table 9m




R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9n



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 9o



Formula Xo

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9p



Formula Xp















R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 9q



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10



Formula XI

$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10a




Formula XIa

$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl




Table 10b



$R^{11c1}$	$R^{11c2}$	$R^{11c1}$	$R^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10c



Formula XIc

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10d



Formula XId

$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10e



Formula XIe














$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10f



Formula XIc


R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10g



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10h



Formula XIh














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10i




Formula XII

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl



Table 10j



Formula XIj














$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10k



Formula XIk

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10(I)



Formula XI(1)

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10m



Formula XIIm

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10n

















$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10o



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 10p



Formula XIp

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 10q



Formula XIq














$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl




Table 11



Formula XII


$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11a




R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11b




R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11c



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 11d



Formula XIIId

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 11e



Formula XIIe

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 11f



Formula XIIIf

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl

Table 11g




Formula XIIg

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl




Table 11h




Formula XIIIh

$R^{11c1}$	$R^{11c2}$	$R^{11c1}$	$R^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl




R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl


Table 11k



Formula XIIk

R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH2Ph	H	-NH2	H
OCH2CH2OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO2Me	H	SO2Me	Cl

Table 11(l)



Formula XII(l)















$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$	$\text{R}^{11\text{c}1}$	$\text{R}^{11\text{c}2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11m



R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11n
















$\text{R}^{11c1}$	$\text{R}^{11c2}$	$\text{R}^{11c1}$	$\text{R}^{11c2}$
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11o



Formula XIIo














R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
			
			
			
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl




Table 11p




R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Table 11q




R <sup>11c1</sup>	R <sup>11c2</sup>	R <sup>11c1</sup>	R <sup>11c2</sup>
H	H	OMe	H
Me	Me	OMe	Me
Br	Me	F	H
H	Cl	Et	H
Me	Cl	-OH	Me
Et	Cl	Et	Br
OCH <sub>2</sub> Ph	H	-NH <sub>2</sub>	H
OCH <sub>2</sub> CH <sub>2</sub> OMe	H	H	Et
Me	F	Et	F
Me	H	Me	Et
SO <sub>2</sub> Me	H	SO <sub>2</sub> Me	Cl

Also preferred are compounds according to Tables 1 through Table 11q, wherein



the biphenylene portions of their formulae:

are each replaced with the following ring structure:




wherein each of the A and D ring portions are as follows:

A	D	A	D

Even more preferred compounds are set forth in Tables 12-24, below.

**Table 12**



**Table 13**




Table 14




Table 15




Table 16








Table 17



wherein A—Q— is selected from the group consisting of:

t-Bu ; O-t-Bu ;  $-(CH_2)_{0-5}$ -amino ; OH ;



carboxylic acid ester ; carboxamide ;








Table 18





**Table 19**


**Table 20**

Table 21




Table 22





Table 23



and

Table 24




Table 25




Table 26






Table 27




Wherein:

Q is a direct link, and A is a member selected from the group:




or Q is a  $-\text{C}(=\text{NH})-$  group, and A is a member selected from the group:




G is a direct link;

J is a member selected from the group:



Y-L is a member selected from the group:



This invention also encompasses all pharmaceutically acceptable isomers, salts, hydrates and solvates of the compounds of formula I. In addition, the compounds of formula I can exist in various isomeric and tautomeric forms, and all such forms are meant to be included in the invention, along with pharmaceutically acceptable salts, hydrates and



solvates of such isomers and tautomers.

The compounds of this invention may be isolated as the free acid or base or converted to salts of various inorganic and organic acids and bases. Such salts are within the scope of this invention. Non-toxic and physiologically compatible salts are particularly useful although other less desirable salts may have use in the processes of isolation and purification.

A number of methods are useful for the preparation of the salts described above and are known to those skilled in the art. For example, the free acid or free base form of a compound of one of the formulas above can be reacted with one or more molar equivalents of the desired acid or base in a solvent or solvent mixture in which the salt is insoluble, or in a solvent like water after which the solvent is removed by evaporation, distillation or freeze drying. Alternatively, the free acid or base form of the product may be passed over an ion exchange resin to form the desired salt or one salt form of the product may be converted to another using the same general process.

#### Prodrug Derivatives of Compounds

This invention also encompasses prodrug derivatives of the compounds contained herein. The term "prodrug" refers to a pharmacologically inactive derivative of a parent drug molecule that requires biotransformation, either spontaneous or enzymatic, within the organism to release the active drug. Prodrugs are variations or derivatives of the compounds of this invention which have groups cleavable under metabolic conditions. Prodrugs become the compounds of the invention which are pharmaceutically active *in vivo*, when they undergo solvolysis under physiological conditions or undergo enzymatic degradation. Prodrug compounds of this invention may be called single, double, triple etc., depending on the number of biotransformation steps required to release the active drug within the organism, and indicating the number of functionalities present in a precursor-type form. Prodrug forms often offer advantages of solubility, tissue compatibility, or delayed release in the mammalian organism (see, Bundgard, Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985 and Silverman, The Organic Chemistry of Drug Design and Drug Action, pp. 352-401, Academic Press, San Diego, CA, 1992). Prodrugs commonly known in the art include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acids

with a suitable alcohol, or amides prepared by reaction of the parent acid compound with an amine, or basic groups reacted to form an acylated base derivative. Moreover, the prodrug derivatives of this invention may be combined with other features herein taught to enhance bioavailability.

As mentioned above, the compounds of this invention find utility as therapeutic agents for disease states in mammals which have disorders of coagulation such as in the treatment or prevention of unstable angina, refractory angina, myocardial infarction, transient ischemic attacks, thrombotic stroke, embolic stroke, disseminated intravascular coagulation including the treatment of septic shock, deep venous thrombosis in the prevention of pulmonary embolism or the treatment of reocclusion or restenosis of reperfused coronary arteries. Further, these compounds are useful for the treatment or prophylaxis of those diseases which involve the production and/or action of factor Xa/prothrombinase complex. This includes a number of thrombotic and prothrombotic states in which the coagulation cascade is activated which include but are not limited to, deep venous thrombosis, pulmonary embolism, myocardial infarction, stroke, thromboembolic complications of surgery and peripheral arterial occlusion.

Accordingly, a method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprises administering to the mammal a therapeutically effective amount of a compound of this invention. In addition to the disease states noted above, other diseases treatable or preventable by the administration of compounds of this invention include, without limitation, occlusive coronary thrombus formation resulting from either thrombolytic therapy or percutaneous transluminal coronary angioplasty, thrombus formation in the venous vasculature, disseminated intravascular coagulopathy, a condition wherein there is rapid consumption of coagulation factors and systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the microvasculature leading to widespread organ failure, hemorrhagic stroke, renal dialysis, blood oxygenation, and cardiac catheterization.

The compounds of the invention also find utility in a method for inhibiting the coagulation of biological samples, (*e.g.* blood) which comprises the administration of a compound of the invention.

The compounds of the present invention may also be used in combination with

other therapeutic or diagnostic agents. In certain preferred embodiments, the compounds of this invention may be coadministered along with other compounds typically prescribed for these conditions according to generally accepted medical practice such as anticoagulant agents, thrombolytic agents, or other antithrombotics, including platelet aggregation inhibitors, tissue plasminogen activators, urokinase, prourokinase, streptokinase, heparin, aspirin, or warfarin. The compounds of the present invention may act in a synergistic fashion to prevent reocclusion following a successful thrombolytic therapy and/or reduce the time to reperfusion. These compounds may also allow for reduced doses of the thrombolytic agents to be used and therefore minimize potential hemorrhagic side-effects. The compounds of this invention can be utilized *in vivo*, ordinarily in mammals such as primates, (e.g. humans), sheep, horses, cattle, pigs, dogs, cats, rats and mice, or *in vitro*.

The biological properties of the compounds of the present invention can be readily characterized by methods that are well known in the art, for example by the *in vitro* protease activity assays and *in vivo* studies to evaluate antithrombotic efficacy, and effects on hemostasis and hematological parameters, such as are illustrated in the examples.

Diagnostic applications of the compounds of this invention will typically utilize formulations in the form of solutions or suspensions. In the management of thrombotic disorders the compounds of this invention may be utilized in compositions such as tablets, capsules or elixirs for oral administration, suppositories, sterile solutions or suspensions or injectable administration, and the like, or incorporated into shaped articles. Subjects in need of treatment (typically mammalian) using the compounds of this invention can be administered dosages that will provide optimal efficacy. The dose and method of administration will vary from subject to subject and be dependent upon such factors as the type of mammal being treated, its sex, weight, diet, concurrent medication, overall clinical condition, the particular compounds employed, the specific use for which these compounds are employed, and other factors which those skilled in the medical arts will recognize.

Formulations of the compounds of this invention are prepared for storage or administration by mixing the compound having a desired degree of purity with physiologically acceptable carriers, excipients, stabilizers etc., and may be provided in sustained release or timed release formulations. Acceptable carriers or diluents for

therapeutic use are well known in the pharmaceutical field, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co., (A.R. Gennaro edit. 1985). Such materials are nontoxic to the recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, acetate and other organic acid salts, antioxidants such as ascorbic acid, low molecular weight (less than about ten residues) peptides such as polyarginine, proteins, such as serum albumin, gelatin, or immunoglobulins, hydrophilic polymers such as polyvinylpyrrolidinone, amino acids such as glycine, glutamic acid, aspartic acid, or arginine, monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose or dextrans, chelating agents such as EDTA, sugar alcohols such as mannitol or sorbitol, counterions such as sodium and/or nonionic surfactants such as Tween, Pluronics or polyethyleneglycol.

Dosage formulations of the compounds of this invention to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile membranes such as 0.2 micron membranes, or by other conventional methods. Formulations typically will be stored in lyophilized form or as an aqueous solution. The pH of the preparations of this invention typically will be 3-11, more preferably 5-9 and most preferably 7-8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of cyclic polypeptide salts. While the preferred route of administration is by injection, other methods of administration are also anticipated such as orally, intravenously (bolus and/or infusion), subcutaneously, intramuscularly, colonically, rectally, nasally, transdermally or intraperitoneally, employing a variety of dosage forms such as suppositories, implanted pellets or small cylinders, aerosols, oral dosage formulations and topical formulations such as ointments, drops and dermal patches. The compounds of this invention are desirably incorporated into shaped articles such as implants which may employ inert materials such as biodegradable polymers or synthetic silicones, for example, Silastic, silicone rubber or other polymers commercially available.

The compounds of the invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of lipids, such as cholesterol, stearylamine or phosphatidylcholines.



The compounds of this invention may also be delivered by the use of antibodies, antibody fragments, growth factors, hormones, or other targeting moieties, to which the compound molecules are coupled. The compounds of this invention may also be coupled with suitable polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidinone, pyran copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxyethyl-aspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, compounds of the invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels. Polymers and semipermeable polymer matrices may be formed into shaped articles, such as valves, stents, tubing, prostheses and the like.

Therapeutic compound liquid formulations generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by hypodermic injection needle.

Therapeutically effective dosages may be determined by either *in vitro* or *in vivo* methods. For each particular compound of the present invention, individual determinations may be made to determine the optimal dosage required. The range of therapeutically effective dosages will be influenced by the route of administration, the therapeutic objectives and the condition of the patient. For injection by hypodermic needle, it may be assumed the dosage is delivered into the body's fluids. For other routes of administration, the absorption efficiency must be individually determined for each compound by methods well known in pharmacology. Accordingly, it may be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. The determination of effective dosage levels, that is, the dosage levels necessary to achieve the desired result, will be readily determined by one skilled in the art. Typically, applications of compound are commenced at lower dosage levels, with dosage levels being increased until the desired effect is achieved.

The compounds of the invention can be administered orally or parenterally in an

effective amount within the dosage range of about 0.1 to 100 mg/kg, preferably about 0.5 to 50 mg/kg and more preferably about 1 to 20 mg/kg on a regimen in a single or 2 to 4 divided daily doses and/or continuous infusion.

Typically, about 5 to 500 mg of a compound or mixture of compounds of this invention, as the free acid or base form or as a pharmaceutically acceptable salt, is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, dye, flavor etc., as called for by accepted pharmaceutical practice. The amount of active ingredient in these compositions is such that a suitable dosage in the range indicated is obtained.

Typical adjuvants which may be incorporated into tablets, capsules and the like are binders such as acacia, corn starch or gelatin, and excipients such as microcrystalline cellulose, disintegrating agents like corn starch or alginic acid, lubricants such as magnesium stearate, sweetening agents such as sucrose or lactose, or flavoring agents. When a dosage form is a capsule, in addition to the above materials it may also contain liquid carriers such as water, saline, or a fatty oil. Other materials of various types may be used as coatings or as modifiers of the physical form of the dosage unit. Sterile compositions for injection can be formulated according to conventional pharmaceutical practice. For example, dissolution or suspension of the active compound in a vehicle such as an oil or a synthetic fatty vehicle like ethyl oleate, or into a liposome may be desired. Buffers, preservatives, antioxidants and the like can be incorporated according to accepted pharmaceutical practice.

#### Preparation of Compounds

The compounds of the present invention may be synthesized by either solid or liquid phase methods described and referenced in standard textbooks, or by a combination of both methods. These methods are well known in the art. See, Bodanszky, "The Principles of Peptide Synthesis", Hafner, *et al.*, Eds., Springer-Verlag, Berlin, 1984.

Starting materials used in any of these methods are commercially available from chemical vendors such as Aldrich, Sigma, Nova Biochemicals, Bachem Biosciences, and


the like, or may be readily synthesized by known procedures.

Reactions are carried out in standard laboratory glassware and reaction vessels under reaction conditions of standard temperature and pressure, except where otherwise indicated.

During the synthesis of these compounds, the functional groups of the amino acid derivatives used in these methods are protected by blocking groups to prevent cross reaction during the coupling procedure. Examples of suitable blocking groups and their use are described in "The Peptides: Analysis, Synthesis, Biology", Academic Press, Vol. 3 (Gross, *et al.*, Eds., 1981) and Vol. 9 (1987), the disclosures of which are incorporated herein by reference.


Non-limiting exemplary synthesis schemes are outlined directly below, and specific steps are described in the Examples. The reaction products are isolated and purified by conventional methods, typically by solvent extraction into a compatible solvent. The products may be further purified by column chromatography or other appropriate methods.

Scheme 1







Scheme 2




Scheme 3




Scheme 4




Scheme 5




Scheme 6




Scheme 7




Scheme 8



Scheme 9




Scheme 10







Scheme 11




Scheme 12




Scheme 13




Scheme 14



Scheme 15



Scheme 16



### Compositions and Formulations

The compounds of this invention may be isolated as the free acid or base or converted to salts of various inorganic and organic acids and bases. Such salts are within the scope of this invention. Non-toxic and physiologically compatible salts are particularly useful although other less desirable salts may have use in the processes of isolation and purification.

A number of methods are useful for the preparation of the salts described above and are known to those skilled in the art. For example, reaction of the free acid or free base form of a compound of the structures recited above with one or more molar equivalents of the desired acid or base in a solvent or solvent mixture in which the salt is insoluble, or in a solvent like water after which the solvent is removed by evaporation, distillation or freeze drying. Alternatively, the free acid or base form of the product may be passed over an ion exchange resin to form the desired salt or one salt form of the product may be converted to another using the same general process.

Diagnostic applications of the compounds of this invention will typically utilize formulations such as solution or suspension. In the management of thrombotic disorders the compounds of this invention may be utilized in compositions such as tablets, capsules or elixirs for oral administration, suppositories, sterile solutions or suspensions or injectable administration, and the like, or incorporated into shaped articles. Subjects in need of treatment (typically mammalian) using the compounds of this invention can be administered dosages that will provide optimal efficacy. The dose and method of administration will vary from subject to subject and be dependent upon such factors as the type of mammal being treated, its sex, weight, diet, concurrent medication, overall clinical condition, the particular compounds employed, the specific use for which these compounds are employed, and other factors which those skilled in the medical arts will recognize.

Formulations of the compounds of this invention are prepared for storage or administration by mixing the compound having a desired degree of purity with physiologically acceptable carriers, excipients, stabilizers etc., and may be provided in sustained release or timed release formulations. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical field, and are described, for example, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., (A.R. Gennaro edit. 1985). Such materials are nontoxic to the recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, acetate and other organic acid salts, antioxidants such as ascorbic acid, low molecular weight (less than about ten residues) peptides such as polyarginine, proteins, such as serum albumin, gelatin, or immunoglobulins, hydrophilic polymers such as polyvinylpyrrolidinone, amino acids such as glycine, glutamic acid, aspartic acid, or arginine, monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose or dextrans, chelating agents such as EDTA, sugar alcohols such as mannitol or sorbitol, counterions such as sodium and/or nonionic surfactants such as Tween, Pluronics or polyethyleneglycol.

Dosage formulations of the compounds of this invention to be used for therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile membranes such as 0.2 micron membranes, or by other conventional methods. Formulations typically will be stored in lyophilized form or as an aqueous solution. The pH of the preparations of this invention typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of cyclic polypeptide salts. While the preferred route of administration is by injection, other methods of administration are also anticipated such as intravenously (bolus and/or infusion), subcutaneously, intramuscularly, colonically, rectally, nasally or intraperitoneally, employing a variety of dosage forms such as suppositories, implanted pellets or small cylinders, aerosols, oral dosage formulations and topical formulations such



as ointments, drops and dermal patches. The compounds of this invention are desirably incorporated into shaped articles such as implants which may employ inert materials such as biodegradable polymers or synthetic silicones, for example, Silastic, silicone rubber or other polymers commercially available.

The compounds of this invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of lipids, such as cholesterol, stearylamine or phosphatidylcholines.

The compounds of this invention may also be delivered by the use of antibodies, antibody fragments, growth factors, hormones, or other targeting moieties, to which the compound molecules are coupled. The compounds of this invention may also be coupled with suitable polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxy-propyl-methacrylamide-phenol, polyhydroxyethyl-aspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the factor Xa inhibitors of this invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels. Polymers and semipermeable polymer matrices may be formed into shaped articles, such as valves, stents, tubing, prostheses and the like.

Therapeutic compound liquid formulations generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by hypodermic injection needle.

Therapeutically effective dosages may be determined by either *in vitro* or *in vivo* methods. For each particular compound of the present invention, individual determinations may be made to determine the optimal dosage required. The range of

therapeutically effective dosages will naturally be influenced by the route of administration, the therapeutic objectives, and the condition of the patient. For injection by hypodermic needle, it may be assumed the dosage is delivered into the body's fluids. For other routes of administration, the absorption efficiency must be individually determined for each inhibitor by methods well known in pharmacology. Accordingly, it may be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. The determination of effective dosage levels, that is, the dosage levels necessary to achieve the desired result, will be within the ambit of one skilled in the art. Typically, applications of compound are commenced at lower dosage levels, with dosage levels being increased until the desired effect is achieved.

A typical dosage might range from about 0.001 mg/kg to about 1000 mg/kg, preferably from about 0.01 mg/kg to about 100 mg/kg, and more preferably from about 0.10 mg/kg to about 20 mg/kg. Advantageously, the compounds of this invention may be administered several times daily, and other dosage regimens may also be useful.

Typically, about 0.5 to 500 mg of a compound or mixture of compounds of this invention, as the free acid or base form or as a pharmaceutically acceptable salt, is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, dye, flavor etc., as called for by accepted pharmaceutical practice. The amount of active ingredient in these compositions is such that a suitable dosage in the range indicated is obtained.

Typical adjuvants which may be incorporated into tablets, capsules and the like are a binder such as acacia, corn starch or gelatin, and excipient such as microcrystalline cellulose, a disintegrating agent like corn starch or alginic acid, a lubricant such as magnesium stearate, a sweetening agent such as sucrose or lactose, or a flavoring agent. When a dosage form is a capsule, in addition to the above materials it may also contain a liquid carrier such as water, saline, a fatty oil. Other materials of various types may be

used as coatings or as modifiers of the physical form of the dosage unit. Sterile compositions for injection can be formulated according to conventional pharmaceutical practice. For example, dissolution or suspension of the active compound in a vehicle such as an oil or a synthetic fatty vehicle like ethyl oleate, or into a liposome may be desired. Buffers, preservatives, antioxidants and the like can be incorporated according to accepted pharmaceutical practice.

In practicing the methods of this invention, the compounds of this invention may be used alone or in combination, or in combination with other therapeutic or diagnostic agents. In certain preferred embodiments, the compounds of this inventions may be coadministered along with other compounds typically prescribed for these conditions according to generally accepted medical practice, such as anticoagulant agents, thrombolytic agents, or other antithrombotics, including platelet aggregation inhibitors, tissue plasminogen activators, urokinase, prourokinase, streptokinase, heparin, aspirin, or warfarin. The compounds of this invention can be utilized in vivo, ordinarily in mammals such as primates, such as humans, sheep, horses, cattle, pigs, dogs, cats, rats and mice, or *in vitro*.

The preferred compounds of the present invention are characterized by their ability to inhibit thrombus formation with acceptable effects on classical measures of coagulation parameters, platelets and platelet function, and acceptable levels of bleeding complications associated with their use. Conditions characterized by undesired thrombosis would include those involving the arterial and venous vasculature.

With respect to the coronary arterial vasculature, abnormal thrombus formation characterizes the rupture of an established atherosclerotic plaque which is the major cause of acute myocardial infarction and unstable angina, as well as also characterizing the occlusive coronary thrombus formation resulting from either thrombolytic therapy or percutaneous transluminal coronary angioplasty (PTCA).

With respect to the venous vasculature, abnormal thrombus formation characterizes the condition observed in patients undergoing major surgery in the lower extremities or the abdominal area who often suffer from thrombus formation in the venous vasculature resulting in reduced blood flow to the affected extremity and a predisposition to pulmonary embolism. Abnormal thrombus formation further characterizes disseminated intravascular coagulopathy commonly occurs within both vascular systems during septic shock, certain viral infections and cancer, a condition wherein there is rapid consumption of coagulation factors and systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the microvasculature leading to widespread organ failure.


The compounds of this present invention, selected and used as disclosed herein, are believed to be useful for preventing or treating a condition characterized by undesired thrombosis, such as (a) the treatment or prevention of any thrombotically mediated acute coronary syndrome including myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, (b) the treatment or prevention of any thrombotically mediated cerebrovascular syndrome including embolic stroke, thrombotic stroke or transient ischemic attacks, (c) the treatment or prevention of any thrombotic syndrome occurring in the venous system including deep venous thrombosis or pulmonary embolus occurring either spontaneously or in the setting of malignancy, surgery or trauma, (d) the treatment or prevention of any coagulopathy including disseminated intravascular coagulation (including the setting of septic shock or other infection, surgery, pregnancy, trauma or malignancy and whether associated with multi-organ failure or not), thrombotic thrombocytopenic purpura, thromboangiitis obliterans, or thrombotic disease associated with heparin induced thrombocytopenia, (e) the treatment or prevention of thrombotic complications associated with extracorporeal circulation (e.g. renal dialysis, cardiopulmonary bypass or other oxygenation procedure, plasmapheresis), (f) the treatment or prevention of thrombotic complications associated with instrumentation (e.g.

cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve), and (g) those involved with the fitting of prosthetic devices.

Anticoagulant therapy is also useful to prevent coagulation of stored whole blood and to prevent coagulation in other biological samples for testing or storage. Thus the compounds of this invention can be added to or contacted with any medium containing or suspected to contain factor Xa and in which it is desired that blood coagulation be inhibited, e.g., when contacting the mammal's blood with material such as vascular grafts, stents, orthopedic prostheses, cardiac stents, valves and prostheses, extra corporeal circulation systems and the like.


Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. The following working examples therefore, specifically point out preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.



EXAMPLESExample 1

Part 1. Into a chilled suspension of 4-amino-3-nitrobenzoic acid (3.6 g, 20 mmol) in dry methanol (45 mL) was bubbled gaseous HCl for 10 min. The yellow suspension was stirred at room temperature for 2 days. The reaction mixture was filtered, washed with cold methanol and dried under high vacuum to give the HCl salt of methyl 4-amino-3-nitrobenzoate (3.25 g, 70%) as a yellow solid.

Part 2. To a chilled suspension of methyl 4-amino-3-nitrobenzoate (2.56 g, 11 mmol) and 4-dimethylaminopyridine (68 mg, 0.56 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (40 mL) and N,N-diisopropylethylamine (1.9 mL, 11 mmol) was added a solution of di-tert-butylidicarbonate (2.2 g, 10 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) dropwise over 20 min. The suspension was stirred at room temperature for 2 hours. Another solution of di-tert-butylidicarbonate (1.2 g, 5.5 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) was added, and the resulting solution was stirred for another hour. The reaction was then concentrated, diluted with ethyl acetate, washed with 5% citric acid, water and brine, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by silica gel chromatography using 5% ethyl acetate in CH<sub>2</sub>Cl<sub>2</sub> as eluent to yield methyl 4-[(tert-butoxy)carbonylamino]-3-nitrobenzoate (2.4 g, 74%). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.54 (s, 9H); 3.93 (s, 3H); 8.19-8.22 (d, 1H); 8.66-8.68 (d, 1H); 8.86 (s, 1H); 9.87 (s, 1H).

Example 2


Part 1. To a solution of methyl 4-[(tert-butoxy)carbonylamino]-3-nitrobenzoate (0.44 g, 1.5 mmol) in methanol (2 mL), ethyl acetate (3 mL) and triethylamine (0.21 mL, 1.5

mmol) was added 10% Pd on carbon (159 mg, 0.15 mmol). The reaction mixture was hydrogenated under 1 atm H<sub>2</sub> for 3.5 hours, filtered, and concentrated *in vacuo* to yield methyl 3-amino-4-[(tert-butoxycarbonyl)amino]benzoate (0.40 g, 99%) as a white solid. ES-MS (M+H-C<sub>4</sub>H<sub>8</sub>)<sup>+</sup> = 211.1.


Part 2. To a chilled solution of methyl 3-amino-4-[(tert-butoxy)carbonylamino]benzoate (0.40 g, 1.5 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and triethylamine (0.21 mL, 1.5 mmol) was added a solution of triphosgene (161 mg, 0.54 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (6 mL) over 5 min. The reaction was stirred at room temperature for 2 hr, diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed with water and brine, dried over sodium sulfate and concentrated *in vacuo*. The crude product was purified by silica gel chromatography using 50% ethyl acetate in CH<sub>2</sub>Cl<sub>2</sub> as eluent to give methyl 1-[(tert-butyl)oxycarbonyl]-2-oxo-(3H)-benzimidazole-5-carboxylate (0.27 g, 61%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.68 (s, 9H); 3.91 (s, 3H); 7.73 (s, 1H); 7.79-7.81 (d, 1H); 7.85-7.87 (dd, 1H); 8.9 (s, 1H). ES-MS (M+H-C<sub>4</sub>H<sub>8</sub>)<sup>+</sup> = 237.1.

### Example 3




Part 1. To a chilled solution of methyl 1-[(tert-butyl)oxycarbonyl]-2-oxo-(3H)-benzimidazole-5-carboxylate (0.37 g, 1.28 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (8 mL) and N,N-diisopropylethylamine (0.23 mL, 1.3 mmol) was added a solution of benzyl chloroformate (0.208 mL, 1.46 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL). The reaction was stirred at room temperature for 2.5 hr, diluted with ethyl acetate, washed with water and brine, dried over sodium sulfate and concentrated *in vacuo*. This residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (9 mL) and treated with neat trifluoroacetic acid (2 mL) at 0 °C for 40 min, diluted with ethyl acetate, washed with 5% NaHCO<sub>3</sub>, water and brine, dried over sodium sulfate and concentrated to yield methyl 2-oxo-3-[benzyloxycarbonyl]-(3H)-benzimidazole-5-carboxylate (0.37 g, 89%) as an off-white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 3.51 (s, 3H); 5.14 (s, 2H); 6.67-6.69 (d, 1H); 6.97-7.06 (m, 5H); 7.17-7.18 (d, 1H); 7.51-7.53 (d, 1H); 8.07 (s, 1H). ES-MS (M+Na)<sup>+</sup> = 349.0.

Example 4

To a chilled solution of methyl 2-oxo-3-(benzyloxycarbonyl)-(3H)-benzimidazole-5-carboxylate (110 mg, 0.337 mmol) in DMF (3 mL) was added cesium carbonate (0.275g, 0.84 mmol) followed by 7-(bromomethyl)naphthalene-2-carbonitrile (108 mg, 0.44 mmol). The reaction was stirred at room temperature for 1.5 hours, diluted with ethyl acetate, washed with water and brine, dried over sodium sulfate and concentrated *in vacuo*. This residue was redissolved in methanol (4 mL), CH<sub>3</sub>CN (3 mL) and 1N LiOH (0.7 mL). The reaction mixture was stirred for 1 hour, acidified with 1N HCl (0.7 mL), filtered and dried to yield methyl 1-[(7-cyano(2-naphthyl))methyl]-2-oxo-(3H)-benzimidazole-5-carboxylate (84 mg, 70%) as a white solid. ES-MS (M+H)<sup>+</sup>=358.1.

To a suspension of methyl 1-[(7-cyano-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxylate (82 mg, 0.23 mmol) in CH<sub>3</sub>CN (2.5 mL) and methanol (2 mL) was added 1N LiOH (2.3 mL, 10 eq.). The reaction mixture was stirred at room temperature for 3 days, acidified with 1N HCl (2.4 mL), filtered and dried to yield 1-[(7-cyano-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxylic acid (72 mg, 91%) as a white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 5.16 (s, 2H); 6.77-6.79 (d, 1H); 7.50-7.54 (t, 2H); 7.62-7.64 (dd, 1H); 7.68 (s, 1H); 7.72 (s, 1H); 7.78-7.84 (dd, 2H); 8.11 (s, 1H); 10.8 (s, 1H). ES-MS (M+H)<sup>+</sup>=344.1.


Example 5

A solution of 1-[(7-cyano-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxylic acid (70 mg, 0.20 mmol) and hydroxylamine hydrochloride (28 mg, 0.40 mmol) in dry




ethanol (4 mL) and DIEA (0.11 mL, 0.61 mmol) was stirred at 55 °C for 8 hours. The reaction was concentrated *in vacuo*, and the resulting residue redissolved in glacial acetic acid (4 mL). To this solution was added acetic anhydride (0.038 mL, 0.40 mmol). The reaction mixture was stirred at room temperature for 1 hr, followed by addition of methanol (3 mL) and 10% Pd on carbon (24 mg, 0.023 mmol). The mixture was hydrogenated under 1 atm H<sub>2</sub> for 17 hours, filtered, and concentrated *in vacuo*. Purification on a Vydac C<sub>18</sub> HPLC column yielded 1-[(7-carboxamidino-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxylic acid (30 mg, 41%) as a white fluffy solid after lyophilization. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 5.29 (s, 2H); 7.17-7.19 (d, 1H); 7.56 (s, 1H); etc. ES-MS (M+H)<sup>+</sup>=361.1.

#### Example 6




To a suspension of 1-[(7-carboxamidino-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxylic acid (7 mg, 0.019 mmol) and BOP reagent (10 mg, 0.023 mmol) in DMF (0.8 mL) and N,N-diisopropylethylamine (10 :L, 0.057 mmol) was added neat pyrrolidine (2 :L, 0.024 mmol). The resulting solution was stirred at room temperature under argon for 4 hours. The reaction was concentrated and purified by HPLC to yield 7-{[2-oxo-5-(pyrrolidinylcarbonyl)-(3H)-benzimidazolyl]methyl}naphthalene-2-carboxamidine (6 mg, 75%) as a white fluffy solid after lyophilization. ES-MS (M+H)<sup>+</sup>=414.1.

#### Example 7




Using a method similar method to that used in Example 6, 1-[(7-carboxamidino-2-naphthyl)methyl]-2-oxo-(3H)-benzimidazole-5-carboxamide was synthesized from the carboxylic acid via a BOP coupling with ammonium hydroxide. ES-MS (M+H)<sup>+</sup>=360.1.

#### Example 8




Using a method similar to that used in Example 6, 7-[[5-(N-methylcarbamoyl)-2-oxo-(3H)-benzimidazolyl]methyl]naphthalene-2-carboxamidine was synthesized from the carboxylic acid via a BOP coupling with methylamine. ES-MS (M+H)<sup>+</sup>=374.1.

#### Example 9



Using a method similar to that used in Example 6, 7-[[5-(N,N-dimethylcarbamoyl)-2-oxo-(3H)-benzimidazolyl]methyl]naphthalene-2-carboxamidine was synthesized from the carboxylic acid via a BOP coupling with dimethylamine. ES-MS (M+H)<sup>+</sup>=388.1.


#### Example 10



Using a method similar to that used in Example 6, 3-{3-[2-oxo-5-(pyrrolidinylcarbonyl)-(3H)-benzimidazolyl]propoxy}benzamidine was synthesized by alkylation with 3-(3-


bromopropoxy)benzenecarbonitrile, saponification, BOP coupling with pyrrolidine, and hydroxylamine method for conversion of nitrile to amidine. ES-MS  $(M+H)^+=408.1$ .

### Example 11




Using a method similar to that used in Example 6, 3-(2-methyl-5-([2-oxo-5-(pyrrolidinylcarbonyl)-(3H)-benzimidazolyl])methyl)-1,3-thiazol-4-yl)-benzamidinium was synthesized by alkylation with 3-[5-(chloromethyl)-2-methyl-1,3-thiazol-4-yl]benzenecarbonitrile, saponification, BOP coupling with pyrrolidine, and hydroxylamine method for conversion of nitrile to amidine. ES-MS  $(M+H)^+=461.0$ .

### Example 12



Using a method similar to that used in Example 6, 1-ethyl-2-([2-oxo-5-(pyrrolidinylcarbonyl)-(3H)-benzimidazolyl])methyl}indole-6-carboxamidinium was synthesized by alkylation with 2-(chloromethyl)-1-ethylindole-6-carbonitrile, saponification, BOP coupling with pyrrolidine, and hydroxylamine method for conversion of nitrile to amidine. ES-MS  $(M+H)^+=431.1$ .

### Example 13




Part 1. To a solution of 2*H*-1,4-Benzoxazin-3(4*H*)-one (1.86 g, 12.5 mmol, 1.0 equiv) in 25 mL of CHCl<sub>3</sub> at 0 °C was added Br<sub>2</sub> (2.0 g, 1.0 equiv) dropwise. After stirring at room temperature overnight, the solvent was evaporated and the residue was recrystallized from EtOH/H<sub>2</sub>O to give the product in 95% yield. LRMS found for C<sub>8</sub>H<sub>7</sub>BrNO<sub>2</sub> (M+H)<sup>+</sup>: 227.9.

Part 2. A solution of the product from Part 1 (103 mg, 0.4 mmol, 1.0 equiv), 2-(*t*-butylaminosulfonyl)phenylboronic acid (91.2 mg, 1.0 equiv), PdCl<sub>2</sub>(dppf) (32.6 mg, 0.1 equiv), and triethylamine (279  $\mu$ L, 5.0 equiv) in 10 mL of DME was degassed with argon for 15 min, then heated to reflux overnight. After cooling to room temperature, the mixture was diluted with ethyl acetate, washed with water, dried over MgSO<sub>4</sub>, and concentrated. Flash chromatography on silica gel gave the product in 21% yield. LRMS found for C<sub>18</sub>H<sub>21</sub>N<sub>2</sub>O<sub>4</sub>S (M+H)<sup>+</sup>: 361.1.


Part 3: A solution of the product from Part 2 (36 mg, 0.1 mmol, 1.0 equiv) in 2 mL of DMF was treated with 2-bromomethyl-7-cyanonaphthalene (40 mg, 75%, 1.2 equiv) and Cs<sub>2</sub>CO<sub>3</sub> (65 mg, 2 equiv) for 0.5h. The mixture was diluted with ethyl acetate, washed with water and subjected to flash column chromatography on silica gel to give the desired product in 95% yield. LRMS found for C<sub>30</sub>H<sub>28</sub>N<sub>3</sub>O<sub>4</sub>S (M+H)<sup>+</sup>: 526.2.

Step 4: The compound obtained in Step 3 (50 mg, 0.1 mmol, 1.0 equiv) was dissolved in 5 mL of methanol. The reaction mixture was cooled to 0 °C and HCl gas was bubbled in until saturation, and the mixture was stirred at room temperature overnight. The solvent was evaporated and the resulting residue was treated with ammonium acetate and 10 mL of methanol at reflux temperature for 2 hr. The solvent was removed at reduced pressure and the crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give the desired salt in 80% yield. LRMS found for C<sub>26</sub>H<sub>23</sub>N<sub>4</sub>O<sub>4</sub>S (M+H)<sup>+</sup>: 487.1.

Example 14

Step 1: The product from Example 13, Step 2 (36 mg, 0.1 mmol, 1.0 equiv) in 2 mL of DMF was treated with 2-bromomethyl-2'-cyanobiphenyl (41 mg, 1.5 equiv) and  $\text{Cs}_2\text{CO}_3$  (65 mg, 2 equiv) for 0.5 hr. The mixture was diluted with ethyl acetate, washed with water and purified by flash column chromatography to give the desired product in 91% yield. LRMS found for  $\text{C}_{32}\text{H}_{30}\text{N}_3\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 552.2.

Step 2: The product from Step 1 (50 mg, 0.09 mmol, 1.0 equiv) was dissolved in 5 mL of methanol. The reaction mixture was cooled to 0 °C, HCl gas was bubbled in until saturation, and the mixture was stirred at room temperature overnight. The solvent was evaporated and the resulting residue was treated with ammonium acetate and 10 ml methanol at reflux temperature for 2 hr. The solvent was removed under reduced pressure and the crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  to give the desired salt in 77% yield. LRMS found for  $\text{C}_{28}\text{H}_{25}\text{N}_4\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 513.2.

Example 15 and 16

Part 1. A solution of 2-nitro-5-bromophenol (2.53 g, 18.2 mmol) in 15 mL of ethanol was treated with  $\text{SnCl}_2 \cdot \text{H}_2\text{O}$ , and the reaction was heated at reflux for 3h. The solvent was evaporated to give a white residue, which was used in the next step without further purification. LRMS:  $\text{C}_9\text{H}_{10}\text{NO}_3$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 180.1.

Part 2. A solution of 2-amino-5-bromophenol (2.53 g, 18.2 mmol, 1.0 equiv) in 15 mL of isobutylmethyl ketone and 15 mL of water was cooled to 0 °C,  $\text{NaHCO}_3$  (3.67 g, 2.4 equiv) and then chloroacetyl chloride (2.36 g, 1.67 mL, 1.15 equiv) were added. The mixture was heated to reflux overnight, then cooled to room temperature. The mixture was diluted with ethyl acetate, washed with water, dried over  $\text{MgSO}_4$ , and concentrated *in vacuo*. Flash chromatography on silica gel gave 7-bromo-3,4-dihydro-2H-1,4-benzoxazin-3-one in 77% yield. LRMS found for  $\text{C}_9\text{H}_{10}\text{NO}_3$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 180.1.

Part 3. A solution of 7-bromo-3,4-dihydro-2H-1,4-benzoxazin-3-one (103 mg, 0.4 mmol, 1.0 equiv), 2-t-butylaminosulfonyl phenyl boronic acid (91.2 mg, 1.0 equiv),  $\text{PdCl}_2(\text{dppf})$  (32.6 mg, 0.1 equiv), and triethylamine (279  $\mu\text{L}$ , 5.0 equiv) in 10 mL of DME was degassed with argon for 15 min, then heated to reflux overnight. After cooling to room temperature, the mixture was diluted with ethyl acetate, washed with water, dried over  $\text{MgSO}_4$ , evaporated. Flash chromatography on silica gel gave the product in 21% yield. LRMS found for  $\text{C}_{18}\text{H}_{21}\text{N}_2\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 361.1.


Part 4. The product from Part 3 (36 mg, 0.1 mmol, 1.0 equiv) in 2 mL DMF was treated with 1-(3-cyanophenyl)-3-methyl-5-chloromethyl pyrazole (35 mg, 1.5 equiv) and  $\text{Cs}_2\text{CO}_3$  (65 mg, 2 equiv) for 0.5 hr. The mixture was diluted with ethyl acetate, washed with water and purified over silica gel to give the desired product in 91% yield. LRMS found for  $\text{C}_{30}\text{H}_{29}\text{N}_5\text{O}_4\text{S}$  ( $\text{M}+\text{H}$ )<sup>+</sup>: 556.1.

Part 5. The compound from Part 4 (50 mg, 0.09 mmol, 1.0 equiv) was dissolved in 5 mL of methanol. The reaction mixture was cooled to 0 °C,  $\text{HCl}$  gas was bubbled in until saturation, and the mixture was stirred at room temperature overnight. The solvent was evaporated and the resulting residue was treated with ammonium acetate and 10 mL of methanol at reflux temperature for 2 hr. The reaction was concentrated *in vacuo* and the crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in




H<sub>2</sub>O/CH<sub>3</sub>CN to give the desired amidine (Example 15) in 77% yield. LRMS found for C<sub>26</sub>H<sub>24</sub>N<sub>6</sub>O<sub>4</sub>S (M+H)<sup>+</sup>: 516.2. Example 16 was obtained as a byproduct in 15% yield. LRMS found for C<sub>26</sub>H<sub>23</sub>N<sub>5</sub>O<sub>5</sub>S (M+H)<sup>+</sup>: 517.1.

### Example 17




Part 1: The product (36 mg, 0.1 mmol, 1.0 equiv) obtained in Part 3 of Example 15 in 2 mL DMF was treated with 1-(3-cyanophenyl)-3-methyl-5-chloromethyl pyrazole (41 mg, 1.5 equiv) and Cs<sub>2</sub>CO<sub>3</sub> 65 mg, 2 equiv) for 0.5h. The mixture was diluted with ethyl acetate, washed with water and purified over silica gel to give the desired product in 91% yield. LRMS found for C<sub>32</sub>H<sub>30</sub>N<sub>3</sub>O<sub>4</sub>S (M+H)<sup>+</sup>: 552.2.

Part 2: The compound obtained in Part 1 (50 mg, 0.09 mmol, 1.0 equiv) was dissolved in 5 mL of methanol. The reaction mixture was cooled to 0 °C, HCl gas was bubbled in until saturation, and the mixture was stirred at room temperature overnight. The solvent was evaporated and the resulting residue was treated with ammonium acetate and 10 ml of methanol at reflux temperature for 2 hr. The solvent was removed at reduced pressure and the crude benzamidine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give the desired salt in 77% yield. LRMS found for C<sub>28</sub>H<sub>25</sub>N<sub>4</sub>O<sub>4</sub>S (M+H)<sup>+</sup>: 513.2.


Example 18

To a solution of tert-butylamine (41.4g, 566 mmol) and triethylamine (118 mL, 849 mmol) in DCM (1000 mL) in an ice bath, was added benzenesulfonyl chloride (100 g, 566 mmol) dropwise, and the mixture was stirred at room temperature overnight. Water was added to the mixture and organic layer was washed with water, sat. NaCl, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and filtrate evaporated *in vacuo* to give the title compound as light yellow solid (117.63 g, 97.6%). ES-MS (M+H)<sup>+</sup> = 214.5.


Example 19

A solution of the compound from example 18 (53.25 g, 250 mmol) in THF (600 mL) was cooled with an ice water bath, and a 2.5 M solution of n-butyllithium in hexane (200 mL, 500 mmol) was added dropwise. A thick precipitate was formed when the reaction mixture was warmed up to 10 °C. Triisopropylborate was added, keeping the temperature below 35 °C. After 1 hr., the mixture was cooled in an ice bath, 1N HCl (405 mL) was added, and the mixture was stirred overnight. The mixture was extracted with ether (3 x 100 mL), and the combined organic extracts were extracted with 1N NaOH (3 x 130 mL). The aqueous extracts were acidified to pH 1 with 12 N HCl, and then extracted with ether (3 x 140 mL). The combined ether extracts were dried over MgSO<sub>4</sub>, and solvents were evaporated *in vacuo*. Hexane and ether were added and a white precipitate formed. The solid was collected and washed with 10% ether/hexane to give the title compound. ES-MS (M+H)<sup>+</sup> = 257.8.




Example 20


To a solution of 5-bromoindole (1.96 g, 10 mmol) in DME (40 mL) and H<sub>2</sub>O (10 mL), was added the compound from Example 19 (3.85 g, 15 mmol), NaHCO<sub>3</sub> (1.68g, 20 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.58g, 0.5 mmol). The mixture was heated to reflux overnight, then cooled to room temperature and diluted with ethyl acetate. The organic layer was washed with water, dried with MgSO<sub>4</sub>, filtered and concentrated. The residue was purified by silica gel column chromatography using 25% ethyl acetate in hexane as eluent to give the title compound (1.52g, 46%). ES-MS (M+Na)<sup>+</sup> 351.1.

Example 21

To a solution of the compound from Example 20 (328 mg, 1 mmol) in DMF (10 mL) was added 7-cyano-2-bromomethylnaphthalene (296 mg, 1.2 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (1.3g, 4 mmol), and the mixture was stirred at room temperature overnight. The reaction mixture was partitioned between water and ethyl acetate, and the organic layer was washed with water, 1N HCl, sat. NaCl, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The residue was subjected to flash column chromatography on silica gel using 50% ethyl acetate in CH<sub>2</sub>Cl<sub>2</sub> followed by 100% CH<sub>2</sub>Cl<sub>2</sub> as eluent to give the title compound (237 mg, 48%). ES-MS (M+H)<sup>+</sup> = 494.2.

Example 22

A solution of the compound from example 21 (115 mg, 0.23 mmol) in 20%MeOH/ethyl acetate (10 mL) was treated with a stream of HCl gas for 10 min. at 0 °C. The resulting solution was capped, stirred at room temperature overnight, then concentrated *in vacuo*. The residue was reconstituted in MeOH (10 mL) and the mixture was treated with ammonium acetate (350 mg, 4.6 mmol). The reaction mixture was heated at reflux for 2 hrs, then concentrated *in vacuo*. The residue was purified by prep HPLC to give the title compound as a white powder. ES-MS (M+H)<sup>+</sup> = 511.2.

Example 23

Part 1. A solution of 4-bromo-2-nitroaniline (2.97 g, 13.7 mmol) in 100 mL of dichloromethane was treated with N,N-diisopropylethylamine (2.38 mL, 13.7 mmol) and 4-dimethylaminopyridine (167 mg, 1.37 mmol). Di-tert-butylidicarbonate (3.88 g, 17.8 mmol) was then added in small portions. The mixture was stirred at room temperature overnight under argon, then diluted with 400 mL of dichloromethane and washed with water (x 2). The organic phase was dried over MgSO<sub>4</sub>, concentrated *in vacuo* and purified by flash column chromatography. The desired product was obtained as a yellow solid in 48% yield (2.07 g).

Part 2. To a mixture of the product from Part 1 (1.02 g, 3.22 mmol), the boronic acid from Example 19 (827 mg, 3.22 mmol), and tetrakis(triphenylphosphine)palladium(0) (186 mg, 0.161 mmol) in 60 mL of benzene was added a solution of sodium hydroxide (515 mg, 12.9 mmol) in 5 mL of water. The mixture was degassed using an argon stream for 15 minutes and then heated at reflux for 24 hours. The solution was concentrated *in vacuo*, the residue was dissolved in ethyl acetate, passed through a bed of celite, and washed with water (x 2). The organic phase was dried over  $\text{MgSO}_4$ , concentrated *in vacuo*, then purified by flash column chromatography. The desired product was obtained as a solid in 60% yield (0.87 g).

Part 3. The product from Part 2 (72 mg, 0.16 mmol) and the bromide from Example 24 (80mg, 0.24 mmol) were dissolved in 4 mL of dry DMF, and cesium carbonate (156 mg, 0.48 mmol) was added. The resulting mixture was stirred overnight, then diluted with 100 mL of ethyl ether, washed with water (x 2), and dried over  $\text{MgSO}_4$ . Filtration and concentration *in vacuo* gave a residue, which was subjected to flash column chromatography. The desired product was obtained in 43% yield (48 mg).


Part 4. The product from Part 3 (90 mg, 0.13 mmol) was dissolved in 5 mL of ethyl alcohol, and tin(II) chloride dihydrate (116 mg, 0.52 mmol) was added. The mixture was heated to reflux for 3 hours, and the ethanol was removed *in vacuo*. The residue was dissolved in ethyl acetate, washed with 1N aqueous NaOH and water. The organic phase was dried over  $\text{MgSO}_4$ , concentrated *in vacuo* and pumped to dryness to give the desired product in quantitative yield (84 mg).

Part 5. The crude product from Part 4 (84 mg) was dissolved in 7 mL glacial acetic acid, and the mixture was stirred at 80 °C overnight. The mixture was concentrated with toluene to remove traces of acetic acid, and the residue was purified by preparative HPLC.

Part 6. The product from Part 5 (20 mg, 0.034 mmol) was dissolved in 5 mL anhydrous methanol, and a vacuum-distilling adapter, equipped with a rubber septum and with a balloon on its side-arm, was placed on the reaction flask. The solution was chilled in ice, and HCl gas was bubbled through the solution via a long needle immersed in the solution

until the balloon expanded, and this reaction mixture was stirred overnight. The reaction was concentrated to dryness *in vacuo*, then dissolved in 5 mL of anhydrous methanol, and dry ammonium acetate (21 mg, 0.27 mmol) was added. The mixture was heated at reflux for 2 hours, then purified by preparative HPLC to afford the title compound. LRMS  $(M+H)^+ = 553$ .

#### Example 24




Part 1. A mixture of 1,1,1-trifluoro-2,4-pentanedione (5.43 mL, 44.7 mmol) and 3-bromophenylhydrazine HCl (10.0 g, 44.7 mmol) in 100 mL of ethanol was heated to reflux overnight. The reaction mixture was concentrated *in vacuo*, and the residue was dissolved in ethyl ether and washed with water (x 2). The organic phase was dried over  $MgSO_4$ , concentrated and subjected to flash column chromatography, giving 12.83 g (94%) of the desired product as an oil, which was a mixture of the two possible regioisomers.

Part 2. The mixture of isomers from part 1 (12.83 g, 41.9 mmol), KCN (5.45 g, 83.8 mmol), CuI (798 mg, 4.19 mmol) and tetrakis(triphenylphosphine)palladium(0) (2.42 g, 2.09 mmol) in 100 mL of acetonitrile was degassed using an argon stream for 30 minutes. The mixture was heated at reflux for 15 hours, then passed through a silica plug on a Buchner funnel using ethyl acetate as eluent. The filtrate was concentrated and dissolved in dichloromethane, washed with water (x 2), and dried over  $MgSO_4$ . Filtration and concentration gave a residue, which was subjected to column chromatography. The regioisomeric products were separable at this stage, giving 3.93 g (37%) of the 3-methyl and 4.0 g (38%) of the 5-methyl isomers.

Part 3. A mixture of the 5-methyl isomer (1.67 g, 6.63 mmol), NBS (1.41 g, 7.95 mmol), and AIBN (543 mg, 3.31 mmol) in 50 mL carbon tetrachloride was degassed using an argon stream for 15 minutes. The reaction was heated at reflux for 2 days, periodically

adding additional small amounts of AIBN. The resulting mixture was diluted with dichloromethane, washed with water (x 2), and dried over  $\text{MgSO}_4$ . Filtration and concentration gave a residue, which was subjected to flash column chromatography. The desired bromomethyl product was obtained in 1.06 g (48%), along with 1.06 g of recovered starting material.

#### Examples 25 and 26



Part 1. The nitro compound from Part 2 of Example 23 (2.70 g, 6.01 mmol) was dissolved in 60 mL of ethanol, and tin(II) chloride dihydrate (5.43 g, 24.0 mmol) was added. The mixture was heated at reflux for 1 hour, then concentrated on the rotovap. The residue was washed through a plug of silica gel and concentrated to give another residue, which was dissolved in 20 mL of formic acid (88%) and stirred at 55 °C for 1 hour. Traces of formic acid were removed by evaporation with toluene *in vacuo*, and the desired product was obtained by flash column chromatography to give 516 mg (26%) of the desired benzimidazole. LRMS  $(\text{M}+\text{H})^+ = 330$ .


Part 2. The benzimidazole from Part 1 (62 mg, 0.188 mmol) was dissolved in 5 mL of DMF, followed by the addition of 2-bromomethyl-7-cyanonaphthalene (93 mg, 0.376 mmol) and cesium carbonate (184 mg, 0.564 mmol). The mixture was stirred overnight, then diluted with ethyl acetate and washed with water (x 2). The organic phase was dried over  $\text{MgSO}_4$ , concentrated *in vacuo*, and the desired products were obtained as a mixture of two regiosomers by flash column chromatography.

Part 3. The mixture of isomers from Part 2 (90 mg, 0.18 mmol) was dissolved in 10 mL of dry methanol. A vacuum-distilling adapter, equipped with a rubber septum and with a balloon on its side-arm, was placed on the reaction flask. The solution was chilled in ice,



and HCl gas was introduced via a long needle immersed in the solution until the balloon began to inflate. The mixture was stirred overnight and then concentrated to dryness *in vacuo*. The residue was dissolved in 5 mL anhydrous methanol, and dry ammonium acetate (139 mg, 1.8 mmol) was added. The mixture was heated at reflux for 2 hours, then purified by preparative HPLC to afford the two title compounds. LRMS (M+H)<sup>+</sup> = 457.

### Example 27




Part 1. To a solution of 5-bromoindoline (1.0g, 5mmol) in 10ml of dioxane was added 5 mL of a 1N NaOH solution and 5 mL of water. This mixture was cooled with an ice bath, and di-tert-butyl dicarbonate (1.2g, 5.5mmol) was added in one portion. The reaction was allowed to warm to room temperature, stirred for 4 hours, and then concentrated. The residue was extracted with ethyl acetate (2 x 25 mL), and the combined organic phases were washed with water (2 x 25mL), saturated aqueous NaCl (2 x 25ml), then dried over MgSO<sub>4</sub>. Filtration and concentration *in vacuo* gave N-Boc-5-bromoindoline (1.33g, 89%) as a light brown powder after drying. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.541 (s, 9H); 3.03-3.08 (t, 2H); 3.92-3.96 (t, 2H); 7.23-7.27 (m, 3H).

Part 2. To a solution of N-Boc-5-bromoindoline (104 mg, 0.35 mmol) in 5ml of anhydrous dioxane were added the boronic acid from Example 19 (100 mg, 0.39 mmol), cesium carbonate (228mg, 0.7mmol) and tris(dibenzylideneacetone)dipalladium(0)-chloroform adduct (10 mg, 0.01 mmol). The reaction flask was thoroughly flushed with argon, and tri-tert-butylphosphine (0.006 mL, 0.025 mmol) was added via syringe. The reaction was heated to 80 °C and stirred overnight. The reaction was then cooled to room temperature, diluted with 25 mL of diethyl ether, flushed through a pad of Celite and concentrated *in vacuo* to give a brown residue. This residue was subjected to flash column chromatography on silica gel using 25% ethyl acetate in hexane to give the desired biphenylamine (150 mg, 100%) as a light brown oil after drying. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.02 (s, 9H); 1.22 (s, 9H); 3.10-3.15 (t, 2H); 4.02, (t, 2H); 7.25-8.14 (m, 7H).

Part 3. The product from Part 2 was treated with 10ml of 20% trifluoroacetic acid in dichloromethane for 10 minutes. The reaction mixture was concentrated *in vacuo*, redissolved in 25 mL of dichloromethane, washed with 1N NaOH (2x25ml), dried over magnesium sulfate, filtered and concentrated *in vacuo* to give the desired biphenylamine as a brown oil after drying. ES-MS ( $M+H^+$ ): 330.5

#### Example 28



Part 1. 3-Cyano-4-fluoronitrobenzene (9.50 g, 57 mmol) was dissolved in 250 mL of ethanol, 5% Pd/C (2.0 g) was added, and the black slurry was stirred under a hydrogen balloon for 2 days. The reaction mixture was filtered through a celite bed and concentrated *in vacuo* to give the amino compound in 95% yield (7.37 g). LRMS ( $M+H^+$ )  $m/z$  137.  $R_f$  0.62 (2:1 ethyl acetate/hexane).

Part 2. The amino compound from Part 1 (4.00 g, 29 mmol) was slurried in 28 mL of conc. HCl and chilled in an ice bath. Sodium nitrite (2.00 g, 29 mmol) was dissolved in 10 mL of water and chilled in ice bath, and this solution was added dropwise to the cold slurry of the amino compound. The reaction mixture was then stirred at 0 °C for 30 min. Tin chloride dihydrate (19.7 g, 29 mmol) was dissolved in 10 mL of conc. HCl and chilled in ice bath, and this solution was added dropwise to the previous reaction mixture. After completion, the mixture was chilled and the solid hydrazine product was isolated by filtration through a Buchner funnel. This material was washed with cold brine (60 mL) and cold hexane (60 mL) to give 5.85 g of a yellow solid, which was used in the next step without further purification.

Part 3. Ethyl 2,4-dioxovalerate (20.0 g, 127 mmol) was dissolved in 100 mL of ethanol, and methoxylamine hydrochloride (11.1 g, 133 mmol) was added. The mixture was heated at reflux for 20 hours, followed by concentration of the reaction mixture *in vacuo*. The residue was dissolved in ethyl acetate, and washed with water and brine. The separated organic phase was dried over  $\text{MgSO}_4$  and evaporated *in vacuo* to give 19.3 g (81%) of the desired imine as an oil. LRMS  $(\text{M}+\text{H})^+$   $m/z$  188.

Part 4. A solution of the product from Part 2 (5.85 g) and the product from Part 3 (3.80 g, 20.3 mmol) in acetic acid (100 mL) and THF (50 mL) was heated at reflux overnight, and the reaction mixture was concentrated *in vacuo*. The residue was taken up in ethyl acetate and washed with water and brine. The organic phase was dried, filtered, and concentrated and the residue was subjected to flash column chromatography to yield the desired 5-carboethoxy isomer (4.45 g, 56% for two steps) and the undesired 3-carboethoxy isomer (1.00 g, 12% over two steps). LRMS  $(\text{M}+\text{H})^+$   $m/z$  274. 5-Carbomethoxy isomer  $R_f$  0.66 (1:1 ethyl acetate/hexane). 3-Carbomethoxy isomer  $R_f$  0.50 (1:1 ethyl acetate/hexane).


Part 5. The biphenylamine from Example 27, Part 3 (90 mg, 0.27 mmol) was dissolved in 2 mL of dichloromethane, and trimethylaluminum (0.7 mL of a 2.0 M solution in hexane, 1.35 mmol) was added at room temperature. The mixture was stirred for 30 minutes, and then a solution of the 5-carbomethoxy ester from Part 4 (77 mg, 0.28 mmol) in 3 mL of dichloromethane was added dropwise to the reaction. The resulting mixture was stirred under argon overnight. The reaction was then quenched by the careful addition of aqueous sat. Rochelle's salt, and the reaction was extracted with dichloromethane (x 3). The organic layer was concentrated and the residue was subjected to flash column chromatography to afford the desired coupling product (90 mg, 60%). LRMS  $(\text{M}+\text{H})^+$   $m/z$  558.  $R_f$  0.29 (1:1 ethyl acetate/hexane).

Part 6. The product from part 5 (83 mg, 0.15 mmol) was dissolved in 2 mL of ethanol, hydrazine monohydrate (40 :L, 0.75 mmol) was added dropwise, and the reaction mixture was heated at reflux overnight. Concentration *in vacuo* gave a residue, which was dissolved in 2 mL of trifluoroacetic acid and heated at reflux for 30 minutes. The resulting




mixture was concentrated and purified by prep HPLC [(C18 column, standard H<sub>2</sub>O to CH<sub>3</sub>CN gradient (0.1% TFA)] to give the desired compound. LRMS (M+H)<sup>+</sup> m/z 514.

#### Example 29



This compound was prepared from the 3-carbomethoxy intermediate from Part 4 of Example 28 in a manner similar to that in Parts 4 and 5, Example 28.

#### Example 30



Part 1. 3-Bromophenylhydrazine hydrochloride (6.80 g, 30.4 mmol) and the product from Example 28, Part 3 (2.84 g, 15.2 mmol) were dissolved in 60 mL of acetic acid and 30 mL of THF, and the reaction mixture was heated at reflux overnight. The solvent was removed *in vacuo*, the residue was dissolved in ethyl acetate, and the solution was washed with water and sat. NaCl. The organic layer was dried over MgSO<sub>4</sub>, evaporated *in vacuo*, and purified by flash column chromatography to give the desired pyrazole in quantitative yield. LRMS (M+H)<sup>+</sup> m/z 309, 311. R<sub>f</sub> 0.47 (1:4 ethyl acetate/hexane).


Part 2. The product from Part 1 was dissolved in 60 mL of acetonitrile, and KCN (2.91 g, 44.8 mmol), CuI (0.43 g, 2.2 mmol) and Pd(PPh<sub>3</sub>)<sub>4</sub> (1.3 g, 1.1 mmol) were added. The

mixture was degassed by bubbling argon through the solution for 30 minutes, and the reaction mixture was heated under argon at reflux overnight. The black mixture was then passed through a silica plug, and the filtrate was concentrated to give a residue, which was taken up in ethyl acetate and washed with water (x 2). The organic layer was dried, concentrated and subjected to flash chromatography to yield the desired nitrile as a white solid (2.69 g, 10.5 mmol, 69% for two steps). LRMS (M+H)<sup>+</sup> m/z 256. R<sub>f</sub> 0.68 (1:1 ethyl acetate/hexane).

Part 3. The product from Part 2 (2.00 g, 7.8 mmol) was dissolved in 40 mL of methanol, and water (20 mL) and LiOH monohydrate (0.66 g, 15.7 mmol) were added. The mixture was stirred for 1 hour, then acidified with 5N HCl to a pH of 1. The methanol was removed *in vacuo*, and the mixture was extracted with ethyl acetate (x 3). The combined organic phases were dried and evaporated to give the desired acid as a white solid (1.38 g, 78%). LRMS (M+H)<sup>+</sup> m/z 228, (M+Na)<sup>+</sup> m/z 250.

Part 4. The acid from part 3 (200 mg, 0.88 mmol), the biphenyl amine from Example 25, Part 3 (290 mg, 0.88 mmol), and 4,4-dimethylaminopyridine (10 mg) were dissolved in 4 mL of pyridine and chilled in ice bath. To the mixture was added POCl<sub>3</sub> (0.25 mL, 2.6 mmol), and the reaction was stirred in the cold for 1 hour. The reaction was quenched with ice water, diluted with ethyl acetate, and the organic phase was washed with sat. NaCl (x 2). The organic layer was dried, concentrated, and subjected to flash column chromatography to yield the desired amide (240 mg, 51%). LRMS (M+H)<sup>+</sup> m/z 540. R<sub>f</sub> 0.37 (1:1 ethyl acetate/hexane).

Part 5. The product from Part 4 (140 mg, 0.26 mmol) was dissolved in 10 mL of anhydrous DMF and chilled in ice bath. To the reaction mixture was added NaBH<sub>4</sub> (80 mg, 2.08 mmol) and anhydrous CoCl<sub>2</sub> beads (68 mg, 0.52 mmol). The mixture was stirred for 30 minutes in the cold, then quenched with ice water and diluted with ethyl acetate. The mixture was filtered through a pad of celite, and the filtrate was washed with sat. NaCl (x 2). The organic phase was dried, and concentrated *in vacuo* to give a residue, which was dissolved in 2 mL of trifluoroacetic acid and stirred at 60 °C for 30 minutes. The mixture was purified by prep HPLC [(C18 column, standard H<sub>2</sub>O to CH<sub>3</sub>CN gradient (0.1% TFA)] to afford the desired aminomethyl compound. LRMS (M+H)<sup>+</sup> m/z 488.

Examples 31 and 32


Part 1. A solution of the product from Example 27, Part 2 (80 mg, 0.2 mmol) in 4 mL of CH<sub>2</sub>Cl<sub>2</sub> was treated with 1 mL of trifluoroacetic acid, and the solution was stirred at room temperature for 10 min. The reaction mixture was then concentrated, and again concentrated from heptane to remove traces of trifluoroacetic acid. The residue was dissolved in 5 mL of DMF, and the bromomethyl compound from Example 24, Part 3 (80 mg, 0.24 mmol) was added, together with 195 mg of cesium carbonate, and the solution was stirred at room temperature for 1 hr. The reaction was diluted with ethyl acetate (25 mL), washed with water and sat. NaCl, and the organic layer was dried over MgSO<sub>4</sub>. Filtration and concentration gave a brown oil, which was used in the next reaction without further purification.

Part 2. The material from Part 1 (0.2 mmol) was taken up in 10 mL of absolute ethanol, and hydroxylamine hydrochloride (164 mg, 1.5 mmol) and triethylamine (0.2 mL, 1.5 mmol) were added. The solution was heated to 60 °C for 5 hours, then concentrated to give a residue, which was taken up in 5 mL of acetic acid, and acetic anhydride (0.2 mL) was added. The reaction mixture was stirred at room temperature for 30 min, then concentrated to dryness, followed by concentration with heptane to remove traces of acetic acid. The residue was used in the next reaction without further purification.

Part 3. The crude product from Part 2 was taken up in 5 mL of absolute ethanol, 1 drop of acetic acid was added, followed by 10% Pd/C (5 mg), and the reaction was placed under a balloon of hydrogen. After 4 hours, the starting material was gone by HPLC, so the reaction mixture was filtered and subjected directly to preparative HPLC [C18 column,


standard H<sub>2</sub>O to CH<sub>3</sub>CN (0.1% TFA) gradient]. Two products were isolated, the expected indoline product, and a product where the indoline ring had been oxidized to an indole ring. Indoline product: LRMS (M+H)<sup>+</sup> m/z 541.1. Indole product: LRMS (M+H)<sup>+</sup> m/z 539.1.

### Example 33




A solution of the product from Example 30, Part 4 (75 mg, 0.19 mmol) was treated with methanol and HCl, followed by ammonium acetate in a manner similar to that in Example 23, Part 6. Preparative HPLC gave the desired product.


### Example 34




Treatment of the product from Example 30, Part 4 with hydroxylamine hydrochloride and triethylamine gave the above amidoxime.

Example 35


A solution of 4-bromoaniline (15 g, 87 mmol) and  $K_2CO_3$  (181 g) in 170 mL of water was cooled to 5 °C. A solution of cinnamoyl chloride (18.2 g, 109 mmol) in acetone (87 mL) was added dropwise, resulting in the formation of a light brown solid. After 1 hour, the solid was recovered by filtration, washed with cold water, once with cold ether and dried overnight at 80 °C at 20 mm Hg to give the desired product (23.8 g, 90%). A portion of this amide (10 g, 33 mmol) was suspended in chlorobenzene (220 mL) and aluminum chloride (26 g, 195 mmol) was added portionwise. The solution was heated at reflux for 2 hr, the solution was concentrated *in vacuo* to one half of its original volume, and the solution was poured into 1L of ice water. The resulting solid was collected by filtration and recrystallized from methanol to give 6-bromoquinolin-2-one (5.0 g, 71%) as a brown solid.

Example 36

A solution of 6-bromohydroquinolin-2-one (1 g, 4.46 mmol), 2-bromomethyl-7-napthonitrile (1.16 g, 4.46 mmol), and cesium carbonate (1.5 g, 5.35 mmol) in 15 mL of DMF was stirred at room temperature for 18 h. The reaction mixture was diluted with water and ethyl acetate and filtered through celite. The organic layer was washed with water (x 2), then with sat. NaCl, dried over  $MgSO_4$ , and concentrated *in vacuo*. The residue was subjected to flash column chromatography on silica gel, using 100%  $CH_2Cl_2$  as eluent to give 7-[(6-bromo-2-oxohydroquinolyl)methyl]naphthalene-2-carbonitrile (876 mg, 50%) as a white solid. ES-MS  $(M+H)^+ = 389, 391$ .


Example 37

To a solution of 7-[(6-bromo-2-oxohydroquinolyl)methyl]naphthalene-2-carbonitrile (30 mg, 0.077 mmol), 2-[[[(tert-butyl)amino]sulfonyl]phenyl]boronic acid (20 mg, 0.077 mmol), Pd<sub>2</sub>(dba)<sub>3</sub> (1 mg, 1.5 mol%), and cesium carbonate (25 mg, 0.093 mmol) in dry dioxane (300 :L) was added tri-*t*-butylphosphine (0.7 uL, 3.6 mol%). The reaction was heated to 75 °C for 11 h, cooled to room temperature, and diluted with CH<sub>2</sub>Cl<sub>2</sub> (2 mL). The solution was then filtered through celite, washed with 1N HCl and sat. NaCl, dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to afford 7-[[6-(2-[[[(tert-butyl)amino]sulfonyl]phenyl)-2-oxohydroquinolyl]methyl}naphthalene-2-carbonitrile (34 mg, 85%). ES-MS (M+H)<sup>+</sup> = 522.2.


Example 38

A solution of the nitrile from Example 37 (34 mg, 0.64 mmol) in ethyl acetate (5mL) containing MeOH (150 :L) was cooled to -78 °C and HCl gas was bubbled in until saturation. The solution was stirred while warming to room temperature over 18 h. The solvent was removed *in vacuo*, the residue was taken up in methanol (2 mL), and dried. NH<sub>4</sub>OAc (50 mg, 0.64 mmol) was added. The mixture was heated to 80 °C for 1h, cooled, and purified by preparative HPLC [(C18 column, water/CH<sub>3</sub>CN gradient (0.1% TFA)]. The appropriate fractions were combined and lyophilized to give the desired amidine (5.6 mg, 19%) as a white powder. LRMS (M+H)<sup>+</sup> m/z 484.



Example 39

A solution of the nitrile from Example 37 (53 mg, 0.10 mmol) in methanol (0.5 mL) and DMF (0.25 mL) was treated with hydroxylamine hydrochloride (97 mg, 0.20 mmol), N,N-diisopropylethylamine (71  $\mu$ L, 0.20 mmol). The solution was heated to 40 °C for 18h, then concentrated *in vacuo*. The residue was stirred with TFA (2 mL) for 1h, then concentrated *in vacuo*, and this residue was purified by preparative HPLC [(C18 column, water/CH<sub>3</sub>CN gradient (0.1% TFA))]. The appropriate fractions were combined and lyophilized to give the desired amidoxime (17 mg, 32%) as a white powder. LRMS (M+H)<sup>+</sup> m/z 499.

Example 40

## Part 1. 7-Bromoisoquinoline

This compound was prepared as a 60:40 mixture with 5-bromoisoquinoline as in *J. Am. Chem. Soc.*, 1939, 61, 183.

## Part 2. 7-Bromoisoquinoline N-oxide hydrochloride

This compound was prepared by a procedure analogous to that for 6-bromoisoquinoline N-oxide hydrochloride as in PCT WO 98/47876. A solution of 7.8 g (37.5 mmol) of a 60:40 mixture of 7-bromo and 5-bromoisoquinoline in 125 mL of CH<sub>2</sub>Cl<sub>2</sub> was treated portionwise with 9.7 g (~39.4 mmol) of 3-chloroperoxybenzoic acid (~70% purity). The solution, which was initially homogeneous, deposited a voluminous precipitate over 1 hr.

Then 100 mL of methanol were added, and the reaction was concentrated to a volume of about 100 mL. Gaseous HCl was then bubbled through the solution for about 10 min, during which time the solution became warm and all of the precipitate dissolved. A few minutes later, another voluminous precipitate began to form. To this solution was added 100 mL of ether, and the mixture was stirred in an ice-water bath for 20 minutes. The resulting product was isolated by filtration, washed thoroughly with ether, and air-dried to give 8.07 g (83%) of the desired compound as a white solid, which was still a 60:40 mixture of the 7- and 5-bromo isomers.

### Part 3. 7-Bromo-1-chloroisoquinoline

This compound was prepared by a procedure analogous to that for 6-bromo-1-chloroisoquinoline as in PCT WO 98/47876. A solution of 8.07 g (31 mmol) of the mixture from Part B was taken up in 50 mL of POCl<sub>3</sub>, and the mixture was heated at 90 °C for 2 hr. The reaction mixture was concentrated to remove most of the POCl<sub>3</sub>, and the residue was taken up in 100 mL of CH<sub>2</sub>Cl<sub>2</sub>. The solution was carefully basified to pH 10 by the slow addition of 1N NaOH, and the organic layer was washed with 100 mL of H<sub>2</sub>O, 100 mL of sat. NaCl, and dried over MgSO<sub>4</sub>. Filtration and concentration gave a light yellow solid, which was subjected to flash column chromatography on silica gel first with 5% and then with 10% ethyl acetate in hexanes. A total of 3.62 g (48%) of the desired 7-bromo-1-chloroisoquinoline was isolated from this chromatography free of the 5-bromo isomer.



**Part 4. 7-Bromo-1-phenoxyisoquinoline**

A solution of 3.60 g (14.8 mmol) of 7-bromo-1-chloroisoquinoline and 1.5 g of solid KOH in 11.2 g of phenol was heated at 140 °C for 2 hr. The reaction was cooled to room temperature, then partitioned between 100 mL of CH<sub>2</sub>Cl<sub>2</sub> and 100 mL of 3N NaOH. The organic layer was washed with another 2 x 100 mL of 3N NaOH, then with 100 mL of H<sub>2</sub>O, and dried over MgSO<sub>4</sub>. Filtration and concentration gave a yellow oil, which was subjected to flash column chromatography on silica gel 30% CH<sub>2</sub>Cl<sub>2</sub> in hexanes, giving 3.42 g (77%) of the desired product as a light yellow solid.

**Part 5. 1-Amino-7-bromoisoquinoline**

A mixture of 3.40 g (11.3 mmol) of 1-amino-7-bromoisoquinoline and 7.65 g of ammonium acetate was heated at 150 °C for 15 hr. The reaction was cooled, and the residue was partitioned between 200 mL of ethyl acetate and 200 mL of 3N NaOH. The organic layer was extracted with 2 x 100 mL of 2N HCl, and the combined aqueous extracts were basified to pH 10 using 50% NaOH. This solution was extracted with 2 x 100 mL of ethyl acetate, and the organics were then washed with 100 mL of sat. NaCl and dried over MgSO<sub>4</sub>. Filtration and concentration gave 1.68 g (66%) of the desired amino compound as a yellow solid.

**Part 6. 1-[Bis(t-butoxycarbonyl)amino]-7-bromoisoquinoline**

A solution of 740 mg (3.32 mmol) of 1-amino-7-bromoisoquinoline in 50 mL of acetonitrile was treated with 1.4 mL of N,N-diisopropylethylamine and 100 mg of 4-(N,N-dimethylamino)pyridine, followed by 3.0 g (4.1 eq) of di-t-butylidicarbonate, and the reaction was stirred at 40 °C for 1 hr. By HPLC analysis, there was still some starting amino compound that remained, so another 1.0 g of di-t-butylidicarbonate were added, and the reaction was stirred at 40 °C for another 30 min. The reaction mixture was concentrated to give a dark oil, which was subjected to flash column chromatography on silica gel with 20% ethyl acetate in hexanes to give 736 mg of the desired product as a light yellow solid. Also isolated were 156 mg of product as a somewhat less pure light yellow solid, making the total yield 64%.

## Part 7. 1-[Bis(t-butoxycarbonyl)amino]isoquinoline-7-carboxaldehyde

A solution of 400 mg (0.95 mmol) of 1-[bis(t-butoxycarbonyl)amino]-7-bromoisquinoline in 50 mL of anhydrous THF was cooled with a liquid nitrogen/methanol slush bath (-98 °C), and 0.55 mL of a 2.43 M solution of n-BuLi in hexanes (1.3 eq) was added dropwise over 1 min. The solution was stirred in the cold for 5 min, then a solution of 5 mL of anhydrous DMF in 10 mL of anhydrous THF was added rapidly. The solution was allowed to warm to about 0 °C, then poured into 50 mL of 0.5 N HCl, and 50 mL of ethyl acetate were added. The aqueous layer was brought to pH 6 with 1N NaOH, 25 mL of sat. NaCl were added, and the layers were shaken and separated. The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated to give an oily residue. This residue was subjected to flash column chromatography on silica gel with 20% ethyl acetate in hexanes to give 190 mg (54%) of the desired aldehyde as a yellow semisolid.

## Part 8. (2Z)-3-{[1-bis(t-butoxycarbonyl)amino]isoquinolin-7-yl} acrylic acid, 2-(trimethylsilyl)ethyl ester


A solution of 117 mg (0.29 mmol) of [bis(2,2,2-trifluoroethoxy)phosphinyl]acetic acid, 2-(trimethylsilyl)ethyl ester (*J. Org. Chem.*, 1991, 56, 4204) and 400 mg of 18-crown-6 in 25 mL of anhydrous THF was cooled with a dry ice-acetone bath under argon, and 0.75 mL of a 0.5 M solution of potassium bis(trimethylsilyl)amide in toluene were added dropwise over 2 min. The reaction was stirred in the cold for 15 min, then a solution of 100 mg (0.27 mmol) of 1-[bis(t-butoxycarbonyl)amino]isoquinoline-7-carboxaldehyde in 25 mL of anhydrous THF was added dropwise over 10 min. The reaction was then allowed to warm to room temperature overnight, then partitioned between 100 mL of CH<sub>2</sub>Cl<sub>2</sub> and 50 mL of H<sub>2</sub>O. The organics were washed with aqueous NaCl, and dried over MgSO<sub>4</sub>. Filtration and concentration gave an oily residue, which was subjected to flash column chromatography on silica gel with 25% ethyl acetate in hexanes to give 33 mg of the desired product as a clear, colorless oil.

Part 9. (2Z)-N-[5-(2{[(N-1,1-dimethylethyl)amino]sulfonyl}phenyl)-1-indoliny]-3-{[1-bis(t-butoxycarbonyl)amino]isoquinolin-7-yl}acrylamide


A solution of 498 mg (0.97 mmol) of (2Z)-3-{[1-bis(t-butoxycarbonyl)amino]isoquinolin-7-yl}acrylic acid, 2-(trimethylsilyl)ethyl ester in 5 mL of DMF was treated at room temperature with 1.2 mL (1.25 eq) of a 1.0 M tetrabutylammonium fluoride in THF, and the reaction was stirred for 5 hr. The reaction mixture was diluted with 100 mL of ether, and the solution was washed with 100 mL of water. The aqueous layer was again extracted with 100 mL of ether, and the combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration and concentration gave 350 mg of an off-white solid, which was used without further purification. A solution of 50 mg of this carboxylic acid and 44 mg of 2-indolin-5-ylbenzenesulfonamide from Example 27, Part 3 in 3 mL of DMF was treated with 100  $\mu$ L of N,N-diisopropylethylamine and 60 mg of HATU, and the reaction was stirred at room temperature overnight. The reaction mixture was diluted with 100 mL of ethyl acetate and washed with sat. NaHCO<sub>3</sub> (2 x 25 mL), and the organic layer was dried over MgSO<sub>4</sub>. Filtration and concentration gave an orange oil, which was subjected to flash column chromatography on silica gel, using 20% ethyl acetate in hexanes as eluent to give 62 mg of the desired product as a yellow oil.

Part 10. (2Z)-N-[4-(2{aminosulfonyl}phenyl)-1-indoliny]-3-{aminoisoquinolin-7-yl}acrylamide


A solution of the yellow oil from Part 9 in 2 mL of trifluoroacetic acid was stirred at room temperature overnight. The reaction mixture was concentrated with CH<sub>2</sub>Cl<sub>2</sub> to remove most of the TFA, then purified directly by prep HPLC to give 15 mg of the desired product was obtained as an off-white solid. LRMS (M + H)<sup>+</sup> = 470.

Example 41

This compound was prepared by a method similar to that used in Example 23, Part 6.

Example 42

This compound was prepared by a method similar to that used in Example 23, Part 6.

Example 43

This compound was prepared by catalytic hydrogenation of the compound from Example 42.

### BIOLOGICAL ACTIVITY EXAMPLES

Evaluation of the compounds of this invention is guided by in vitro protease activity assays (see below) and in vivo studies to evaluate antithrombotic efficacy, and effects on hemostasis and hematological parameters.

The compounds of the present invention are dissolved in buffer to give solutions containing concentrations such that assay concentrations range from 0 to 100  $\mu\text{M}$ . In the assays for thrombin, prothrombinase and factor Xa, a synthetic chromogenic substrate is added to a solution containing test compound and the enzyme of interest and the residual catalytic activity of that enzyme is determined spectrophotometrically. The  $\text{IC}_{50}$  of a compound is determined from the substrate turnover. The  $\text{IC}_{50}$  is the concentration of test compound giving 50% inhibition of the substrate turnover. The compounds of the present invention desirably have an  $\text{IC}_{50}$  of less than 500 nM in the factor Xa assay, preferably less than 200 nM, and more preferred compounds have an  $\text{IC}_{50}$  of about 100 nM or less in the factor Xa assay. The compounds of the present invention desirably have an  $\text{IC}_{50}$  of less than 4.0  $\mu\text{M}$  in the prothrombinase assay, preferably less than 200 nM, and more preferred compounds have an  $\text{IC}_{50}$  of about 10 nM or less in the prothrombinase assay. The compounds of the present invention desirably have an  $\text{IC}_{50}$  of greater than 1.0  $\mu\text{M}$  in the thrombin assay, preferably greater than 10.0  $\mu\text{M}$ , and more preferred compounds have an  $\text{IC}_{50}$  of greater than 100.0  $\mu\text{M}$  in the thrombin assay.

#### Amidolytic Assays for determining protease inhibition activity

The factor Xa and thrombin assays are performed at room temperature, in 0.02 M Tris·HCl buffer, pH 7.5, containing 0.15 M NaCl. The rates of hydrolysis of the para-nitroanilide substrate S-2765 (Chromogenix) for factor Xa, and the substrate Chromozym TH (Boehringer Mannheim) for thrombin following preincubation of the enzyme with inhibitor for 5 minutes at room temperature, and were determined using the Softmax 96-well plate reader (Molecular Devices), monitored at 405 nm to measure the time dependent appearance of p-nitroaniline.



The prothrombinase inhibition assay is performed in a plasma free system with modifications to the method described by Sinha, U. *et al.*, Thromb. Res., 75, 427-436 (1994). Specifically, the activity of the prothrombinase complex is determined by measuring the time course of thrombin generation using the p-nitroanilide substrate Chromozym TH. The assay consists of preincubation ( 5 minutes) of selected compounds to be tested as inhibitors with the complex formed from factor Xa (0.5 nM), factor Va (2 nM), phosphatidyl serine:phosphatidyl choline (25:75, 20  $\mu$ M) in 20 mM Tris·HCl buffer, pH 7.5, containing 0.15 M NaCl, 5 mM CaCl<sub>2</sub> and 0.1% bovine serum albumin. Aliquots from the complex-inhibitor mixture are added to prothrombin (1 nM) and Chromozym TH (0.1 mM). The rate of substrate cleavage is monitored at 405 nm for two minutes. Eight different concentrations of inhibitor are assayed in duplicate. A standard curve of thrombin generation by an equivalent amount of untreated complex are used for determination of percent inhibition.

#### Antithrombotic Efficacy in a Rabbit Model of Venous Thrombosis

A rabbit deep vein thrombosis model as described by Hollenbach, S. et al., Thromb. Haemost. 71, 357-362 (1994), is used to determine the in-vivo antithrombotic activity of the test compounds. Rabbits are anesthetized with I.M. injections of Ketamine, Xylazine, and Acepromazine cocktail. A standardized protocol consists of insertion of a thrombogenic cotton thread and copper wire apparatus into the abdominal vena cava of the anesthetized rabbit. A non-occlusive thrombus is allowed to develop in the central venous circulation and inhibition of thrombus growth is used as a measure of the antithrombotic activity of the studied compounds. Test agents or control saline are administered through a marginal ear vein catheter. A femoral vein catheter is used for blood sampling prior to and during steady state infusion of test compound. Initiation of thrombus formation begins immediately after advancement of the cotton thread apparatus into the central venous circulation. Test compounds are administered from time = 30 min to time = 150 min at which the experiment is terminated. The rabbits are euthanized and the thrombus excised by surgical dissection and characterized by weight and

histology. Blood samples are analyzed for changes in hematological and coagulation parameters.

#### Effects of Compounds in Rabbit Venous Thrombosis model

Administration of compounds in the rabbit venous thrombosis model demonstrates antithrombotic efficacy at the higher doses evaluated. There are no significant effects of the compound on the aPTT and PT prolongation with the highest dose (100 µg/kg + 2.57 µg/kg/min). Compounds have no significant effects on hematological parameters as compared to saline controls. All measurements are an average of all samples after steady state administration of vehicle or (D)-Arg-Gly-Arg-thiazole. Values are expressed as mean  $\pm$  SD.


Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described.

It should be understood that the foregoing discussion and examples merely present a detailed description of certain preferred embodiments. It will be apparent to those of ordinary skill in the art that various modifications and equivalents can be made without departing from the spirit and scope of the invention. All the patents, journal articles and other documents discussed or cited above are herein incorporated by reference.

The claimed invention is:

1. A compound according to the formula:



wherein:

A is selected from:

- (a) C<sub>1</sub>-C<sub>6</sub>-alkyl;
- (b) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;
- (c) -N(-R<sup>2</sup>, -R<sup>3</sup>), R<sup>3</sup>-C(=N-R<sup>2</sup>)-, (-R<sup>2</sup>, -R<sup>3</sup>)N-C(=N-R<sup>2</sup>)-, (-R<sup>2</sup>, -R<sup>3</sup>)N-C(=N-R<sup>2</sup>)-N(-R-)-
- (d) phenyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;
- (e) naphthyl, which is independently substituted with 0-2 R<sup>1</sup> substituents; and
- (f) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1</sup> substituents;

R<sup>1</sup> is selected from:

Halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, R<sup>2</sup>-C(=N-R<sup>3</sup>)-, (-R<sup>2</sup>, -R<sup>3</sup>)N-C(=N-R<sup>2</sup>)-, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -C(=O)-N(-R<sup>2</sup>, -R<sup>3</sup>), -SO<sub>2</sub>N(-R<sup>2</sup>, -R<sup>3</sup>), -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1</sub>-C<sub>4</sub>-alkyl, -CN, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;



$R^2$  and  $R^3$  are independently selected from the group consisting of:

H,  $-OR^2$ ,  $-N(-R^2, -R^3)$ ,  $-C_{1-4}alkyl$ ,  $-C_{2-6}alkenyl$ ,  $-C_{2-6}alkynyl$ ,  $-C_{3-8}cycloalkyl$ ,  $-C_{0-4}alkylC_{3-8}cycloalkyl$ ,  $-C_{0-4}alkylphenyl$  and  $-C_{0-4}alkylnaphthyl$ , wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $-C_{1-4}alkyl$ ,  $-C_{2-6}alkenyl$ ,  $-C_{2-6}alkynyl$ ,  $-C_{3-8}cycloalkyl$ ,  $-C_{0-4}alkylC_{3-8}cycloalkyl$ ,  $-CN$ , and  $-NO_2$ ;

or  $R^2$  and  $R^3$  taken together can form a 3-8 membered cycloalkyl or a heterocyclic ring system, wherein the heterocyclic ring system may have from 5 to 10 ring atoms, with 1 to 2 rings being in the ring system and contain from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the heterocyclic ring system may be independently replaced with a member selected from the group consisting of halo,  $C_1-C_4-alkyl$ ,  $-CN$ ,  $-C_{1-4}alkyl$ ,  $-C_{2-6}alkenyl$ ,  $-C_{2-6}alkynyl$ ,  $-C_{3-8}cycloalkyl$ ,  $-C_{0-4}alkylC_{3-8}cycloalkyl$  and  $-NO_2$ ;

m is an integer of 0-2;

Q is a member selected from the group consisting of:

a direct link,  $-CH_2-$ ,  $-C(=O)-$ ,  $-N(R^4)-$ ,  $-N(R^4)CH_2-$ ,  $-C=N(R^4)-$ ,  $-C(=O)-N(R^4)-$ ,  $-N(R^4)-C(=O)-$ ,  $-SO_2-$ ,  $-O-$ ,  $-SO_2-N(R^4)-$  and  $-N(R^4)-SO_2-$ ;

$R^4$  is selected from:

H,  $-C_{1-4}alkyl$ ,  $-C_{2-6}alkenyl$ ,  $-C_{2-6}alkynyl$ ,  $-C_{3-8}cycloalkyl$ ,  $-C_{0-4}alkylC_{3-8}cycloalkyl$ ,  $-C_{0-4}alkylphenyl$  and  $-C_{0-4}alkylnaphthyl$ , wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $-C_{1-4}alkyl$ ,  $-C_{2-6}alkenyl$ ,  $-C_{2-6}alkynyl$ ,  $-C_{3-8}cycloalkyl$ ,  $-C_{0-4}alkylC_{3-8}cycloalkyl$ ,  $-CN$ , and  $-NO_2$ ;

D is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1a}$  substituents; and
- (b) an aromatic six-membered heterocyclic ring having from 1-2 ring nitrogen atoms, and wherein the ring atoms may be substituted with 0-2  $R^{1a}$  substituents;

$R^{1a}$  is selected from:

Halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2a</sup>R<sup>3a</sup>, SO<sub>2</sub>NR<sup>2a</sup>R<sup>3a</sup>, SO<sub>2</sub>R<sup>2a</sup>, CF<sub>3</sub>, OR<sup>2a</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>2a</sup> and R<sup>3a</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

M is a member selected from the group consisting of:

-N(R<sup>16</sup>)-C(=O)-, -N(R<sup>16</sup>)-C(=S)-, -C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-, -C(-R<sup>19</sup>, -R<sup>19a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-, -C(-R<sup>17</sup>)-C(-R<sup>18</sup>)-C(=O)-, -C(-R<sup>17</sup>)-C(-R<sup>18</sup>)-C(=S)-, -C(-R<sup>17</sup>)-C(-R<sup>18</sup>)-, -O-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -O-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S(=O)<sub>2</sub>-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S(=O)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -S-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S(=O)<sub>2</sub>-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -S(=O)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(=O)-C(=O)-, -N(R<sup>16</sup>)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=O)-, -N(R<sup>16</sup>)-C(-R<sup>17</sup>, -R<sup>18</sup>)-C(=S)-, -C(=S)-C(=S)-, -C(=S)-C(=O)-, -C(=O)-C(=S)-, -N=C(-R<sup>17</sup>)-C(=O)-, -N=C(-R<sup>17</sup>)-C(=S)-, -C(-R<sup>17</sup>)=N-, -N(-R<sup>16</sup>)-C(=O)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -O-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -N(-R<sup>16</sup>)-C(=S)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -S(=O)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -S(=O)<sub>2</sub>-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-, -C(=C(R<sup>17b</sup>, -R<sup>17c</sup>))-C(=O)-, -C(=C(R<sup>17b</sup>, -R<sup>17c</sup>))-C(=S)-, -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-R<sup>17</sup>, -R<sup>17a</sup>)-C(=S)-, -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-(N(-H, -R<sup>18b</sup>)), -R<sup>17a</sup>)-C(=O)-; -N=C(-R<sup>17</sup>)- and -N(-R<sup>16</sup>)-C(-R<sup>18</sup>, -R<sup>18a</sup>)-C(-(N(-H, -R<sup>18b</sup>)), -R<sup>17a</sup>)-C(=S)-; wherein the first named atom of the chain is directly attached to D, and wherein D, M and the N atom attached to the last chain atom of M collectively form a bicyclic ring structure;

R<sup>16</sup>, R<sup>17</sup>, R<sup>17a</sup>, R<sup>18</sup>, R<sup>18a</sup>, R<sup>18b</sup>, R<sup>19</sup>, and R<sup>19a</sup> are each independently selected from the group consisting of:

hydrogen, halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>R<sup>2</sup>, CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, -CN, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

R<sup>17b</sup> and R<sup>17c</sup> are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy, -C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the cycloalkyl, the phenyl ring, or the aromatic heterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, -CN, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

E is a member selected from the group consisting of:

a direct link, -C(=O)-, -C(=O)-N(R<sup>5</sup>)-, -C(-R<sup>5a</sup>, -R<sup>6a</sup>)- and -C(-R<sup>5b</sup>, -R<sup>6b</sup>)-C(-R<sup>5c</sup>, -R<sup>6c</sup>)-;

wherein R<sup>5</sup>, R<sup>5a</sup>, R<sup>6a</sup>, R<sup>5b</sup>, R<sup>6b</sup>, R<sup>5c</sup> and R<sup>6c</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylheteroaryl, C<sub>1-4</sub>alkylCOOH and C<sub>1-4</sub>alkylCOOC<sub>1-4</sub>alkyl, wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl,

C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -OH, -O-C<sub>1-4</sub>alkyl, -SH, -S-C<sub>1-4</sub>alkyl, -CN and -NO<sub>2</sub>;

G is selected from:

a direct link, -C(R<sup>7</sup>,R<sup>8</sup>)-, -C(R<sup>7a</sup>,R<sup>8a</sup>)-C(R<sup>7b</sup>,R<sup>8b</sup>)- and -C(R<sup>7c</sup>)=C(R<sup>8c</sup>)-;

wherein R<sup>7</sup>, R<sup>8</sup>, R<sup>7a</sup>, R<sup>8a</sup>, R<sup>7b</sup>, R<sup>8b</sup>, R<sup>7c</sup> and R<sup>8c</sup> are independently a member selected from from the group consisting of:

hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)OR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, a naturally occurring or synthetic amino acid side chain, and C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the C<sub>0-4</sub>alkylheterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>9</sup> and R<sup>10</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>3-8</sub>cycloalkyl, and C<sub>1-4</sub>alkyl-O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl-COOH wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>, and wherein R<sup>9</sup> and R<sup>10</sup> taken together can form a 5-8 membered heterocyclic ring;

J is a member selected from the group consisting of:

a direct link, -O-, -O-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -S-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)<sub>2</sub>-(-R<sup>11</sup>, -R<sup>11a</sup>)-, -C(=O)-, -C(=O)-N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(=O)-, -N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(-R<sup>11</sup>, -R<sup>11a</sup>)- and a monocyclic aromatic or

non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>11</sup> substituents;

R<sup>11</sup>, R<sup>11a</sup>, R<sup>11b</sup>, and R<sup>11</sup> are a member independently selected from the group consisting of:

hydrogen, halo, -CF<sub>3</sub>, -CN, -NR<sup>9</sup>R<sup>10</sup>, -SO<sub>2</sub>Me, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>2-6</sub>alkenyl, -O-C<sub>2-6</sub>alkynyl, -O-C<sub>3-8</sub>cycloalkyl, -O-C<sub>1-4</sub>alkyl-O-C<sub>1-4</sub>alkyl, -O-C<sub>1-4</sub>alkyl-COOH, -O-C<sub>1-4</sub>alkyl-phenyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>2-6</sub>alkenyl, -C(=O)-O-C<sub>2-6</sub>alkynyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, C<sub>0-4</sub>alkylC(=O)OR<sup>9</sup>, C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl; wherein from 1-4 hydrogen atoms on the C<sub>0-4</sub>alkylheterocyclic ring may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-phenyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OR<sup>9</sup>, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents;
- (b) naphthyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents; and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1b</sup> substituents;



$R^{1b}$  is a member selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN,  $-NO_2$ ,  $NR^{2b}R^{3b}$ ,  $SO_2NR^{2b}R^{3b}$ ,  $SO_2R^{2b}$ ,  $CF_3$ ,  $OR^{2b}$ ,  $O-CH_2-CH_2-OR^{2b}$ ,  $O-CH_2-COOR^{2b}$ ,  $N(R^{2b})-CH_2-CH_2-OR^{2b}$ ,  $N(-CH_2-CH_2-OR^{2b})_2$ ,  $N(R^{2b})-C(=O)R^{3b}$ ,  $N(R^{2b})-SO_2-R^{3b}$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl-phenyl, -CN and  $-NO_2$ ;

$R^{2b}$  and  $R^{3b}$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $-OR^9$ ,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN and  $-NO_2$ ;

L is selected from:

H, -CN,  $C(=O)NR^{12}R^{13}$ ,  $-(CH_2)_nNR^{12}R^{13}$ ,  $C(=NR^{12})NR^{12}R^{13}$ ,  $OR^{12}$ ,  $-NR^{12}C(=NR^{12})NR^{12}R^{13}$ , and  $NR^{12}C(=NR^{12})-R^{13}$ ;

n is an integer from 0 to 8;

$R^{12}$  and  $R^{13}$  are independently selected from:

hydrogen,  $-OR^{14}$ ,  $-NR^{14}R^{15}$ ,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $COOC_{1-4}$ alkyl,  $COO-C_{0-4}$ alkylphenyl and  $COO-C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH,  $-O-C_{1-4}$ alkyl,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN, and  $-NO_2$ ;

$R^{14}$  and  $R^{15}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on

the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

2. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 1.
3. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 1.
4. The method of claim 3, wherein the condition is selected from the group consisting of:  
acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.
5. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 1.

6. A compound of claim 1,

wherein:

A is selected from:

- (a) C<sub>1</sub>-C<sub>6</sub>-alkyl;
- (b) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;
- (c) phenyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;
- (d) naphthyl, which is independently substituted with 0-2 R<sup>1</sup> substituents; and
- (e) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from the group consisting of N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1</sup> substituents;

R<sup>1</sup> is selected from:

halo, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, -C<sub>1</sub>-C<sub>4</sub>-alkyl, -CN, -C<sub>1-4</sub>alkyl, -C<sub>2-6</sub>alkenyl, -C<sub>2-6</sub>alkynyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl and -NO<sub>2</sub>;

R<sup>2</sup> and R<sup>3</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

m is an integer of 0-2;

Q is a member selected from the group consisting of:



a direct link,  $\text{-C(=O)-}$ ,  $\text{-N(R}^4\text{)-}$ ,  $\text{-C(=O)-N(R}^4\text{)-}$ ,  $\text{-N(R}^4\text{)-C(=O)-}$ ,  $\text{-SO}_2\text{-}$ ,  $\text{-O-}$ ,  $\text{-SO}_2\text{-N(R}^4\text{)-}$  and  $\text{-N(R}^4\text{)-SO}_2\text{-}$ ;

$\text{R}^4$  is selected from:

H,  $\text{C}_{1-4}\text{alkyl}$ ,  $\text{C}_{2-6}\text{alkenyl}$ ,  $\text{C}_{2-6}\text{alkynyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{0-4}\text{alkylC}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{0-4}\text{alkylphenyl}$  and  $\text{C}_{0-4}\text{alkylnaphthyl}$ , wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $\text{C}_{1-4}\text{alkyl}$ ,  $\text{C}_{2-6}\text{alkenyl}$ ,  $\text{C}_{2-6}\text{alkynyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{0-4}\text{alkylC}_{3-8}\text{cycloalkyl}$ ,  $\text{-CN}$ , and  $\text{-NO}_2$ ;

D is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $\text{R}^{1a}$  substituents; and
- (b) an aromatic six-membered heterocyclic ring having from 1-2 ring nitrogen atoms, and wherein the ring atoms may be substituted with 0-2  $\text{R}^{1a}$  substituents;


$\text{R}^{1a}$  is selected from:

halo,  $\text{C}_{1-4}\text{alkyl}$ ,  $\text{C}_{2-6}\text{alkenyl}$ ,  $\text{C}_{2-6}\text{alkynyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{0-4}\text{alkylC}_{3-8}\text{cycloalkyl}$ ,  $\text{-CN}$ ,  $\text{-NO}_2$ ,  $(\text{CH}_2)_m\text{NR}^{2a}\text{R}^{3a}$ ,  $\text{SO}_2\text{NR}^{2a}\text{R}^{3a}$ ,  $\text{SO}_2\text{R}^{2a}$ ,  $\text{CF}_3$ ,  $\text{OR}^{2a}$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $\text{C}_{1-4}\text{alkyl}$ ,  $\text{C}_{2-6}\text{alkenyl}$ ,  $\text{C}_{2-6}\text{alkynyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{0-4}\text{alkylC}_{3-8}\text{cycloalkyl}$ ,  $\text{-CN}$  and  $\text{-NO}_2$ ;

$\text{R}^{2a}$  and  $\text{R}^{3a}$  are independently selected from the group consisting of:

H,  $\text{C}_{1-4}\text{alkyl}$ ,  $\text{C}_{2-6}\text{alkenyl}$ ,  $\text{C}_{2-6}\text{alkynyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{0-4}\text{alkylC}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{0-4}\text{alkylphenyl}$  and  $\text{C}_{0-4}\text{alkylnaphthyl}$ , wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $\text{C}_{1-4}\text{alkyl}$ ,  $\text{C}_{2-6}\text{alkenyl}$ ,  $\text{C}_{2-6}\text{alkynyl}$ ,  $\text{C}_{3-8}\text{cycloalkyl}$ ,  $\text{C}_{0-4}\text{alkylC}_{3-8}\text{cycloalkyl}$ ,  $\text{-CN}$  and  $\text{-NO}_2$ ;

M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



and the like, wherein the aromatic carbocyclic ring corresponding to the D portion for each of the bicyclic rings may be replaced with an aromatic heterocyclic ring as defined above for D, and wherein 0 to 2 of the hydrogen atoms on the D portion of the bicyclic ring may be replaced by  $R^{1a}$  substituents as defined above;

$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl, -CN,  $-NO_2$ ,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$ ,  $SO_2R^2$ ,  $CF_3$ ,  $OR^2$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo,  $C_1$ - $C_4$ -alkyl, -CN,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl $C_{3-8}$ cycloalkyl and  $-NO_2$ ;

$R^{17b}$  and  $R^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy,  $-C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $-C_{2-6}alkynyl$ ,  $-C_{3-8}cycloalkyl$ ,  $-C_{0-4}alkyl-C_{3-8}cycloalkyl$ , -CN,  $-NO_2$ ,  $-(CH_2)_mNR^2R^3$ ,  $-SO_2NR^2R^3$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the cycloalkyl, the phenyl ring, or the aromatic heterocyclic ring may be independently replaced with a member selected from the group consisting of halo,  $C_1-C_4-alkyl$ , -CN,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylC_{3-8}cycloalkyl$  and  $-NO_2$ ;

E is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-(-C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-)$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylC_{3-8}cycloalkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $C_{0-4}alkylheteroaryl$ ,  $C_{1-4}alkylCOOH$  and  $C_{1-4}alkylCOOC_{1-4}alkyl$ , wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylC_{3-8}cycloalkyl$ , -OH,  $-O-C_{1-4}alkyl$ , -SH,  $-S-C_{1-4}alkyl$ , -CN and  $-NO_2$ ;

G is selected from:

a direct link,  $-C(R^7, R^8)-$ ,  $-C(R^{7a}, R^{8a})-C(R^{7b}, R^{8b})-$  and  $-C(R^{7c})=C(R^{8c})-$ ;

wherein  $R^7$ ,  $R^8$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{7b}$ ,  $R^{8b}$ ,  $R^{7c}$  and  $R^{8c}$  are independently a member selected from the group consisting of:

hydrogen, halogen,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $-OR^9$ ,  $-N(R^9R^{10})$ ,  $-C_{0-4}alkylCOOR^9$ ,  $-C_{0-4}alkylC(=O)NR^9R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9-CH_2-CH_2-O-R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9(-CH_2-CH_2-O-R^{10})_2$ ,  $-N(R^9)COR^{10}$ ,  $-N(R^9)C(=O)R^{10}$ ,  $-N(R^9)SO_2R^{10}$ , and a naturally occurring or synthetic amino acid side chain, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $-OR^9$ ,  $C_{1-4}alkyl$ ,  $C_{2-6}alkenyl$ ,  $C_{2-6}alkynyl$ ,  $C_{3-8}cycloalkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ , -CN and  $-NO_2$ ;

$R^9$  and  $R^{10}$  are independently selected from:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl, -CN and -NO<sub>2</sub>, and wherein  $R^9$  and  $R^{10}$  taken together can form a 5-8 membered heterocyclic ring;

J is a member selected from the group consisting of:

a direct link, -O-, -O-C( $R^{11}$ ,  $R^{11a}$ )-, -S-, -S(=O)-, -S(=O)<sub>2</sub>-, -S-C( $R^{11}$ ,  $R^{11a}$ )-, -S(=O)-C( $R^{11}$ ,  $R^{11a}$ )-, -S(=O)<sub>2</sub>-C( $R^{11}$ ,  $R^{11a}$ )-, -C(=O)-, -C(=O)-N( $R^{11b}$ )-, -N( $R^{11b}$ )-C(=O)-, -N( $R^{11b}$ )-, -N( $R^{11b}$ )-C( $R^{11}$ ,  $R^{11a}$ )- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{11c}$  substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$  and  $R^{11c}$  are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O- $C_{1-4}$ alkyl, -O- $C_{2-6}$ alkenyl, -O- $C_{2-6}$ alkynyl, -O- $C_{3-8}$ cycloalkyl, -COOH, -C(=O)-O- $C_{1-4}$ alkyl, -C(=O)-O- $C_{2-6}$ alkenyl, -C(=O)-O- $C_{2-6}$ alkynyl, -C(=O)-O- $C_{3-8}$ cycloalkyl,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkylphenyl,  $C_{0-4}$ alkylnaphthyl,  $C_{0-4}$ alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2  $R^{1b}$  substituents;
- (b) naphthyl, which is independently substituted with 0-2  $R^{1b}$  substituents; and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{1b}$  substituents;

$R^{1b}$  is a member selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl,  $C_{0-4}$ alkyl- $C_{3-8}$ cycloalkyl, -CN, -NO<sub>2</sub>,  $NR^{2b}R^{3b}$ ,  $SO_2NR^{2b}R^{3b}$ ,  $SO_2R^{2b}$ , CF<sub>3</sub>,  $OR^{2b}$ , O-CH<sub>2</sub>-CH<sub>2</sub>- $OR^{2b}$ , O-CH<sub>2</sub>-COOR<sup>2b</sup>, N( $R^{2b}$ )-CH<sub>2</sub>-CH<sub>2</sub>- $OR^{2b}$ , N(-CH<sub>2</sub>-CH<sub>2</sub>- $OR^{2b}$ )<sub>2</sub>, N( $R^{2b}$ )-C(=O)R<sup>3b</sup>, N( $R^{2b}$ )-

SO<sub>2</sub>-R<sup>3b</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>2b</sup> and R<sup>3b</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

L is selected from:

H, -CN, C(=O)NR<sup>12</sup>R<sup>13</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup>, C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, OR<sup>12</sup>, -NR<sup>12</sup>C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, and NR<sup>12</sup>C(=NR<sup>12</sup>)-R<sup>13</sup>;

n is an integer from 0 to 8;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

hydrogen, -OR<sup>14</sup>, -NR<sup>14</sup>R<sup>15</sup>, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, COOC<sub>1-4</sub>alkyl, COO-C<sub>0-4</sub>alkylphenyl and COO-C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH, -O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

R<sup>14</sup> and R<sup>15</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl, wherein from 1-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.



7. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 6.
8. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 6.
9. The method of claim 8, wherein the condition is selected from the group consisting of:  
acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.
10. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 6.

11. A compound of claim 1,

wherein:

A is selected from:

- (a) C<sub>1</sub>-C<sub>6</sub>-alkyl;
- (b) C<sub>3</sub>-C<sub>8</sub>-cycloalkyl;
- (c) phenyl, which is independently substituted with 0-2 R<sup>1</sup> substituents;
- (d) naphthyl, which is independently substituted with 0-2 R<sup>1</sup> substituents; and
- (e) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1</sup> substituents;

R<sup>1</sup> is selected from:

halo, C<sub>1-4</sub>alkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>R<sup>2</sup>, CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

R<sup>2</sup> and R<sup>3</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

m is an integer of 0-2;

B is a member selected from the group consisting of:

a direct link, -C(=O)-, -N(R<sup>4</sup>)-, -C(=O)-N(R<sup>4</sup>)-, -N(R<sup>4</sup>)-C(=O)-, -SO<sub>2</sub>-, -O-, -SO<sub>2</sub>-N(R<sup>4</sup>)- and -N(R<sup>4</sup>)-SO<sub>2</sub>-;

R<sup>4</sup> is selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

D is phenyl, which is independently substituted with 0-2 R<sup>1a</sup> substituents;


R<sup>1a</sup> is selected from:

halo,  $C_{1-4}$ alkyl,  $-CN$ ,  $-NO_2$ ,  $-(CH_2)_mNR^{2a}R^{3a}$ ,  $-SO_2NR^{2a}R^{3a}$ ,  $-SO_2R^{2a}$ ,  $CF_3$ ,  $-OR^{2a}$ , and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

$R^{2a}$  and  $R^{3a}$  are independently selected from the group consisting of:

H,  $C_{1-4}$ alkyl,  $C_{0-4}$ alkylphenyl and  $C_{0-4}$ alkylnaphthyl;

M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



wherein 0 to 2 of the hydrogen atoms on the D portion of the bicyclic ring may be replaced by  $R^{1a}$  substituents as defined above;

$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:



halo, C<sub>1-4</sub>alkyl, -CN, -NO<sub>2</sub>, (CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, SO<sub>2</sub>R<sup>2</sup>, CF<sub>3</sub>, OR<sup>2</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S;

R<sup>17b</sup> and R<sup>17c</sup> are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy, -C<sub>1-4</sub>alkyl, -CN, -NO<sub>2</sub>, -(CH<sub>2</sub>)<sub>m</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>NR<sup>2</sup>R<sup>3</sup>, -SO<sub>2</sub>R<sup>2</sup>, -CF<sub>3</sub>, -OR<sup>2</sup>, phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-4 heteroatoms selected from N, O and S;

m is an integer from 0-6;

E is a member selected from the group consisting of:

a direct link, -C(=O)-, -C(=O)-N(R<sup>5</sup>)-, -C(-R<sup>5a</sup>, -R<sup>6a</sup>)- and -(-C(-R<sup>5b</sup>, -R<sup>6b</sup>)-C(-R<sup>5c</sup>, -R<sup>6c</sup>)-;

wherein R<sup>5</sup>, R<sup>5a</sup>, R<sup>6a</sup>, R<sup>5b</sup>, R<sup>6b</sup>, R<sup>5c</sup> and R<sup>6c</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, C<sub>0-4</sub>alkylheteroaryl, C<sub>1-4</sub>alkylCOOH and C<sub>1-4</sub>alkylCOOC<sub>1-4</sub>alkyl, wherein from 0-2 hydrogen atoms on the ring atoms of the phenyl, naphthyl and heteroaryl moieties may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, -OH, -O-C<sub>1-4</sub>alkyl, -SH, -S-C<sub>1-4</sub>alkyl, -CN and -NO<sub>2</sub>;

G is selected from:

a direct link, -C(R<sup>7</sup>, R<sup>8</sup>)-, -C(R<sup>7a</sup>, R<sup>8a</sup>)-C(R<sup>7b</sup>, R<sup>8b</sup>)- and -C(R<sup>7c</sup>)=C(R<sup>8c</sup>)-;

wherein R<sup>7</sup>, R<sup>8</sup>, R<sup>7a</sup>, R<sup>8a</sup>, R<sup>7b</sup>, R<sup>8b</sup>, R<sup>7c</sup> and R<sup>8c</sup> are independently a member selected from from the group consisting of:

hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)-<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, and a naturally occurring or synthetic amino acid side chain;

R<sup>9</sup> and R<sup>10</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

J is a member selected from the group consisting of:

a direct link, -O-, -O-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S-, -S(=O)<sub>2</sub>-, -S-C(-R<sup>11</sup>, -R<sup>11a</sup>)-, -S(=O)<sub>2</sub>-(-R<sup>11</sup>, -R<sup>11a</sup>)-, -C(=O)-N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(-R<sup>11</sup>, -R<sup>11a</sup>)- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>11c</sup> substituents;

R<sup>11</sup>, R<sup>11a</sup>, R<sup>11b</sup> and R<sup>11c</sup> are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>3-8</sub>cycloalkyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, and a C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents;
- (b) naphthyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents; and
- (c) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>1b</sup> substituents;

R<sup>1b</sup> is a member selected from the group consisting of:

halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, -NO<sub>2</sub>, NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>NR<sup>2b</sup>R<sup>3b</sup>, SO<sub>2</sub>R<sup>2b</sup>, CF<sub>3</sub>, OR<sup>2b</sup>, O-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, O-CH<sub>2</sub>-COOR<sup>2b</sup>, N(R<sup>2b</sup>)-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>, N(-CH<sub>2</sub>-CH<sub>2</sub>-OR<sup>2b</sup>)<sub>2</sub>, N(R<sup>2b</sup>)-C(=O)R<sup>3b</sup>, N(R<sup>2b</sup>)-SO<sub>2</sub>-R<sup>3b</sup>, and a 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from N, O and S, wherein from 1-4 hydrogen atoms on the aromatic heterocyclic system may be independently replaced with a member selected from the group consisting of halo, C<sub>1-4</sub>alkyl, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN and -NO<sub>2</sub>;

R<sup>2b</sup> and R<sup>3b</sup> are independently selected from the group consisting of:

H, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

L is selected from:

H, -CN, C(=O)NR<sup>12</sup>R<sup>13</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup>, C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, OR<sup>12</sup>, -NR<sup>12</sup>C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>, and NR<sup>12</sup>C(=NR<sup>12</sup>)-R<sup>13</sup>;

n is an integer from 0 to 6;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

hydrogen, -OR<sup>14</sup>, -NR<sup>14</sup>R<sup>15</sup>, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, COOC<sub>1-4</sub>alkyl, COO-C<sub>0-4</sub>alkylphenyl and COO-C<sub>0-4</sub>alkylnaphthyl, wherein from 0-4 hydrogen atoms on the ring atoms of the phenyl and naphthyl moieties may be independently replaced with a member selected from the group consisting of halo, -OH, -O-C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, -CN, and -NO<sub>2</sub>;

R<sup>14</sup> and R<sup>15</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylC<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

12. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 11.
13. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 11.
14. The method of claim 13, wherein the condition is selected from the group consisting of:

acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.

15. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 11.

16. A compound of claim 1,

wherein:

A is selected from:

- (a) phenyl, which is independently substituted with 0-2  $R^1$  substituents; and
- (b) a monocyclic or fused bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^1$  substituents;

$R^1$  is selected from:

halo,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$  and  $SO_2R^2$ ;

$R^2$  and  $R^3$  are independently selected from the group consisting of:

H and  $C_{1-4}$ alkyl;

m is an integer of 0-2;

Q is a member selected from the group consisting of:


a direct link,  $-C(=O)-$ ,  $-SO_2-$ , and  $-O-$ ;

D is phenyl, which is independently substituted with 0-2  $R^{1a}$  substituents;

$R^{1a}$  is selected from:

halo and  $C_{1-4}$ alkyl;

M, D and N collectively form a bicyclic ring structure selected from the group consisting of:



$R^{16}$ ,  $R^{17}$ ,  $R^{17a}$ ,  $R^{18}$ ,  $R^{18a}$ ,  $R^{18b}$ ,  $R^{19}$  and  $R^{19a}$  are each independently selected from the group consisting of:

halo,  $C_{1-4}$ alkyl,  $-CN$ ,  $-NO_2$ ,  $(CH_2)_mNR^2R^3$ ,  $SO_2NR^2R^3$ ,  $SO_2R^2$ ,  $CF_3$  and  $OR^2$ ;

$R^{17b}$  and  $R^{17c}$  are each independently a member selected from the group consisting of:

hydrogen, -halo, hydroxy,  $-C_{1-4}alkyl$ ,  $-CN$ ,  $-NO_2$ ,  $-(CH_2)_mNR^2R^3$ ,  $-SO_2NR^2R^3$ ,  $-SO_2R^2$ ,  $-CF_3$ ,  $-OR^2$ , phenyl, and a 5-6 membered aromatic heterocyclic ring containing from 1-3 N atoms;

E is a member selected from the group consisting of:

a direct link,  $-C(=O)-$ ,  $-C(=O)-N(R^5)-$ ,  $-C(-R^{5a}, -R^{6a})-$  and  $-(-C(-R^{5b}, -R^{6b})-C(-R^{5c}, -R^{6c})-)$ ;

wherein  $R^5$ ,  $R^{5a}$ ,  $R^{6a}$ ,  $R^{5b}$ ,  $R^{6b}$ ,  $R^{5c}$  and  $R^{6c}$  are independently selected from:

H,  $C_{1-4}alkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $C_{0-4}alkylheteroaryl$ ,  $C_{1-4}alkylCOOH$  and  $C_{1-4}alkylCOOC_{1-4}alkyl$ ;

G is selected from:

a direct link,  $-C(R^7, R^8)-$ ,  $-C(R^{7a}, R^{8a})-C(R^{7b}, R^{8b})-$  and  $-C(R^{7c})=C(R^{8c})-$ ;

wherein  $R^7$ ,  $R^8$ ,  $R^{7a}$ ,  $R^{8a}$ ,  $R^{7b}$ ,  $R^{8b}$ ,  $R^{7c}$  and  $R^{8c}$  are independently a member selected from the group consisting of:

hydrogen, halogen,  $C_{1-4}alkyl$ ,  $C_{0-4}alkyl-C_{3-8}cycloalkyl$ ,  $C_{0-4}alkylphenyl$ ,  $C_{0-4}alkylnaphthyl$ ,  $-OR^9$ ,  $-N(R^9R^{10})$ ,  $-C_{0-4}alkylCOOR^9$ ,  $-C_{0-4}alkylC(=O)NR^9R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9-CH_2-CH_2-O-R^{10}$ ,  $-C_{0-4}alkylC(=O)NR^9(-CH_2-CH_2-O-R^{10})_2$ ,  $-N(R^9)COR^{10}$ ,  $-N(R^9)C(=O)R^{10}$ ,  $-N(R^9)SO_2R^{10}$ , and a naturally occurring or synthetic amino acid side chain;

$R^9$  and  $R^{10}$  are independently selected from:

H,  $C_{1-4}alkyl$ ,  $C_{0-4}alkylphenyl$  and  $C_{0-4}alkylnaphthyl$ ;

J is a member selected from the group consisting of:

a direct link,  $-O-$ ,  $-S-$ ,  $-C(=O)-N(R^{11b})-$ ,  $-N(R^{11b})-$ ,  $-N(R^{11b})-C(-R^{11}, -R^{11a})-$  and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2  $R^{11c}$  substituents;

$R^{11}$ ,  $R^{11a}$ ,  $R^{11b}$  and  $R^{11c}$  are a member independently selected from the group consisting of:



hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -O-C<sub>3-8</sub>cycloalkyl, -COOH, -C(=O)-O-C<sub>1-4</sub>alkyl, -C(=O)-O-C<sub>3-8</sub>cycloalkyl, C<sub>1-4</sub>alkyl, C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, and a C<sub>0-4</sub>alkylheterocyclic ring having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S, CH<sub>2</sub>COOC<sub>1-4</sub>alkyl, CH<sub>2</sub>COOC<sub>1-4</sub>alkylphenyl and CH<sub>2</sub>COOC<sub>1-4</sub>alkylnaphthyl;

Y is a member selected from the group consisting of:

- (a) phenyl, which is independently substituted with 0-2 R<sup>1b</sup> substituents;
- (b) an aromatic heterocyclic ring having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring may be substituted with 0-2 R<sup>1b</sup> substituents;
- (c) a fused aromatic bicyclic heterocyclic ring system having from 5 to 10 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the bicyclic ring system may be substituted with 0-2 R<sup>1b</sup> substituents;

R<sup>1b</sup> is a member selected from the group consisting of:

halo, -C<sub>1-4</sub>alkyl, -OH, -OBn, -O-CH<sub>2</sub>-CH<sub>2</sub>-OH, -O-CH<sub>2</sub>-CH<sub>2</sub>-OCH<sub>3</sub>, -O-CH<sub>2</sub>-COOH, -O-CH<sub>2</sub>-C(=O)-O-CH<sub>3</sub>, -NH<sub>2</sub>, -NH-CH<sub>2</sub>-CH<sub>2</sub>-O-CH<sub>3</sub>, -NH-C(=O)-O-CH<sub>3</sub> and -NH-SO<sub>2</sub>-CH<sub>3</sub>;

L is selected from:

H, -C(=O)NR<sup>12</sup>R<sup>13</sup>, -(CH<sub>2</sub>)<sub>n</sub>NR<sup>12</sup>R<sup>13</sup> and -C(=NR<sup>12</sup>)NR<sup>12</sup>R<sup>13</sup>;

n is an integer from 0 to 6;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:


hydrogen and C<sub>1-4</sub>alkyl;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

17. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 16.
18. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 16.
19. The method of claim 18, wherein the condition is selected from the group consisting of:  
acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.
20. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 16.



21. A compound of claim 1, wherein A is a member selected from the group consisting of:





hydrogen, halogen, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkyl-C<sub>3-8</sub>cycloalkyl, C<sub>0-4</sub>alkylphenyl, C<sub>0-4</sub>alkylnaphthyl, -OR<sup>9</sup>, -N(R<sup>9</sup>R<sup>10</sup>), -C<sub>0-4</sub>alkylCOOR<sup>9</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>, -C<sub>0-4</sub>alkylC(=O)NR<sup>9</sup>(-CH<sub>2</sub>-CH<sub>2</sub>-O-R<sup>10</sup>)-<sub>2</sub>, -N(R<sup>9</sup>)COR<sup>10</sup>, -N(R<sup>9</sup>)C(=O)R<sup>10</sup>, -N(R<sup>9</sup>)SO<sub>2</sub>R<sup>10</sup>, and a naturally occurring or synthetic amino acid side chain;

R<sup>9</sup> and R<sup>10</sup> are independently selected from:

H, C<sub>1-4</sub>alkyl, C<sub>0-4</sub>alkylphenyl and C<sub>0-4</sub>alkylnaphthyl;


J is a member selected from the group consisting of:

a direct link, -O-, -S-, -C(=O)-N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-, -N(R<sup>11b</sup>)-C(-R<sup>11</sup>, -R<sup>11a</sup>)- and a monocyclic aromatic or non-aromatic heterocyclic ring having from 5 to 8 ring atoms, wherein 1-4 ring atoms of the ring system are selected from N, O and S, and wherein the ring system may be substituted with 0-2 R<sup>11c</sup> substituents;

R<sup>11</sup>, R<sup>11a</sup>, R<sup>11b</sup> and R<sup>11c</sup> are a member independently selected from the group consisting of:

hydrogen, halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -C<sub>1-4</sub>alkyl, -COOH, phenyl, and benzyl wherein the aromatic ring of the phenyl or benzyl is substituted with 0-2 members independently selected from the group consisting of halo, -CN, -NO<sub>2</sub>, -OH, -O-C<sub>1-4</sub>alkyl, -C<sub>1-4</sub>alkyl, -COOH and -C(=O)-O-C<sub>1-4</sub>alkyl;


Y and L taken together are a member selected from the group consisting of:




and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

22. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a therapeutically acceptable carrier and a therapeutically effective amount of a compound of claim 16.
23. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of claim 16.
24. The method of claim 18, wherein the condition is selected from the group consisting of:  
acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.
25. A method for inhibiting the coagulation of biological samples, comprising the administration of a compound of claim 16.


26. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{11b}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^7$  is selected from the group consisting of:


H; Me; Et; Ph;



27. A compound of claim 1, having the formula:




wherein the following portion of said formula






is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{11c1}$  is selected from the group consisting of:

H; Me; Et; Ph; OMe; CF<sub>3</sub>; F; OH; Br; NH<sub>2</sub>; SO<sub>2</sub>Me; OCH<sub>2</sub>Ph; OCH<sub>2</sub>CH<sub>2</sub>OMe;




and wherein  $R^{11c2}$  is selected from the group consisting of:


H; Me; Et; Cl; Br; F; Ph;




28. A compound of claim 1, having the formula:



wherein the following portion of said formula




is selected from the group consisting of:






wherein  $R^{11c1}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^{11c2}$  is selected from the group consisting of:


H; Me; Et; Ph;




29. A compound of claim 1, having the formula:




wherein the following portion of said formula:



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:






wherein  $R^{1c1}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^{1c2}$  is selected from the group consisting of:


H; Me; Et; Ph;




30. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{7c}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^{8c}$  is selected from the group consisting of:


H; Me; Et; Ph;




31. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{7a}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^{7b}$  is selected from the group consisting of:


H; Me; Et; Ph;




32. A compound of claim 1, having the formula:



wherein the following portion of said formula




is selected from the group consisting of:






wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{7a}$  is selected from the group consisting of:

H, Me, Ph, OMe, F, OH, Br,  $NH_2$ ,  $OCH_2Ph$ ,  $OCH_2CH_2OMe$ ,




and wherein  $R^{7b}$  is selected from the group consisting of:


H; Me; Et; Ph;




33. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{6c}$  is selected from the group consisting of:

H; Me; Et; Ph; OMe; F; OH; Br; NH<sub>2</sub>; SO<sub>2</sub>Me; OCH<sub>2</sub>Ph; OCH<sub>2</sub>CH<sub>2</sub>OMe;




and wherein  $R^{11b}$  is selected from the group consisting of:


H; Me; Et; Cl; Br; F; Ph;




34. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{11c1}$  is selected from the group consisting of:

H; Me; Et; Ph; OMe; F; OH; Br;  $NH_2$ ;  $SO_2Me$ ;  $OCH_2Ph$ ;  $OCH_2CH_2OMe$ ;




and wherein  $R^{11c2}$  is selected from the group consisting of:


H; Me; Et; Cl; Br; F; Ph;



35. A compound of claim 1, having the formula:




wherein the following portion of said formula






is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{11c1}$  is selected from the group consisting of:

H; Me; Et; Ph; OMe; F; OH; Br; NH<sub>2</sub>; SO<sub>2</sub>Me; OCH<sub>2</sub>Ph; OCH<sub>2</sub>CH<sub>2</sub>OMe;




and wherein  $R^{11c2}$  is selected from the group consisting of:


H; Me; Et; Cl; Br; F; Ph;




36. A compound of claim 1, having the formula:




wherein the following portion of said formula



is selected from the group consisting of:




wherein Y and L, taken together, are selected from the group consisting of:




wherein  $R^{1c1}$  is selected from the group consisting of:

H; Me; Et; Ph; OMe; F; OH; Br; NH<sub>2</sub>; SO<sub>2</sub>Me; OCH<sub>2</sub>Ph; OCH<sub>2</sub>CH<sub>2</sub>OMe;




and wherein  $R^{1c2}$  is selected from the group consisting of:


H; Me; Et; Cl; Br; F; Ph;




37. A compound of claim 1, selected from the group consisting of:




38. A compound of claim 1, selected from the group consisting of:




39. A compound of claim 1, selected from the group consisting of:



40. A compound of claim 1, selected from the group consisting of:




41. A compound of claim 1, selected from the group consisting of:








42. A compound according to the formula:




wherein A—Q— is selected from the group consisting of:


t-Bu ; O-t-Bu ;  $-(CH_2)_{0-5}$ -amino ; OH ; carboxylic acid ester ; carboxamide ;




43. A compound of claim 1, selected from the group consisting of:




44. A compound of claim 1, selected from the group consisting of:




45. A compound of claim 1, selected from the group consisting of:











46. A compound of claim 1, selected from the group consisting of:



47. A compound of claim 1, selected from the group consisting of:




48. A compound of claim 1, selected from the group consisting of:




and

49. A compound of claim 1, selected from the group consisting of:







50. A compound of claim 1, selected from the group consisting of:




51. A compound of claim 1, selected from the group consisting of:




52. A compound, selected from the group consisting of:



wherein Q is a direct link, and A is a member selected from the group:



or Q is a -C(=NH)- group, and A is a member selected from the group:




G is a direct link;

J is a member selected from the group:



Y-L is a member selected from the group:



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/21742

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(7) : C07D 209/04, 215/00, 235/04, 241/36, 265/36, 403/02, 498/02

US CL : 544/105, 354; 546/165; 548/305.4, 364.7, 490

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 544/105, 354; 546/165; 548/305.4, 364.7, 490

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
EAST, Registry Online, CA Online**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 129:330730,	1-36, 52
---	RIES, U. 'Preparation of disubstituted bicyclic heterocyclic antithrombotics and	-----
Y	anticoagulants,' abstract, DE 19718181, 05 November 1998.	37-51
X	Database CA on STN, Chemical Abstracts (Columbus, OH, USA), No. 128:16433,	1-36, 52
---	SUZUKI, N. 'Preparation of thienopyridininones as GNRH agonists and antagonists,'	-----
Y	abstract, WO 97/40846, November 1997.	37-51
X	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 127:135807,	1-36, 52
---	SUZUKI, N. 'Preparation of condensed bicyclic compounds as prolactin production	-----
Y	inhibitors,' abstract, EP 781774, July 1997.	37-51
X	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 127:50652,	1-36, 52
---	FRECHETTE, R. 'Preparation of benzoxazine antimicrobial agents,' abstract, WO	-----
Y	97/17333, May 1997.	37-51
X	Database CA on STN, Chemical Abstracts, (Columbus, OH, USA), No. 126:343504,	1-36, 52
---	FURUYA, S. 'Preparation of quinolin-4-ones as GnRH antagonists,' abstract WO	-----
Y	97/14682, April 1997.	37-51



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:		"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A"	document defining the general state of the art which is not considered to be of particular relevance	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E"	earlier application or patent published on or after the international filing date	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&"	document member of the same patent family
"O"	document referring to an oral disclosure, use, exhibition or other means		
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

06 November 2000 (06.11.2000)

Date of mailing of the international search report

25 JAN 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Dorothea Lawrence  
D. Margaret Seaman

Telephone No. 703-308-1235

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/21742

## C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus. OH, USA), No 125:167737, KATRITZKY, A.R. 'A new indole synthesis via [3+3] annulation of 2-(benzotriazol-1-ylmethyl)pyrroles with .alpha.,.beta.-unsaturated aldehydes and ketones,' abstract Tetrahedron Lett, Vol.37, No. 32, pages 5641-5644, 1996.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus. OH, USA), No. 124:317157, SUMMERS, J.B. 'Platelet activating factor antagonists: imidazopyridine indoles,' abstract, US patent #5.486,525, January 1996.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus. OH, USA), No. 124:175907, SHEPPARD, G.S. 'Synthesis and evaluation of water soluble indole pyrrolothiazole PAF antagonists,' abstract, Bioorg. Med. Chem. Lett. Vol. 5, No. 23, pages 2913-1918, 1995.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus. OH, USA), No. 123:340121, SUMMERS, J.B. 'Preparation of 3-[(imidazopyridylalkyl)benzoyl]indoles and analogs as platelet activating factor antagonists,' abstract, WO 95/16687, June 1995.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus. OH, USA), No. 123:32955, BACH, N.J. 'Preparation of 1H-Indole-3-Acetic acid hydrazides as sPLA2 inhibitors,' abstract, EP 620214, October 1994.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus. OH, USA) No. 122:290852, BOYD, D. B. 'Preparation of arylindoles, benzimidazoles, and -indazoles as angiotensin II angatonists,' abstract, EP 574174, December 1993.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus. OH, USA), No. 119:49384, SUMMERS, J.B. 'Preparation of 7-indol-3-yl carbonyl pyrrolo 1,2-c thiazoles and related compounds as platelet activating factor antagonists,' abstract, WO 93/01813, February 1992.	1-36, 52 ----- 37-51
X --- Y	Database CA on STN, Chemical Abstracts, (Columbus. OH, USA) No. 99:157, KOLASA, K. 'Preliminary pharmacological studies of the central action of phenyl and piperidinomethyl derivatives of 2-benzoxazolone,' abstract, Ann. Univ. mariae Curie-Sklodowska, Sect. D, Vol. 36, pages 73-81, 1981.	1-36, 52 ----- 37-51